Six days after admission his blood count showed 3,890,000 red cells per c.mm.; haemoglobin, 9.5 g. per 100 ml.; white cells 3,100 per c.mm. (46% polymorphs, 50% lymphocytes, 1% basophils, 1% eosinophils); M.C.V., 68 cubic μ ; the film showed hypochromia. The patient had received no iron or other therapy apart from his ascorbic acid.

At this stage he was put on iron and ammonium citrate mixture, 1 g. three times daily, and given totals of 350 mg. of thiamine, 350 mg. of nicotinic acid, and 10 ml. of crude liver extract by injection in the course of 10 days. The red cells increased to 4,600,000 and the haemoglobin to 14.5 g.% on July 3.

His physical condition improved rapidly, and within a fortnight he was up and about and doing odd jobs in the ward. His mental condition unfortunately showed no parallel improvement. He was discharged to the care of a niece, as he was no longer able to look after himself.

COMMENTS

Although mild degrees of nutritional deficiency are frequent enough in old people who live alone, gross deficiency syndromes amounting to clinical scurvy or pellagra are exceedingly rare. Senile purpura is often mistaken for the former, and glossitis originating from dentures or the use of antibiotics combined with peripheral vascular disease can produce a clinical picture which is sometimes mistaken for pellagra in aged persons.

During the past three years we have seen only one other case of scurvy among the aged sick, this patient being a feeble-minded man who lived alone. During the course of two years in Belmont Road Hospital, Liverpool—a hospital which catered for the aged sick of that city—one of us saw four cases of scurvy.

The history in all these cases is much the same. The old person lives alone and never gets a "cooked meal." Among this population, making tea to go with bread-and-butter or frying meat, egg, or fish is not reckoned as cooking a meal. A "cooked meal" includes potatoes and a vegetable.

Sheldon (1948), in his survey of Wolverhampton old people, showed that widows tended to preserve the home and the home routine when the partner died, whereas widowers did not. Elderly women when living alone will cook a meal in the sense described above, whereas elderly men will not. It is not surprising, therefore, that all the scorbutics we have met with in this age group were old men.

There are two institutions in our cities which cater for the needs of lonely old people and help to keep the incidence of scurvy low. For those who can walk there is the local fish-and-chip shop. For those confined to the house there is the mobile meals service. The cooked meal delivered by the local authority three times a week supplies many potential deficiencies, and the potatoes which it always contains are an excellent source of vitamin C. It is frequently reported that these people, rather than cook a meal for themselves, will consume half the meal on the day it is delivered and save half for heating up on the next day. The nutritionist might object to the wastage of vitamin C by this procedure, but in practice it seems that the meal retains sufficient vitamins to supply the needs of the old people.

The case recorded illustrates destitution in the midst of plenty which is occasionally a tragic feature of old age. There was enough money to purchase food, but neither partner was mentally capable of looking after the home. Although some improvement occurred in the patient's mental condition after iron and vitamin C and B complex supplements, there was no cure such as is seen in cases of pellagra. The mental deterioration in both partners obviously antedated the scurvy—it was a cause rather than a consequence of the deficiency.

WILLIAM HUGHES, M.D., M.R.C.P.
DOROTHY C. MACLENNAN, M.B.

Stapleton Hospital, Bristol.

REFERENCE

Sheldon, J. H. (1948). *The Social Medicine of Old Age*. London.

Kwashiorkor Syndrome in a 3-months-old African Infant

Kwashiorkor is believed to be exceedingly rare under the age of 6 months (Trowell and Davies, 1952), and in a study of 1,000 cases recorded in the literature no case was reported under the age of 3 months. Cases in the 4–6 months group were those in which breast-feeding had failed and the infant was fed on starchy foods from birth.

The following case is of interest in that the infant was breast-fed from birth and presented at the age of 3 months with all the signs of advanced kwashiorkor.

CASE REPORT

A female Jalu baby aged 3 months was admitted to hospital on March 5, 1953, weighing 6 lb. (2.7 kg.). The mother gave a history that the child had been born at home one month prematurely (birth weight not known), and although small was apparently healthy. Breast-feeding was established without difficulty and the child thrived. At the age of 1 month she developed oedema of the face and feet and started to have loose stools—an average of four bulky stools being passed in twenty-four hours. The baby continued to feed well, with very occasional vomiting after feeds. There was no history of an infection or other illness during the first three months of life.

On examination she appeared contented and quite mentally alert. There was marked oedema of the face, hands, and feet, but no apparent loss of subcutaneous fat. The skin was pale and depigmented and the hair quite white. Examination of the abdomen showed the liver to be enlarged to three fingerbreadths below the costal margin. No abnormalities were detected clinically in any of the other systems.

Laboratory Investigations.—Blood slide, negative for malaria parasites. Mantoux 1:1,000, negative. Hb, 85%; red blood cells, 3,210,000 per c.mm.; white blood cells, 4,900 per c.mm. (polymorphs 42%, lymphocytes 57%, monocytes 1%). A blood film showed no abnormalities. Urine contained no albumin. Blood urea, 25 mg. per 100 ml. Plasma proteins: total serum protein, 4.02%; albumin 1.73%; α globulin 0.27%; β globulin 1.39%; γ globulin 0.63%. Ide test, negative. Mother's Kahn test, negative. Mother's breast milk: protein, 1.42%; lactose, 7.1%; fat, 2.4%.

Breast-feeding was maintained while the infant was in hospital, as the mother had an adequate supply, an average gain of 3 oz. (85 g.) being recorded after test weighing. An additional supplement of 8 g. of protein in the twenty-four hours was given in the form of dried skimmed milk, which the child took readily.

No clinical improvement occurred, and on March 19 it was noticed that the child had become more oedematous and was disinclined to feed. Tube feeding was instituted for twenty-four hours, with expressed breast milk and dried skimmed milk, after which she seemed better and oral feeding was recommenced. On March 24 the condition deteriorated suddenly. The child became dyspnoeic and was found to have a left-lower-lobe pneumonia. Penicillin was given immediately, but the child died two days later.

At post-mortem examination the clinical finding of a left-lower-lobe pneumonia was confirmed. Examination of the abdomen revealed an enlarged pale liver. Kidneys, spleen, and pancreas were all macroscopically normal. No abnormality was discovered in any other system. The pathological report on sections of liver and pancreas was: "The liver shows generalized fatty change. There is no evidence of cirrhosis. There is some atrophy and fibrosis of the pancreas."

My thanks are due to the Honourable Director of Medical Services, Kenya Colony, for permission to publish and to Mr. Pearson, of the Medical Research Laboratory, Nairobi, for the biochemical investigations.

LORNA G. MACDOUGALL, M.B., Ch.B., D.C.H.,
Colonial Medical Services, Kenya.

REFERENCE

Trowell, H. C., and Davies, J. N. (1952). *British Medical Journal*, 2, 796.