On the day of admission his breathing was rapid for short periods. He passed only small amounts of urine, without pain and with no obvious haematuria.

On admission he was drowsy and fretful, lying curled up when left alone. There was a widespread morbilliform rash, and the face was flushed. Two small bruises were noted in the lumbar region. Many glands were palpable in the neck, and on the right side there was a large soft swelling overlying the middle of the sterno-mastoid muscle and extending forward over the angle of the lower jaw. The axillary and inguinal glands were also palpably enlarged. The liver edge was palpable two fingerbreadths below the costal margin. No other abnormalities were discovered.

A provisional diagnosis of sulphonamide toxicity was made and penicillin administration was begun in a dosage of 1.5 mega units daily. Potassium citrate was also given in a daily dose of 6 g., and the fluid intake was pushed to 2 pints (1,140 ml.) a day. The haematological findings are summarized in the Table.

On the second hospital day the boy's temperature rose to 103° F. (39.4° C.). He was passing urine plentifully—21 oz. (200 ml.) plus two wet beds in the first 24 hours—and drinking freely, but remained very drowsy. The urine contained a trace of albumin, but no red cells or crystals were found; the chemical test for sulphonamides was strongly positive. In the afternoon of this day he passed a very large tarry stool which gave a strongly positive benzidine reaction.

The temperature settled within 12 hours, and in the course of the next few days the child became brighter and more

rested itself by very slight bruising and profuse melaena. The cerebral state was probably also the consequence of sulphonamide intoxication.

Thrombocytopenia after sulphonamide administration has been infrequently reported. In 1943 Kracke and Townsend reviewed the literature and found reports of 10 cases, to which they added two of their own. Since then there have appeared reports of another 21 cases, making 33 in all. The drugs responsible were sulphanilamide, 4 cases; sulphapyridine, 8 cases; sulphathiazole, 12 cases; sulphadiazine, 7 cases; sulphamerazine, 1 case; and sulphafurazole, 1 case. The frequency with which preparations appear on this list reflects their popularity rather than their toxicity, and it is noteworthy that the number of reports, after reaching a peak in 1943, 1944, and 1945, has fallen off sharply—presumably following the more general use of penicillin and other antibiotics. It would appear that almost all of the more soluble and absorbable sulphonamides are capable of producing thrombocytopenia.

In the present case there was an obvious depression of the bone marrow affecting all elements. This was almost certainly due to the sulphonamide in view of the rapid recovery following withdrawal of the drug, the failure to discover any other cause, and the associated symptoms of sulphonamide toxicity. This case underlines the potential toxicity of even the best of the sulphonamides. It would appear to be advisable that these drugs be prescribed in quantities sufficient for a definite short period only, and that opportunities for excessive medication be not given to parents of patients by supplying more than is needed. The over-dosage in this case arose from a misunderstanding, and though the outcome was favourable it might not have been.

I wish to thank Professor W. F. Gaisford for criticism and advice.

REFERENCE

Kracke, R. R., and Townsend, E. W. (1943). *J. Amer. med. Ass.*, **122**, 168.

Haematological Findings

Date	Hb	W.B.C.					Platelets
		Total	Poly-morphs	Lympho-cytes	Eosino-phils	Mono-cytes	
19/1/54	78%	3,980	66%	33%	—	1%	45,000
20/1/54	78%	4,600	48%	51%	—	1%	55,000
21/1/54		5,400	45%	53%	1%	1%	60,000
22/1/54		6,300	38%	60%	1%	1%	110,000
23/1/54		8,700					135,000
25/1/54		12,600					275,000
26/1/54		9,600					360,000
27/1/54	74%	9,000	55%	42%	—	3%	400,000

alert, the rash gradually faded, and the lymph-node swelling subsided. Melaena stools were passed for three days, but on the ninth day after admission the faeces gave a negative benzidine reaction. The urine never contained red cells or crystals, but the chemical test for sulphonamides remained positive until the third day.

A throat swab taken on admission gave a scanty growth of normal upper respiratory flora. A Paul-Bunnell test on the third day gave no agglutination and a 1:1,000 Mantoux test was negative. Chest x-ray films taken on admission showed slight enlargement of the right hilar shadow but no pulmonary lesion. Bone marrow was obtained on the second day in hospital by iliac crest puncture. There was definite hypoplasia—total nucleated cell count of 46,800 per c.mm. (normal for children of 2 to 10 years, 70,000–700,000 per c.mm.)—with general reduction in the number of all types of nucleated cells; myeloid/erythroid ratio 12:1 (normal ratio in children 4:1 to 13:1); megakaryocytes were plentiful—1% of the nucleated cells.

The child was finally discharged after 12 days in hospital in apparently normal health.

Discussion

This child received a total of 30 g. of sulphadimidine in 11 days. He had not, so far as his parents were aware, ever received a sulphonamide before, and there was no family or personal history of allergy or idiosyncrasy. The effects were presumably, therefore, a direct toxic action of the overdose. These effects were fever, rash, lymph-node enlargement, moderate granulocytopenia (total polymorph count of 2,600 per c.mm.), and marked thrombocytopenia. The last mani-

Medical Memoranda

Treponematoses in Gambia: Preliminary Communication

Investigations have been made into the forms of treponematoses occurring in Gambia, and two forms have been encountered. These are yaws and another treponemal disease which shows many similarities with the endemic syphilis described among the Arabs and the people of Bosnia.

This second disease, as found in Gambia, usually appears first on the mucous membranes at the angles of the mouth in young children, or it can appear on the tip of the penis in small boys. These lesions contain large numbers of spirochaetes morphologically identical with *Treponema pallidum*, and the patients give strongly positive Wassermann and Kahn reactions. There is little doubt that the disease is mainly transmitted by the use of a common drinking-cup and, less often, by close personal contact. The use of a common drinking-cup is prevalent in Gambia, and the people themselves, when asked if the disease could be transmitted from one person to another, volunteered the information that they believed the drinking-cup to be responsible for passing on the disease.

The secondary manifestations of the disease which have been observed are (a) florid eruptions on the vulva and in the anal cleft, and (b) circularly arranged groups of papules which may coalesce to form rings; these can occur on any part of the skin. These lesions contain large numbers of spirochaetes.

The most common tertiary lesion observed is a hyperkeratosis of the palms and soles associated at a later stage with depigmentation of the hands and feet. Destructive lesions of the nasal septum are fairly common, but lesions of the long bones are rarely seen.

There are five main tribes in Gambia, and all cases of this disease which have been seen have been restricted to one tribe.

Gambia occupies a unique position in West Africa, as it lies on the fringe of the tropical rain-belt. The average annual rainfall in Gambia is 42 in. (107 cm.), with a dry season extending over eight months, while 45 miles (72 km.) to the south the rainfall is 73 in. (185 cm.), and 83 miles (134 km.) to the north the rainfall is 21 in. (53 cm.). The mean temperature in Gambia is 81° F. (27.2° C.). Therefore it is likely that the climatic conditions in Gambia, which are intermediate between those in tropical West Africa to the south and those in the desert to the north, are suited to the transmission of both yaws and the so-called endemic syphilis. This theory tends to be supported by the present observations on the occurrence of the two diseases side by side in a relatively small territory.

Our thanks are due to Dr. A. E. Wilkinson, of the M.R.C. V.D. Reference Laboratory, who undertook the serological investigations.

JAMES A. MCFADZEAN, M.D.,
National Institute for Medical Research, London,
and Medical Research Council Laboratories Gambia.

JOHN F. MCCOURT, M.B., D.P.H.,
Medical Officer of Health, Gambia.

General Peritonitis after Subgluteal Abscess

The following case is reported as no previous reference to such a case can be found in the literature.

CASE REPORT

A man aged 33 was admitted to hospital as an emergency case with a provisional diagnosis of general peritonitis due to a burst appendicular abscess. Twelve days previously he had gone to bed at home with fever and headache. Three days later, in addition to the fever which continued, he developed a pain in the left buttock. He called in his own doctor, who diagnosed influenza, which was prevalent at the time. Five days later his doctor saw him again. He was still feverish and the pain in the left buttock was worse and was thought to be due to sciatica. He was also constipated and was given a strong aperient, which caused a copious evacuation of the bowels two days later. The pain in the buttock continued to get worse, until on the morning of admission it suddenly disappeared. In the afternoon he developed stomach-ache, felt especially in the right iliac fossa and across the hypogastrium under the ribs. There was no vomiting or cough.

His previous medical history was that he had had a pyogenic folliculitis of the hair of the left leg for six months. This had now cleared up. In 1941 he had suffered from severe non-specific dysentery while serving with the Forces in West Africa. This was followed by a poly-arthritis affecting especially the hip-joints. He was invalided out of the Forces, but had been quite well since working as a telephone supervisor.

On examination in hospital he was seen to be well built and well covered. He looked ill and had rapid gasping respiration. His temperature was 103.8° F. (39.9° C.), and pulse 114. His tongue was dry and the abdomen tender all over, but especially in the right iliac fossa. There was guarding of both recti, but no board-like rigidity. Examination per rectum was painful, but no mass could be felt. The left buttock appeared quite normal and was not tender on pressure. The movements of both hip-joints were full and painless.

At operation the same evening the abdomen was opened through a long right paramedian incision. The peritoneal cavity was found to contain a large quantity of frank yellow pus. Large quantities of pus were sucked from the right paracolic gutter and the pelvis. The appendix, duodenum, stomach, gall-bladder, and bowel were examined without revealing any abnormality. The abdomen was therefore

closed with drains into the pelvis and below the right lobe of the liver. He made a good recovery with the aid of chemotherapy and intravenous fluids.

One week after operation he complained of a recurrence of the pain in the left buttock. Twelve days after operation there was a distinct tender swelling under the gluteus maximus between the great trochanter of the left femur and the sacrum. He was given a general anaesthetic and the fibres of the gluteus maximus were split to drain a large abscess cavity. On inserting a finger, the posterior surface of the neck of the femur was directly felt. X-ray examination of the bony pelvis and femurs showed no sign of osteomyelitis. A culture of the pus from the abscess grew anaerobic streptococci and the coliform organisms. One month later, when the incision in the buttock had healed, he developed a pelvic abscess, which was successfully drained per rectum. A culture of the pus grew non-haemolytic streptococci.

Three weeks later there was a recurrence of pain in the left buttock. The abscess was reopened and drained. Further radiographs showed no infective focus of bones. He then made rapid progress and was discharged home one month later, though the buttock incision was still discharging slightly.

He finally returned to work six months after the onset of the illness. Both hip-joints had full painless movements and there was no residual disability of any sort.

COMMENT

It seems probable that the general peritonitis in this case was caused by pus from the subgluteal abscess tracking along the left pyriformis muscle and bursting into the peritoneal cavity through the pelvic peritoneum. This probability seems strengthened by the fact that a pelvic abscess developed six weeks after the initial drainage of the peritoneal cavity, and shortly after the first external drainage of the abscess had ceased. Presumably a second overflow of pus reached the pelvis, where it was walled off before reaching the general peritoneal cavity.

J. F. H. BULMAN, F.R.C.S.,
Carshalton War Memorial Hospital, Surrey.

Torsion of Hydrosalpinx at 14 Years of Age

Torsion of the Fallopian tube, although not a rarity (de Solderhoff, 1949), is still uncommon enough to be worth reporting. Wolf (1951) recorded that just over 120 cases had been described since 1890. Neel *et al.* (1943) saw torsion of the tube in a child aged 3 years. Tauber (1949) undertook a review of the subject of tubal torsion, and stressed that the history of repeated attacks of pain was the most significant feature. Shaw (1949) believed that the condition differed from torsion of an ovarian cyst in that "shock" was absent. Novak (1949), quoting Anspach, stated that a hydrosalpinx in childhood may be secondary to (a) vulvovaginitis, (b) one of the exanthemata, or (c) tuberculous infection.

CASE REPORT

A girl aged 13 years 8 months was admitted to hospital in June, 1952, complaining of colicky pains in the left iliac fossa for 13 hours. She had vomited four times. There was no disturbance of her bowel or urinary functions. She had suffered from most of the common exanthemata. There was no history of previous attacks of abdominal pain. Her mother stated that from infancy until a year previously the child had had repeated episodes of vulvar inflammation, without any discharge. It had always responded to bathing and the application of talcum powder. There was no relevant family history.

On examination the temperature was 99.4° F. (37.4° C.), pulse 92. Tenderness, guarding, and rigidity were present in the left iliac fossa. Rectal examination revealed no