TWO CASES OF FULMINATING PNEUMONIA IN PATIENTS ON HORMONE THERAPY

BY

J. A. PAGE, M.B., B.S., M.R.C.S.

(From the Rheumatism Research Unit, South-west and
Oxford Regions, Royal National Hospital for
Rheumatic Diseases, Bath)

Many articles have appeared in recent months on the uses of A.C.T.H. and cortisone in the treatment of various acute and chronic diseases. Little emphasis has been laid on the practical results of the well-known disseminating qualities of these drugs on occult infections.

Leading articles in the *British Medical Journal* (1954) and *Lancet* (1951) have drawn attention to this disconcerting quality and discussed at length its theoretical causation. In the latter article it was queried: "In treating the pale rheumatic child with some disorder of his antibody responses, shall we gain only a fit-looking child with no defences against infection, who makes his symptomless way to death from septicaemia?"

It is with attention to the word symptomless that I would like to report two cases of unexpected death and discuss their practical implication.

Case 1

A housewife aged 65 was admitted with an acute generalized exacerbation of her rheumatoid arthritis precipitated by purulent rhinorrhoea and sinusitis. On admission she had cardiac failure, pyrexia, generalized swelling, and joint pain, and was unable to sit up in bed by herself. She had lost about 1½ st. (9.5 kg.) in weight in the past three months. Her E.S.R. (Westergren) was 110 mm. in one hour, haemoglobin 52%, and B.P. 160/90. She ran a stormy course of remissions and exacerbations, eventually being discharged to a convalescent home eight months after admission.

During her stay in hospital she was at first given 50 mg. of cortisone daily, combined with a short course of penicillin and digoxin. In two months she had improved to a degree where a reduction in the dose of cortisone to 37.5 mg. daily was reasonable. However, she rapidly had a further exacerbation of her symptoms, which failed to settle on a dose of 75 mg. of cortisone daily. It was decided that the cortisone should be replaced by A.C.T.H. She was started on 60 mg. of A.C.T.H. daily in October, on which she improved quite rapidly, enabling us to reduce the dose to 20 mg. of A.C.T.H. gel daily within a fortnight. Steady improvement was maintained, and therapy was discontinued. Within a week she had a further exacerbation of her rheumatoid arthritis and purulent rhinorrhoea which necessitated another course of penicillin, combined with 100 mg. of cortisone daily. Her symptoms gradually improved, and a reduction to 75 mg. of cortisone daily was possible. It was found, however, that any further reduction in cortisone resulted in an immediate exacerbation. She was discharged in February, 1954, to continue taking 75 mg. of cortisone daily. At this time she was able to walk with the aid of two sticks and had gained 14 lb. (6.4 kg.) in weight.

During her stay in hospital she had three short courses of penicillin, and two courses of intravenous iron and cortisone and A.C.T.H.

She was seen in the out-patient department a month after discharge and appeared very well, having no joint pains and being able to walk unaided. She was sent home to continue taking 75 mg. of cortisone daily. At this time her E.S.R. was 26 mm. and haemoglobin 80%.

Three weeks later she died in her sleep, having been normally active throughout the day and making no complaint whatsoever. Her brother said that a few days previously she had been better than she had been for years, being more active and able to get about her housework with more ease.

Post-mortem Findings.—Macroscopically there was a bilateral hydrothorax of approximately 2–3 pints (1,140–1,700 ml.) with a bilateral early confluent basal bronchopneumonia. The heart musculature was thin and of poor texture. The pericardium was adherent anteriorly. The liver showed passive congestion. The kidneys appeared normal. There were typical rheumatoid arthritic joint changes. The suprarenals were normal (right 5 g., left 5.5 g.). The pituitary was normal (weight 0.9 g.). Histological examination showed that the suprarenals and pituitary were normal for her age. The myocardium showed mild toxic myocarditis. The liver showed cloudy swelling with an increase of red cells in the portal tract. The kidneys were normal. The sternal marrow was of normal cellularity and the thyroid gland showed normal activity, with small lymphoid accumulations. The lung was not examined histologically.

Case 2

An electrical engineer aged 55 was readmitted in early January, 1954, because of gradual deterioration of his rheumatoid arthritis whilst on 400 mg. of phenylbutazone daily, which he had been taking for about a year. He complained of pain in all his joints, especially the right elbow and both shoulders. The E.S.R. (Westergren) was 12 mm. in one hour; haemoglobin, 95%; W.B.C., 13,400 (polymorphs 84%, lymphocytes 9%, monocytes 7%). The same dose of phenylbutazone was continued together with a short course of A.C.T.H., beginning with 20 mg. gel daily, which was reduced to 10 mg. daily twelve days later. Four days after this he complained of difficulty in respiration, due to inspiratory stridor, which failed to respond immediately to subcutaneous adrenaline. He was given 50 mg. of crystalline A.C.T.H. intramuscularly, this being repeated twice in the next forty-eight hours, by which time the attack had abated. He was given a short course of penicillin because his throat swab revealed haemolytic streptococci, not Lancefield group A or O. The E.N.T. opinion queried chronic inflammatory laryngitis secondary to crico-arytenoid arthritis. The phenylbutazone was discontinued and 10 mg. of A.C.T.H. gel daily was continued for a further fourteen days, in which time he greatly improved and his joints were much easier. On withdrawal of A.C.T.H. he had a mild flare-up of his symptoms, which were brought easily under control by regular aspirin. He was discharged in the latter part of February.

In the latter part of April he had an acute flare-up of all his joints and was admitted to a nursing-home, where he was put on 20 mg. of A.C.T.H. gel daily. Seven days later he was symptom-free and feeling so much better that he was planning his return to work. That evening he went to sleep at a normal time but awoke in the early hours of the morning, short of breath, and died within six hours.

Post-mortem Findings.—Macroscopically the left pleural cavity contained dense pleural adhesions along the anterior margin of the lung due to a fibro-purulent exudate from a fairly recent pleurisy; there was some free fluid present. The right pleural cavity was obliterated by dense adhesions, but there was no free fluid. The right lung was oedematous. The left lung showed congestion and oedema of the lower lobe. The upper lobe was totally consolidated and in a state of grey hepatization. The pharynx and larynx were both healthy and normal. There was moderate enlargement of two mediastinal glands. Both the kidneys were congested. The heart muscle appeared to be normal, but there was a finely patent foramen ovale in the interatrial septum. **Histology:**—Lungs: the presence of acute pneumonia was confirmed, a purulent exudate filling the alveoli; the left lung was in a stage of grey hepatization. Heart: slight coronary atheroma; nothing else abnormal. Liver: moderate fatty

change and evidence of chronic venous congestion. Kidney: evidence of toxæmia, with cloudy swelling of tubular epithelium. Spleen: congestion of sinusoids; Malpighian corpuscles were small, with relative deficiency of lymphocytes. Hilar lymph nodes: reactive hyperplasia and evidence of early pyogenic infection. The suprarenals were normal (right 6.6 g., left 6.7 g.). The pituitary was also normal.

Two similar cases are mentioned in the 1953 annual report of the rheumatological department of the Royal Free Hospital. These were both on long-term cortisone; they remained symptomless until they became explosively ill, and in spite of treatment died a few hours later. These cases will be reported fully in due course.

Comment

The two cases reported display the well-known masking qualities of A.C.T.H. and cortisone over severe bacterial infections. In the light of experimental knowledge of dissemination of infection by these drugs, the second case is of interest because of the short duration of A.C.T.H. treatment and also the low dosage of drug given—20 mg. of gel daily. The absence of gross histological change in the adrenal and pituitary glands, using only routine staining methods, is noted.

The practical implication of patients on long-term cortisone is of most concern to the general practitioner, on whom the ultimate supervision of the patient's health depends. Although the incidence of pneumonia in patients on long-term cortisone is relatively low, its occurrence is, in our limited experience, grave and exceedingly difficult to diagnose early enough for treatment to be satisfactory.

The problem of diagnosis can at this stage only rest upon constant clinical examination at relatively short intervals, at the most a week.

The problem of treatment and general management is as yet unresolved, for in the light of present knowledge it would seem justifiable to institute antibiotic therapy at the earliest symptoms or sign of infection.

Summary

Two cases of fulminating pneumonia in patients undergoing hormone therapy for the control of an exacerbation of rheumatoid arthritis are described.

Both cases display the well-known masking effect of cortisone and A.C.T.H. over severe bacterial infection.

In the first case death occurred during prolonged cortisone therapy. In the second case death occurred a week after commencement of A.C.T.H. therapy.

The problem of diagnosis and treatment is discussed.

I am indebted to Dr. G. D. Kersley for his help and advice in preparing this record and for allowing me to report these two cases; also to Dr. H. J. Gibson for furnishing the post-mortem and histological reports.

REFERENCES

- British Medical Journal*, 1954, 1, 381.
Lancet, 1951, 1, 949.

In his Annual Report for 1953 Dr. Kenneth Fraser, county medical officer of Cumberland, describes a survey of all the spastic children in the county. Fourteen of these were being educated at primary schools, three were classified as ineducable, four were in special schools, and four were being considered for admission to special schools. Of the children under 5 years of age, only three were discovered with a serious degree of spasticity. Eight children in all were classified as being seriously disabled owing to spasticity, and three of these had been affected by erythroblastosis foetalis. Dr. Fraser, commenting on the importance of performing blood tests on expectant mothers, adds: "I hope the significance of this will not be overlooked by those concerned."

Medical Memoranda

Intussusception at 70 Hours

I would like to record this case as that of one of the youngest babies to have an intussusception. It is of interest in demonstrating some of the odd symptoms one may encounter when dealing with cases of intestinal obstruction of the newborn.

CASE REPORT

The patient, a male child of healthy parents, was normally delivered. His blood group was A Rh-positive and the W.R. negative. At birth he was quite active and appeared to be normal. There was some bruising of both feet and one hand that was not due to trauma. Thirty hours after birth a dark stool of meconium plus melaena was passed, as were similar stools at 48 hours. At this period he sucked vigorously, but regurgitated green fluid. A tentative diagnosis of haemorrhagic disease of the newborn was made and 10 mg. of "synkavit" was given. At 58 hours he had another melaena stool and vomited some green fluid, and his general condition began to deteriorate.

He was admitted to the Newcastle-upon-Tyne Babies Hospital at 60 hours and was found to have petechial haemorrhages of the trunk, face, hands, and feet. He was observed for two hours, and during this period he vomited bile-stained fluid and his general condition became much worse. He was therefore transferred to the children's department of the Newcastle-upon-Tyne General Hospital as a possible case of duodenal atresia. He was then 66 hours old, weighed 6 lb. (2.7 kg.), and had a haemoglobin of 88%. He was dehydrated and pale. He had small subcutaneous haemorrhages on the face and bruising of the hands and feet. He continued to vomit greenish fluid. The upper abdomen was distended and no mass could be felt.

An x-ray film was taken, and the radiologist reported: "There was no gas distal to the first part of the duodenum, indicating a complete obstruction. Barium enema showed a normal colon and excluded malrotation. The lesion is most likely to be an atresia."

In view of this a provisional diagnosis of duodenal atresia and haemorrhagic disease of the newborn was made, and he was prepared for operation. A blood transfusion was started and gastric aspiration was carried out. At this stage he passed three small stools of fresh blood.

At 77 hours I operated. The abdomen was opened through a right paramedian incision. On opening the peritoneal cavity it was immediately obvious that one was dealing with a high intussusception. Examination showed it to begin about 10 in. (25 cm.) from the duodeno-jejunal junction and to be about 12 in. (30 cm.) long. Reduction was started in the usual manner, but the emerging bowel was seen to be gangrenous and the intussusciptions began to split. I resected the affected bowel, closed both ends, and did a side-to-side anastomosis, using No. 000 catgut throughout. The baby stood the operation well and made an uneventful recovery.

The post-operative regime was as follows. Gastric aspiration was done hourly and the I.V. drip was continued. Twenty-four hours after operation feeds of 2 dr. (7 ml.) of sterile water were given every two hours, aspiration being carried out before the feed. Glucose water was given in gradually increasing amounts for 48 hours until 96 hours after the operation, when the amount being aspirated was less than the amount taken. A stool containing bile was passed at 72 hours. Gastric aspiration was continued at lengthening intervals until seven days after the operation, when only 2 oz. (57 ml.) had been aspirated in the previous 12 hours. He had full-strength half-cream milk from the fifth day and was on to three-hourly feeds after six days. Penicillin and streptomycin were given for ten days.