Published in final edited form as:

Gene Expr Patterns. 2007 October ; 7(8): 837–845.

The Slit receptor *Robo1* is predominantly expressed via the *Dutt1* alternative promoter in pioneer neurons in the embryonic mouse brain and spinal cord

Hikmet Feyza Nural, W. Todd Farmer, and Grant S. Mastick

Departments of Biology and Biochemistry, University of Nevada, Reno NV 89557

Abstract

Robo1 is a member of the Roundabout (*Robo*) family of receptors for the Slit axon guidance cues. In mice (and humans), the *Robo1* locus has alternative promoters producing two transcript isoforms, *Robo1* and *Dutt1*. These isoforms have unique 5' termini, predicted to encode distinct N-terminal amino acids, but share the rest of their 3' exons. To determine the spatial expression of the *Robo1* and *Dutt1* isoforms, we generated isoform-specific RNA probes, and carried out in situ hybridization on E10.5 mouse embryos, the stage in early neuron differentiation when many major axon pathways are established. The two isoforms had distinct expression patterns that partially overlapped. *Dutt1* was the predominant isoform, with widespread expression in regions of post-mitotic neurons and neuroepithelial cells. The *Robo1* isoform had a distinct expression pattern restricted to subsets of neurons, many of which were *Dutt1*-negative. *Dutt1* was the main isoform expressed in spinal cord commissural neurons. For both probes, the main hybridization signal was limited to two spots in the nuclei of individual cells. This study shows distinct expression patterns for the *Dutt1* and *Robo1* alternative promoters in the embryonic nervous system.

Keywords

axon guidance; neuron; Robo; Slit; longitudinal axon; commissural axon; dorsal root ganglion

1. Results and Discussion

Growth cones navigate by selectively responding to molecular cues in their environment. Different types of axons differ in their responses to cues, depending on their expression of specific combinations of guidance receptors. In contrast to the great diversity in neuron types and their axon projection patterns, a surprisingly small number of axon guidance receptors have been identified to date. However, single receptor genes could potentially give rise to multiple receptor isoforms through mechanisms such as alternative promoters and differential mRNA splicing.

The Slit repellents and their Robo receptors are a major guidance system conserved from invertebrates to vertebrates (Dickson and Gilestro, 2006). The Roundabout (*Robo*) family of

Corresponding author: Grant Mastick, Department of Biology, University of Nevada, Reno NV 89557, Phone: 775-784-6168, FAX: 775-784-1650, gmastick@unr.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

transmembrane receptors was first identified in *Drosophila* in a mutant screen for genes that control midline crossing by pioneer axons (Seeger et al., 1993). Slit/Robo signaling has a primary function in keeping axons out of the Slit+ midline, both in commissural axons to prevent re-crossing, as well as in ipsilateral axons to prevent them from crossing at all. Robo homologs have been identified in mammals and other vertebrates (Kidd et al., 1998b; Sundaresan et al., 1998; Yuan et al., 1999). Robo1, Robo2, Robo3 (Rig1) and Robo4 are the four members found in human, rat and mouse. Vertebrate Robo expression is widespread, suggesting that Slit/Robo signaling has broad roles in guiding a diverse set of commissural, longitudinal, and other axon populations. For example, mice with *Robo* mutations have defects in midline crossing of spinal cord commissural axons (Long et al., 2004). Other axon populations dependent on Slit/Robo signaling include major forebrain tracts and commissures (Andrews et al., 2006; Bagri et al., 2002), retinal axons (Plump et al., 2002; Thompson et al., 2006), and motor axons (Hammond et al., 2005), as well as others.

Recent evidence indicates that the small number of vertebrate *Robo* genes each produces more than one isoform. For example, alternative splicing of Robo2 and 3 transcripts results in isoforms with distinct affinity for Slit binding and/or differences in intracellular signaling (Camurri et al., 2004; Yue et al., 2006). Tissue-specific alternative mRNA splicing appears to contribute to the spatial complexity of Robo2 and 3 expression (Challa et al., 2005; Dalkic et al., 2006).

Robo1 presents a different case, with alternative transcription start sites providing the potential to be independently regulated. A human *Robo* homolog was independently cloned from breast and small cell lung cancer cell lines as a potential tumor suppressor gene and designated as *Dutt1* (Deleted in U-twenty twenty) (Sundaresan et al., 1998). Sequence analysis subsequently established that *Dutt1* cDNA sequences largely overlap with *Robo1*, but each has unique 5'-most exons and start codons. Therefore, *Robo1* and *Dutt1* genes are derived from alternative promoters. Interestingly, the *Robo1* and *Dutt1* promoters appear to have differential spatial and temporal transcriptional activity, based on RT-PCR analyses of several mouse tissues including developing brain (Clark et al., 2002). However, whether the *Robo1* and *Dutt1* isoform expression patterns are overlapping or distinct in the embryonic brain has not been determined. Intriguingly, one familial form of dyslexia has been correlated with a translocation breakpoint that causes *Robo1* haplo-insufficiency while leaving *Dutt1* transcription unaffected (Hannula-Jouppi et al., 2005). Together, these results suggest the potential for distinct patterns for each isoform.

To investigate the spatial expression from the two promoters for the *Robo1* and *Dutt1* isoforms, we generated isoform-specific probes for in situ hybridization, and used these to label the mRNA expression patterns of *Robo1* and *Dutt1* in mouse brain and spinal cord on embryonic day 10.5. E10.5 was chosen as a developmental stage in which several key types of brain stem and spinal cord neuron types were differentiating and extending axons, yet early enough to present a relatively simple spatial pattern of identified populations of neurons with diverse axon projection patterns.

1.1. Generation of isoform-specific RNA probes

In the structure of the mammalian *Robo1* locus, the first two exons, referred to as exon 1a and exon 1b, are unique to the *Robo1* isoform, while exon 2 is unique to *Dutt1* (Figure 1A) (Clark et al., 2002). The isoforms share sequences from exon 3, and thus share the coding sequence for the Ig domains. The first 18 amino acids of *Dutt1* correspond to a signal peptide (Figure 1C). *Robo1* extends further N-terminal with a 24 amino acid signal peptide followed by an isoform-specific 32 unique amino acid sequence. Previous studies showed tissue-specific differential expression of mouse *Dutt1/Robo1* isoforms with RT-PCR (Clark et al., 2002).

Different transcription start sites indicate the possibility of promoters driving transcription in separate subsets of cells.

Our first aim was to identify the 5' sequences unique to each mouse isoform. Sequences for the *Dutt1*-specific exon 2 were represented in mouse cDNA and EST databases. The longest exon 2 sequences were 1046 bp long, based on several independent sequences, suggesting that transcription from the *Dutt1* promoter produced an initial 5'UTR of 992 nucleotides and coding sequence of 54 nucleotides. Sequences for the *Robo1* isoform were more difficult to identify. Beginning with cDNA sequences from rat or human that corresponded to *Robo1*-specific exons 1a and 1b, BLAST searches identified conserved mouse genomic sequences for putative exons 1a and 1b. The first 218 bp of the predicted *Robo1* isoform consisted of exon 1a, containing 5' UTR (50 bp) and the beginning of coding sequence (72 bp), plus exon 1b (96 bp). However, no matches to exon 1a and 1b could be identified in mouse cDNA or EST databases. Furthermore, BLAST searches using the shared exon 3 identified 96 ESTs, several with exon 2 sequences at their 5' ends, but none with exon 1a and 1b sequence. Thus, *Dutt1* sequences are represented in the mouse EST databases, but not *Robo1*-specific sequences. One EST, [AW494633](#), matched exon 3 sequence, but with a different sequence at its 5' end that mapped to a BAC-derived genomic sequence 360 kb upstream of exon 2 (lying between the putative exon 1a and 1b). This suggested a third promoter also poorly represented in EST databases, and was not further characterized.

To verify that the *Dutt1* and *Robo1* mRNAs were present in mouse embryos, and to isolate cDNAs sufficiently long for in situ hybridization probes, we used RT-PCR to amplify *Robo1* and *Dutt1* fragments with exon-specific primers from a cDNA pool from whole E10.5 mouse embryos (Figure 1B). Fragments of the predicted sizes were produced for each isoform, 559 bp for *Dutt1* and 200 bp for *Robo1*, suggesting that the predicted exons were in fact transcribed. Reverse primers included T7 minimal promoter sequences to allow synthesis of antisense riboprobes. *Dutt1*^{ex2} indicates a *Dutt1*-specific probe while *Robo1*^{ex1a+1b} corresponds to a *Robo1*-specific probe. For a common probe to recognize both *Robo1* and *Dutt1* mRNA, we used the previously published rat cDNA subclone that spans the sequence encoding the first three Ig domains (Kidd et al., 1998b), and refer to this as the *Robo1*^{ex3} probe.

1.2 *Robo1* and *Dutt1* are differentially expressed in the embryonic brain

To gain an overview of the expression patterns of *Robo1* and *Dutt1* in developing mouse embryos, we performed whole mount in situ hybridization on E10.5 (Figure 2). This stage initiates CNS development with a relatively small number of neuron types, with neuronal cell bodies differentiating in specific regions (Mastick and Easter, 1996). The common probe, *Robo1*^{ex3}, showed a broad expression of combined *Robo1/Dutt1* with heavy labeling in several regions of the brain (Figure 2A). Using isoform-specific probes, our general observation was that *Dutt1* accounted for most of the combined *Robo1/Dutt1* expression (Figure 2B). In contrast, *Robo1* appeared in a restricted subset of the overall expression domain, overlapping with *Dutt1* in some areas but distinct in others (Figure 2C). The specific patterns of expression are explained in more detail below, from anterior to posterior.

In forebrain, *Robo1/Dutt1* expression extended in patches from the base of the optic stalk through ventral thalamus (Figure 2A). A similar expression pattern was observed for the *Dutt1* probe, but fewer cells were labeled with *Robo1* (Figure 2B, C). In the posterior forebrain and anterior midbrain, *Robo1* and *Dutt1* expression partially overlapped (Figure 2B,C). However, *Dutt1* showed broader expression, while *Robo1* was expressed in discrete patches mainly in anterior midbrain. In the hindbrain, *Dutt1* had a very strong lateral band of expression, additional discrete stripes of cells dorsally, and a ventral block of labeling in rhombomere 4. In contrast, *Robo1* was expressed in patches in the anterior hindbrain, with a band of labeling at the level of the motor columns (Figure 2C).

1.3 *Dutt1* has broad expression in neuronal and non-neuronal tissue

To take a closer look at *Robo1* and *Dutt1* expression patterns, we examined brain and spinal cord sections, as described below from anterior to posterior.

In ventral thalamus, both *Dutt1* and *Robo1* could be seen in the mantle layer (Figure 3A, C). This corresponded with neurons, as revealed by co-labeling with β III-tubulin antibody labeling (Figure 3B, D). The initial ventral thalamic neurons are the tyrosine-hydroxylase- and Pax6-positive neurons of the zona incerta that pioneer descending longitudinal fibers through the thalamus (Mastick, et al, 2001; Nural and Mastick, 2005). *Dutt1* was also present in the deeper neuroepithelial layer where there was no *Robo1* labeling (Figure 3A).

Differences in isoform expression were evident in other parts of the brain, as well. In hindbrain, heavy *Dutt1* staining was seen in the mantle layer throughout the dorsal region (Figure 4A), corresponding with β -tubulin labeling (Figure 4B). Towards the ventral half of the hindbrain, the labeling expanded into a block of neuroepithelial labeling (Figure 4A, C). *Robo1* expression was restricted to a small subset within dorsal and ventral neurons. Thus, *Dutt1* and *Robo1* expression could potentially overlap in a subset of dorsal hindbrain neurons, though there were many more *Dutt1*-positive cells. Both *Robo1* and *Dutt1* were expressed ventrally, in the area of the hindbrain motor neurons, where *Dutt1* expression appeared more medial (Figure 4D), and *Robo1* more lateral (Figure 4G). In midbrain sections (not shown), *Dutt1* was expressed by both the neuroepithelium and by the mantle layer of neurons, with *Robo1* limited to a few scattered cells.

Expression in anterior spinal cord was similar to the hindbrain (Figure 5). *Dutt1* expression was present in cell bodies along the margin, medially in the motor columns, as well as in blocks of neuroepithelium (Figure 5A). Again, *Robo1* expression pattern was limited to the marginal layer but in fewer cells compared to *Dutt1* expression, with lateral motor column expression (Figure 5B, F). *Dutt1* also was expressed by dorsal root ganglia (DRG), but not *Robo1*. DRG expression contrasts with a previous study using probes that detected both *Robo1/Dutt1* isoforms, which failed to label E13–17 rat spinal cord DRGs while a *Robo2* probe did (Brose et al., 1999; Wang et al., 1999). The differing results may reflect a higher sensitivity of our probes, or a species difference. The best studied commissural neurons arise in dorsal spinal cord near the roof plate and project their axons ventrally, crossing the midline under Robo guidance (Long, et al., 2004). In dorsal spinal cord neurons, *Dutt1* expression was much more extensive than *Robo1*, indicating that the *Dutt1* isoform is the predominant isoform in commissural axons.

The neuroepithelial labeling in the brain and spinal cord is consistent with previous expression studies of *Robo1* mRNA (Connor and Key, 2002; Hammond et al., 2005; Sundaresan et al., 1998). However, two independent *Robo1* antibodies, neither isoform-specific, have been used to label spinal cord sections, with labeling in post-crossing commissural axons and not in neuroepithelium (Long et al., 2004; Sundaresan et al., 2004). This suggests that *Dutt1* and *Robo1* mRNAs detected by in situ are post-transcriptionally regulated.

1.4 *Robo1* isoform expression in longitudinal neurons

In addition to commissural axons, *Robo1* isoforms would also be expected to be expressed in longitudinal axons. Therefore, we investigated the expression patterns of *Dutt1* and *Robo1* in sections through the clusters of neurons whose axons pioneer two main longitudinal tracts through the E10.5 brain stem. The first tract, the medial longitudinal fasciculus (MLF), is the most ventral pioneer tract, with axons descending from a ventral cluster of neurons (the interstitial nucleus of Cajal) straddling the midbrain/forebrain boundary (Mastick and Easter, 1996; Wilson and Easter, 1991; Wilson and Easter, 1991b). To identify MLF neurons, we

performed retrograde diI labeling of the MLF from the hindbrain back to the cluster of MLF cell bodies, followed by photoconversion to a stable brown label, in situ hybridization, and sectioning. *Robo1* labeling could be seen in many though not all MLF neuron cell bodies (Figure 6B), whereas *Dutt1* was expressed in neuroepithelial cells surrounding the source of MLF, but clearly not in MLF cell bodies (Figure 6A). The second tract examined was the first dorsal longitudinal tract, the tract of postoptic commissure (TPOC), originating from neurons at the base of the optic stalk and projecting through diencephalon and midbrain (Mastick and Easter, 1996). The isolated location of the TPOC neuron cluster allowed us to identify these neurons using β III tubulin antibody. Sections through the source of TPOC showed labeling for both *Dutt1* and *Robo1* (Figure 6B, D). We also examined dorsal midbrain neurons, which include the tmesV neurons that pioneer a dorsal descending tract (Stainier and Gilbert, 1990), but did not detect expression of either *Robo1* isoform in this region (e.g. Figure 2). Thus, *Robo1* and in some cases *Dutt1* are expressed in at least a subset of pioneer longitudinal neurons.

1.5 Nuclear Localization of *Dutt1* and *Robo1* hybridization signals

In our sections, we noticed puncta (dots) with both *Dutt1* and *Robo1* probes (Figure 7A, D), in addition to the expected diffuse labeling within cells. From comparison with DAPI labeling of the cell nuclei, we infer that the puncta were located in the nucleus (Figure 7B, E), with the caveat that the images were collected on a wide-field microscope. Some of the *Dutt1*- and *Robo1*-positive cells were also positive for β -tubulin. Thus, the same patterns of dots were observed in neurons and non-neuronal cells (Figure 7C, F). In general, two dots per nucleus were detected. There is evidence that nuclear puncta represent nascent transcripts (Lawrence et al., 1989; Xing et al., 1993) suggesting a pair of transcriptional foci at the *Robo1* gene of the maternal and paternal chromosomes. This prompted us to re-examine hybridization signals from other axon guidance receptors such as *Robo2*, *Rig1* and *DCC*, which showed a similar dotted pattern within the nucleus (data not shown), though not with other developmental genes (*Shh*, *Pax6*, *Nkx2.2*, *Lhx1*). In some cases, the puncta may be a consequence of very large transcription units causing delays in export to the cytoplasm.

1.6 Conclusions

Our main finding is that *Dutt1* is the predominant isoform, expressed both in post-mitotic neuronal populations and proliferating neuroepithelial cells, while *Robo1* expression is restricted to a subset of neurons. Although *Dutt1* has broader expression, some neuronal populations express only *Robo1*. Together, the summed expression of the two isoforms is broad, suggesting that *Robo1* isoforms are expressed by most if not all pioneer axons. The differential expression indicates that the alternative *Dutt1* and *Robo1* promoters contribute independently to *Robo1* spatial regulation in nervous system development. In previous studies, *Dutt1* had been associated with lung development and tumorigenesis (Xian et al., 2001).

Interestingly, the *Dutt1* promoter region has CpG islands in both mouse and human (Dallol et al., 2002). Hypermethylation of this promoter region is observed in some tumor cell lines, which results in silencing of the *Dutt1* gene (Dallol et al., 2002; Xian et al., 2004), and could contribute to *Robo1* transcriptional regulation.

There are several examples of Robo isoforms with differing N-terminal sequences. For example, Robo3 isoforms A and B have different signal peptides, and Robo3A has 29 unique amino acids (Camurri et al., 2005; Yuan et al., 1999). These isoforms of Robo3 have different affinities to Slit2 (Camurri et al., 2005). In humans, a point mutation in the Robo3 isoform A signal peptide results in horizontal gaze palsy with progressive scoliosis (HGPPS), even though the Robo3 isoform B remains intact (Jen et al., 2004). This raises an interesting point for future determination of Slit binding characteristics of Robo1 and Dutt1.

2. Experimental Procedures

2.1 Mouse embryo collection

Expression studies were carried out on CD-1 mouse embryos, obtained by timed mating, with noon of the day of the vaginal plug designated as E0.5. Pregnant females were euthanized in a CO₂ chamber and embryos were collected in ice-cold 0.1M PO₄ and fixed in 4% paraformaldehyde (PFA) in 0.1M PO₄, then stored at 4°C. The institutional Animal Care and Use Committee of the University of Nevada, Reno approved all procedures involving mice.

2.2 Generation of *Dutt1*- and *Robo1*- specific probes

Total RNA was prepared from whole E10.5 embryos (100ug) using the RNeasy Kit (Qiagen). First strand synthesis (Superscript-II) was performed using random hexamers. The cDNA pool was used to amplify *Robo1*- and *Dutt1*- specific sequences. With the help of Vector NTI (Invitrogen), isoform-specific primers were designed to amplify the unique first exons of the two isoforms. The primers used were: *Robo1* specific forward 5' GAATTCGCTGGCTGGTGTAAATGTATGGG 3', *Robo1* specific reverse 5' TAATACGACTCACTATAGGTGTGTAGCCCAGCGAA TTCT 3', *Dutt1* specific forward 5' TAGAGACTGTTCTCCCCAGCTGCCG 3', *Dutt1* specific reverse 5' TAATACGACTCACTATAGGGTGAGCAGGCTCCGCGATCA 3'. Reverse primers include a minimal T7 promoter at the 5' end. PCR conditions were as follows: 94°C for 2min, 30 cycles of 94°C for 30sec, 55°C for 30sec, 72°C for 2min, and completed with a 10 min extension at 72°C. The *Dutt1*-specific product size was 559 bp and the *Robo1*-specific product was 200 bp. Anti-sense probes for in situ hybridization were synthesized from the PCR products of *Robo1* and *Dutt1* using the T7 promoters. Transcription reaction (2ul transcription buffer, 2ul Dig labeled nucleotides (Roche), 5ul DNA, 1ul RNasin, 1ul polymerase and 9ul water) was incubated for 2 hours at 37°C. Probe was precipitated with 4M lithium chloride and 3 volumes of 100% ethanol, then resuspended in 40ul of pre-hybridization buffer (50% formamide, 5X SSC, 50ug/ml tRNA, 1.0% SDS, 50ug/ml heparin) and stored in -20 °C. A probe complementary to the shared exon3 was synthesized from a rat cDNA subclone (gift of M. Tessier-Lavigne, Genentech), which we refer to as *Robo1*^{ex3}.

2.3 In situ hybridization

Three or more embryos were analyzed for each probe. Whole embryos fixed in 4% PFA/O. 1M PO₄ were washed in PBT (PBS, 0.1% Triton-X) 3 times for 5 minutes each. Then embryos were passed through ascending (25%, 75 %, and 100%) series of methanol: PBT. After two washes in 100% methanol, a descending series of methanol was applied. After 2 washes in PBT for 5 minutes, embryos were bleached with 6% hydrogen peroxide in PBT for 10 minutes. Following 5 washes in PBT for 5 minutes each, treatment with Proteinase K (20ug/ml in PBT) was carried out for 8 minutes at room temperature. Glycine (2mg/ml in PBT) was applied for 10 minutes to inhibit Proteinase K. After refixing embryos in 0.2% glutaraldehyde in 4% PFA, embryos were placed in prehybridization buffer and incubated for 1–3 hours at 65°C (Wilkinson and Nieto, 1993). After this stage, embryos were incubated overnight with the addition of digoxigenin-labeled RNA probe. Hybridization was continued by washing embryos 3 times for 30min at 65°C in solution 1(50% formamide, 5XSSC, 1%SDS) then 3 times for 30min at 65°C in solution 3 (50% formamide, 5X SSC). Finally embryos were rinsed in TBST 3 times for 5 min at room temperature. Embryos were blocked in 10% heat-treated sheep serum in TBST for 2 hours. Anti-Dig alkaline phosphatase Fab fragments (Roche) were preadsorbed before applying to embryos. To preadsorb antibody, 3–5 mg of embryo powder in 0.5ml of TBST was incubated for 30 minutes at 70°C, then chilled on ice for 15min, followed by addition of 5ul of sheep serum and 1ul of antibody and rocked 60 minutes at 4°C. After centrifugation for 10 minutes, supernatant containing preadsorbed antibody was added to TBST containing 1% sheep serum (1:10000) and applied to embryos overnight at 4°C. After washes in TBST

for a whole day, the next morning a final wash of TBST with 2mM levamisol was done for 15 minutes. Washes were done for 3 times for 10 minutes in NTMT (100mM NaCl, 100mM Tris-HCl pH 9.5, 50mM MgCl₂, 0.1% Tween-20), then NBT/BCIP (Roche) substrate was added for the color development. After the color reaction was completed, embryos were stored in 4% PFA at 4°C.

2.4 Immunohistochemistry

To visualize neurons, after in situ hybridization, embryos (n= three or more for each probe) were sectioned and labeled with β III-tubulin antibody. For sectioning, embryos were incubated in 5% sucrose/95% 0.1M PO₄ for 30 minutes, then placed in 15% sucrose/85% 0.1M PO₄ for 4 hours to overnight. Finally, embryos were transferred into 15% sucrose/7.5% gelatin/0.1M PO₄ and incubated 6 hours to overnight at 37°C. The embryos were embedded in the gelatin solution with one embryo per block (constructed of aluminum foil). The blocks were frozen by immersion in a slurry of 2-methylbutane and dry ice, and stored at -20°C. Embedded embryos were cryosectioned at -24°C. 20 to 30 micron thick sections were placed on Superfrost Plus (Fisher) slides. Sections and whole mount embryos were washed with PGT (0.1M PO₄, 0.25% Triton-X, 2mg/ml Gelatin) several times, and then blocked in PGT plus 0.1M Lysine for an hour. β -tubulin-III antibody (Covance, 1:1000) was diluted in PGT and incubated with embryo overnight. Following washes in PGT, secondary antibody conjugated to horse radish peroxidase was added at 1:200 in PGT for 2 hours. After washes in PGT, embryos and sections were placed in 50mM Tris-HCl pH 7.6 for 30 minutes, followed by incubation with DAB solution (0.3mg/ml DAB/50mM Tris-HCl pH 7.6) for 30 minutes on ice. The peroxidase reaction was started with addition of 1% hydrogen peroxide. Color reaction was stopped in 50mM Tris-HCl and embryos stored in 4% PFA. Embryos were mounted in 80% Glycerol/20% PFA under coverslips for examination with a Leica DMR microscope, using a Leica LEI-750 CCD video camera system. Adobe Photoshop was used to assemble images into figures.

Following in situ hybridization, some of the embryos were sectioned and labeled with DAPI to visualize the nuclei of cells. 1 μ g/ml of DAPI in PBS was applied to sections for 30 seconds and washed 3 times 5 min each in PBS.

2.5 Axon Tracing

The MLF longitudinal tract was traced to its source with lipophilic fluorescent tracer DiI (Molecular Probes). Then, DiI was converted to a stable product for double labeling purposes. Both of the procedures were previously described (Nural and Mastick, 2004). Photoconverted embryos (n=3 for each tract) were labeled by in situ hybridization and sectioned for analyses.

Acknowledgements

We are grateful for helpful discussions and bioinformatics insights from Jim Lauderdale (University of Georgia) and Tom Kidd (University of Nevada). An in situ hybridization probe for the common region of *Robo1* was generously provided by Marc Tessier-Lavigne (Stanford University; Genentech). Funding for this study was provided by NIH RO1 HD38069 and the March of Dimes #1-FY06-387 to GSM.

Literature cited

- Andrews W, Liapi A, Plachez C, Camurri L, Zhang J, Mori S, Murakami F, Parnavelas JG, Sundaresan V, Richards LJ. Robo1 regulates the development of major axon tracts and interneuron migration in the forebrain. *Development* 2006;133:2243–52. [PubMed: 16690755]
- Bagri A, Marin O, Plump AS, Mak J, Pleasure SJ, Rubenstein JL, Tessier-Lavigne M. Slit proteins prevent midline crossing and determine the dorsoventral position of major axonal pathways in the mammalian forebrain. *Neuron* 2002;33:233–48. [PubMed: 11804571]

- Brose K, Bland KS, Wang KH, Arnott D, Henzel W, Goodman CS, Tessier-Lavigne M, Kidd T. Slit proteins bind Robo receptors and have an evolutionarily conserved role in repulsive axon guidance. *Cell* 1999;96:795–806. [PubMed: 10102268]
- Camurri L, Mambetisaeva E, Davies D, Parnavelas J, Sundaresan V, Andrews W. Evidence for the existence of two Robo3 isoforms with divergent biochemical properties. *Mol Cell Neurosci* 2005;30:485–93. [PubMed: 16226035]
- Camurri L, Mambetisaeva E, Sundaresan V. Rlg-1 a new member of Robo family genes exhibits distinct pattern of expression during mouse development. *Gene Expr Patterns* 2004;4:99–103. [PubMed: 14678835]
- Challa AK, McWhorter ML, Wang C, Seeger MA, Beattie CE. Robo3 isoforms have distinct roles during zebrafish development. *Mech Dev* 2005;122:1073–86. [PubMed: 16129585]
- Clark K, Hammond E, Rabbitts P. Temporal and spatial expression of two isoforms of the Dutt1/Robo1 gene in mouse development. *FEBS Lett* 2002;523:12–6. [PubMed: 12123796]
- Connor RM, Key B. Expression and role of Roundabout-1 in embryonic *Xenopus* forebrain. *Dev Dyn* 2002;225:22–34. [PubMed: 12203717]
- Dalkic E, Kescu C, Sucularli C, Aydin IT, Akcali KC, Konu O. Alternatively spliced Robo2 isoforms in zebrafish and rat. *Dev Genes Evol* 2006;216:555–63. [PubMed: 16625395]
- Dallol A, Forgacs E, Martinez A, Sekido Y, Walker R, Kishida T, Rabbitts P, Maher ER, Minna JD, Latif F. Tumour specific promoter region methylation of the human homologue of the *Drosophila* Roundabout gene DUTT1 (ROBO1) in human cancers. *Oncogene* 2002;21:3020–8. [PubMed: 12082532]
- Dickson BJ, Gilestro GF. Regulation of commissural axon pathfinding by slit and its Robo receptors. *Annu Rev Cell Dev Biol* 2006;22:651–75. [PubMed: 17029581]
- Hammond R, Vivancos V, Naeem A, Chilton J, Mambetisaeva E, Andrews W, Sundaresan V, Guthrie S. Slit-mediated repulsion is a key regulator of motor axon pathfinding in the hindbrain. *Development* 2005;132:4483–95. [PubMed: 16162649]
- Hannula-Jouppi K, Kaminen-Ahola N, Taipale M, Eklund R, Nopola-Hemmi J, Kaariainen H, Kere J. The axon guidance receptor gene ROBO1 is a candidate gene for developmental dyslexia. *PLoS Genet* 2005;1:e50. [PubMed: 16254601]
- Jen JC, Chan WM, Bosley TM, Wan J, Carr JR, Rub U, Shattuck D, Salamon G, Kudo LC, Ou J, Lin DD, Salih MA, Kansu T, Al Dhalaan H, Al Zayed Z, MacDonald DB, Stigsby B, Plaitakis A, Dretakis EK, Gottlob I, Pieh C, Traboulsi EI, Wang Q, Wang L, Andrews C, Yamada K, Demer JL, Karim S, Alger JR, Geschwind DH, Deller T, Sicotte NL, Nelson SF, Baloh RW, Engle EC. Mutations in a human ROBO gene disrupt hindbrain axon pathway crossing and morphogenesis. *Science* 2004;304:1509–13. [PubMed: 15105459]
- Kidd T, Brose K, Mitchell KJ, Fetter RD, Tessier-Lavigne M, Goodman CS, Tear G. Roundabout controls axon crossing of the CNS midline and defines a novel subfamily of evolutionarily conserved guidance receptors. *Cell* 1998b;92:205–15. [PubMed: 9458045]
- Lawrence JB, Singer RH, Marselle LM. Highly localized tracks of specific transcripts within interphase nuclei visualized by in situ hybridization. *Cell* 1989;57:493–502. [PubMed: 2541917]
- Long H, Sabatier C, Ma L, Plump A, Yuan W, Ornitz DM, Tamada A, Murakami F, Goodman CS, Tessier-Lavigne M. Conserved roles for Slit and Robo proteins in midline commissural axon guidance. *Neuron* 2004;42:213–23. [PubMed: 15091338]
- Mastick GS, Easter SS Jr. Initial organization of neurons and tracts in the embryonic mouse fore- and midbrain. *Dev Biol* 1996;173:79–94. [PubMed: 8575640]
- Nural HF, Mastick GS. Pax6 guides a relay of pioneer longitudinal axons in the embryonic mouse forebrain. *J Comp Neurol* 2004;479:399–409. [PubMed: 15514979]
- Plump AS, Erskine L, Sabatier C, Brose K, Epstein CJ, Goodman CS, Mason CA, Tessier-Lavigne M. Slit1 and Slit2 cooperate to prevent premature midline crossing of retinal axons in the mouse visual system. *Neuron* 2002;33:219–32. [PubMed: 11804570]
- Stainier DY, Gilbert W. Pioneer neurons in the mouse trigeminal sensory system. *Proc Natl Acad Sci U S A* 1990;87:923–7. [PubMed: 2300585]

- Sundaresan V, Mambetisaeva E, Andrews W, Annan A, Knoll B, Tear G, Bannister L. Dynamic expression patterns of Robo (Robo1 and Robo2) in the developing murine central nervous system. *J Comp Neurol* 2004;468:467–81. [PubMed: 14689480]
- Sundaresan V, Roberts I, Bateman A, Bankier A, Sheppard M, Hobbs C, Xiong J, Minna J, Latif F, Lerman M, Rabbitts P. The DUTT1 gene, a novel NCAM family member is expressed in developing murine neural tissues and has an unusually broad pattern of expression. *Mol Cell Neurosci* 1998;11:29–35. [PubMed: 9608531]
- Thompson H, Barker D, Camand O, Erskine L. Slits contribute to the guidance of retinal ganglion cell axons in the mammalian optic tract. *Dev Biol* 2006;296:476–84. [PubMed: 16828733]
- Wang KH, Brose K, Arnott D, Kidd T, Goodman CS, Henzel W, Tessier-Lavigne M. Biochemical purification of a mammalian slit protein as a positive regulator of sensory axon elongation and branching. *Cell* 1999;96:771–84. [PubMed: 10102266]
- Wilson SW, Easter SS Jr. A pioneering growth cone in the embryonic zebrafish brain. *Proc Natl Acad Sci U S A* 1991;88:2293–6. [PubMed: 2006169]
- Wilson SW, Easter SS Jr. Stereotyped pathway selection by growth cones of early epiphyseal neurons in the embryonic zebrafish. *Development* 1991b;112:723–46. [PubMed: 1935687]
- Xian J, Aitchison A, Bobrow L, Corbett G, Pannell R, Rabbitts T, Rabbitts P. Targeted disruption of the 3p12 gene, Dutt1/Robo1, predisposes mice to lung adenocarcinomas and lymphomas with methylation of the gene promoter. *Cancer Res* 2004;64:6432–7. [PubMed: 15374951]
- Xian J, Clark KJ, Fordham R, Pannell R, Rabbitts TH, Rabbitts PH. Inadequate lung development and bronchial hyperplasia in mice with a targeted deletion in the Dutt1/Robo1 gene. *Proc Natl Acad Sci U S A* 2001;98:15062–6. [PubMed: 11734623]
- Xing Y, Johnson CV, Dobner PR, Lawrence JB. Higher level organization of individual gene transcription and RNA splicing. *Science* 1993;259:1326–30. [PubMed: 8446901]
- Yuan W, Zhou L, Chen JH, Wu JY, Rao Y, Ornitz DM. The mouse SLIT family: secreted ligands for ROBO expressed in patterns that suggest a role in morphogenesis and axon guidance. *Dev Biol* 1999;212:290–306. [PubMed: 10433822]
- Yue Y, Grossmann B, Galetzka D, Zechner U, Haaf T. Isolation and differential expression of two isoforms of the ROBO2/Robo2 axon guidance receptor gene in humans and mice. *Genomics* 2006;88:772–8. [PubMed: 16829019]

Figure 1. The *Robo1/Dutt1* gene encodes two isoforms with different 5' ends

A) Genomic organization of mouse *Robo1* and *Dutt1* sequences, with distances in kilobases (kb) indicated below. Dark blue boxes represent *Robo1*-specific exons; the orange box represents the *Dutt1*-specific exon 2; light blue boxes represent 3' exons shared by both isoforms. The red box indicates additional 5' UTR sequence identified from *Dutt1* ESTs. B) Schematic representation of *Dutt1*- and *Robo1*-specific cDNA sequences. To make *Dutt1*- and *Robo1*-specific in situ probes, *Dutt1*- and *Robo1*-specific sequences were amplified from a mouse embryonic (E10.5) cDNA pool using primers indicated with arrows. *Dutt1*-specific primers generate a 559 bp fragment, while *Robo1*-specific primers generate a 200bp product. C) Predicted translated products of *Dutt1* and *Robo1*. Signal peptides of each isoform are underlined. In bold are shown the 32 amino acids unique to *Robo1* following the signal peptide.

Figure 2. *Dutt1* and *Robo1* are differentially expressed in E10.5 mouse CNS

Whole mount in situ hybridization labeling of E10.5 embryos, with the neural tube dissected free of external tissues. A) *Dutt1/Robo1* shared probe *Robo1*^{ex3}. B) *Dutt1*-specific probe *Dutt1*^{ex2}. C) *Robo1*-specific probe *Robo1*^{ex1a+b}. Overlapping expression domains of both isoforms are indicated with empty arrowheads. Black arrowheads indicate isoform-specific expression regions. White bars show the section planes for the following figure panels indicated. The black bar in 2A represents 500 μ m. FB, forebrain; MB, midbrain; HB, hindbrain; op, optic stalk; MN, motor neuron column.

Figure 3. Thalamic neuroepithelium and neurons express *Dutt1*, while *Robo1* expression is restricted to neurons

Embryos were labeled by whole mount in situ hybridization, then sectioned. Inset in A shows the angle of sections (black line) in A through D. A) *Dutt1* expression pattern in ventral thalamus. *Dutt1* was expressed in cells near the pial surface (N, small bracket) as well as neuroepithelial cells (NE, large bracket). Box shows the area represented in B. B) Double labeling with pan-neuronal β III-tubulin antibody (brown) shows that *Dutt1* was expressed at higher levels in neurons than in neuroepithelial cells. C) *Robo1* expression in ventral thalamus was restricted to the pial surface. D) Double labeling with β III-tubulin shows that *Robo1* expression (dark spots) was largely limited to neurons. VT, ventral thalamus; NE, neuroepithelia; N, neuronal layer; CV, cerebral vesicle. Scale bar in C: 100 μ m for panels A and C; 25 μ m for B and D.

Figure 4. *Dutt1* is the predominant isoform expressed in hindbrain

Transverse sections of hindbrain labeled by in situ hybridization for either *Dutt1*^{ex2} probe (A–D) or *Robo1*^{ex1a+b} probe (E–F) at the level of rhombomere (r) 6. A) *Dutt1* was expressed in the marginal zone of the dorsal half of the hindbrain, but included the ventral half of the ventricular zone. B) β III-tubulin labeling shows that *Dutt1* was expressed in the neurons along the margin of hindbrain. C) Higher magnification of upper boxed area in B, showing neuroepithelial labeling. D) Higher magnification of boxed area in B located close to ventral midline (floor plate, fp). *Dutt1* was expressed in a subset of motor neurons. E) *Robo1* expression was restricted to the dorsal and ventral margins of the hindbrain. F) Double labeling of *Robo1* in situ hybridization with β III-tubulin antibody. G) Higher magnification of boxed area in F. *Robo1* was expressed in motor neurons located close to the margin of the hindbrain. rp, roof plate. Scale bar in 4E represents 100 μ m in A, B, E, F; 30 μ m in C, D; 15 μ m in G.

Figure 5. *Dutt1* is the main isoform expressed in spinal cord commissural neurons

Transverse sections through E10.5 spinal cord, labeled with either *Dutt1*^{ex2} (A–D) or *Robo1*^{ex1a+b} (E–G) probes. A) Expression pattern of *Dutt1* in spinal cord, showing dorsal labeling in region containing commissural neurons (c), a band of neuroepithelial expression at an intermediate position, and a ventral region medial to the motor column (MN). B) Combined label with β III-tubulin antibody and *Dutt1*, showing expression in dorsal commissural neurons, as well as tissue underlying motor neurons. C) Close up of the area indicated by the left box in B. Commissural neurons (c) labeled for β III-tubulin and *Dutt1*. D) Close up of the area in the right box in B. *Dutt1* was expressed in dorsal root ganglia (DRG). E) *Robo1* expression. F) *Robo1* was expressed by motor neurons, contrasting with the more medial expression of *Dutt1*. G) High magnification of boxed area in F. Some commissural neurons expressed *Robo1*. fp, floor plate; rp, roof plate. Scale bar in 5E represents 100 μ m in A, B, E, F, and 20 μ m in C, D, and G.

Figure 6. *Dutt1* and *Robo1* are expressed in pioneer longitudinal neurons

Inset in A shows the projections of two pioneer longitudinal tracts, with the dashed lines indicating planes of section. MLF cell bodies were back labeled with DiI from the hindbrain, and following photoconversion (brown), sections were labeled by in situ (purple) for *Dutt1* (A) or *Robo1* (C). A) *Dutt1* was not expressed in MLF neurons. However, it was expressed in nearby neuroepithelial cells (arrowhead). C) *Robo1* probe labeled a subset of MLF neurons. B, D) β III-tubulin labeling of TPOC neurons, following in situ hybridization with *Dutt1* or *Robo1*. Both *Dutt1* and *Robo1* were expressed in TPOC neurons (white arrowheads). Scale bar in C represents 20 μ m in A and C; 10 μ m in B, D.

Figure 7. *Dutt1* and *Robo1* transcripts accumulate in puncta within the nucleus of cells
 Embryos were labeled with either *Dutt1* (A–C) or *Robo1* (D–F) RNA probe, then sectioned. A, D: Both *Dutt1* and *Robo1* had punctate expression within non-neuronal cells. B, E: DAPI labeling of the same sections suggest that puncta are localized to the nucleus. C, F: βIII-tubulin + neurons had punctate expression for *Dutt1* and *Robo1*. Scale bar in D represents 20 μm in A, B, D, and E; 10 μm in C, F.