Authors' affiliations

M D S Erlewyn-Lajeunesse, L P Hunt, Clinical Sciences at South Bristol, Faculty of Medicine, University of Bristol, UK

K Coppens, P J Chinnick, P Davies, I M Higginson, Children's Emergency Department, Bristol Royal Hospital for Children, Bristol, UK
J R Benger, Faculty of Health and Social Care, University of the West of England, Bristol, UK

Funding: this study was funded by the Anthony Hopkins Memorial Prize, awarded by the Faculty of Accident and Emergency Medicine as an unrestricted award to the Emergency Department

Competing interests: none declared

Correspondence to: Dr M Erlewyn-Lajeunesse, Institute of Child Life & Health, UBHT Education Centre, Upper Maudlin Street, Bristol BS2 8AE, UK; Mich.Lajeunesse@bristol.ac.uk

Accepted 1 February 2006

Published Online First 7 February 2006

REFERENCES

- 1 **Mayoral CE**, Marino RV, Rosenfeld W, et al. Alternating antipyretics: is this an alternative? *Pediatrics* 2000;**105**:1009–12.
- 2 **Uhari M**, Rantala H, Vainionpaa L, et al. Effect of acetaminophen and of low intermittent doses of diazepam on prevention of recurrences of febrile seizures. *J Pediatr* 1995;**126**:991–5.
- 3 **Perrott DA**, Piira T, Goodenough B, et al. Efficacy and safety of acetaminophen vs ibuprofen for treating children's pain or fever: a meta-analysis. *Arch Pediatr Adolesc Med* 2004;**158**:521–6.
- 4 **Moghal NE**, Hegde S, Eastham KM. Ibuprofen and acute renal failure in a toddler. *Arch Dis Child* 2004;**89**:276–7.
- 5 **Richardson J**, Sills J. Hypothermia following fever. *Arch Dis Child* 2004;**89**:1177.

IMAGES IN PAEDIATRICS

doi: 10.1136/adc.2004.068056

An Eiffel penetrating head injury

A 3 year old boy presented to our accident and emergency department with an obvious penetrating head injury. He had tripped and fallen onto a metal model of the Eiffel Tower which then became rigidly lodged into his skull.

On arrival he had a Glasgow coma score of 15 and was neurologically intact. He was then anaesthetised for a computed tomography scan which showed the tip of the metallic model penetrating the skull and lying 11 mm into the brain parenchyma.

He was transferred directly to the neurosurgical theatre for a craniotomy to remove the foreign body and debridement of the wound. Following this procedure he was successfully extubated and made a good recovery on the paediatric intensive care unit.

The following day he was discharged to the ward with regular antibiotics and prophylactic phenytoin.

This case report highlights the dangers to children from seemingly innocuous objects and toys.¹

M George, J Round

St George's Hospital Medical School, Blackshaw Road, London SW17, UK;
 marinageorge@hotmail.com

Competing interests: none declared

Reference

- 1 **Couper GW**, Boddie DE, Eljamel MS, et al. Unguarded electric plugs cause penetrating head injuries in children. *J Accid Emerg Med* 2000;**17**:55.