Published in final edited form as:

Hum Immunol. 2007 October ; 68(10): 839–843.

Search for Quantitative Trait Loci of Atopy-Associated Immune Responses Using Allergen-Specific IgG1 as an ‘Endophenotype’

Duaine R. Jackola¹, Michael B. Miller², Carol L. Liebeler¹, and Malcolm N. Blumenthal¹

¹ *The Asthma and Allergy Center, Department of Medicine, University of Minnesota Medical School, Minneapolis, MN.*

² *Division of Epidemiology, School of Public Health, University of Minnesota Medical School, Minneapolis, MN.*

Abstract

Inherited atopic diseases of humans arise from adverse adaptive humoral responses to non-infectious environmental allergens. We have previously reported that allergen-specific IgG1 provides more reliable heritability estimates for responses to allergens than total IgE. Genome scans were done for 91 Caucasian nuclear families with history of atopy for total IgE and IgG1 produced against a common major allergen from house dust mite, Der p 1. Suggestive associations for Der p 1-IgG1 production were found at 7 quantitative trait loci (QTL) (logarithm of the odds, $\text{LOD} \geq 1.23$; $p \leq 0.009$) with QTL-specific heritabilities of 73% – 80%. Scans using total IgE found suggestive associations for 12 QTL's ($\text{LOD} \geq 1.44$; $p \leq 0.004$), but QTL-specific heritabilities only in the range of 30% – 35%. Allergen-specific IgG1 is a suitable ‘endophenotype’ to be used in searches for genes associated with atopy-associated humoral immune responses to common aeroallergens.

Keywords

Atopy; Allergen-IgG1; Endophenotype; Genome Scan; IgE

INTRODUCTION

Atopic disorders, such as allergic rhinitis or atopic bronchial asthma, are common inherited diseases that result from adverse immune system responses to non-infectious environmental allergens. They are generally characterized by elevated serum total IgE levels or evidence for the production of allergen-specific IgE manifest by positive skin prick test (SPT) results. However, genome scans using these phenotypes have found numerous chromosomal regions putatively associated with these traits, but no clear consensus has arisen [1].

Numerous metabolic pathways contribute to the production of IgE [2] that can be influenced by numerous environmental factors such as history of infectious disease, environmental pollution levels, population health practices and others. Incidence of atopic disease varies widely in different geographic locales due to the impact of these factors [3]. Quantitative IgE

Correspondence: Duaine R. Jackola, Ph. D., The Asthma & Allergy Center, University of Minnesota Medical School, Mayo Mail Code 434, 420 Delaware St. S.E., Minneapolis, Minnesota 55455. Telephone: +1 612-626-4624. FAX: +1 612-624-9188. E-mail: drjackola@q.com..

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

traits associated with atopy show a great amount of phenotypic variability, which probably accounts for much of the observed variation observed in genome scan results [1].

Production of allergen-specific IgE is the result of an adaptive, memory-forming humoral immune response to non-infectious environmental antigens (allergens). Save for specific IgE production, it has been shown that it is equally likely for anyone to mount a T cell-dependent humoral response to allergens involving varying amounts of the IgA and IgG isotype subclasses [4,5]. Of special note, specific IgG1 is elevated among those with atopic sensitization to a particular allergen [6].

In this journal, we have previously shown among nuclear families with medical histories of atopic diseases that the total narrow sense heritability of serum total IgE is about 48% [7]. However, this estimate was significantly influenced by age, gender and atopic clinical status. In contrast, using the major Der p 1 allergen from house dust mite it was found that the total narrow sense heritability of atopy-associated Der p 1-IgG1 responsiveness was about 62%. Importantly, this estimate was not significantly influenced by age, gender or overall atopic status [7]. Thus, for gene searches for atopy-associated immune response, allergen-specific IgG1 may be a suitable 'endophenotype' [8].

We conducted genome-scans in these families using Der p 1-IgG1 as an 'endophenotype,' and compared these to results for total IgE. As with the heritability estimates [7], the results suggest that allergen-specific IgG1 is a more robust phenotype for conducting gene searches for these complex human immune system responses to common aeroallergens.

METHODS

Study Population

Caucasian families residing in the Minneapolis/St. Paul, Minnesota metropolitan area that included two children with a physician's diagnosis of asthma were recruited through media advertisements as part of studies for the genetics of asthma. We evaluated 367 people from 91 nuclear families that included both parents, the asthmatic children and their siblings. A total of 169 parents (mean age = 43.7 years; range = 22–75 years) and 198 children (mean age = 14.9 years; range = 6–42 years) were screened. There were 176 females and 191 males. Each person, or legal guardian, provided written, informed consent according to the guidelines of the Institutional Review Board of the University of Minnesota.

Skin Prick Tests (SPT)

Skin prick testing was done as previously described [9] using 14 standardized allergen extracts for dust mites (2), including Der p allergen extract, ragweed, molds (3), trees (4), grasses (2), animal danders (2). A positive SPT result was recorded for a wheal area $\geq 9 \text{ mm}^2$ above negative saline control.

Der p 1 Allergen

The major allergen Der p 1 from house dust mite was purified by methods previously described [10]. We chose Der p 1 allergen as a model, as we have previously shown that the atopic response to any particular allergen is not obligately inherited. Rather, it is a random variable independent of any family history of atopic allergen sensitivity [11,12]. Der p 1 is a common non-seasonal indoor allergen that induced the highest incidence of atopic sensitization among all of the common indoor allergens that were tested in the SPT screens.

Serum Antibody Tests

Serum total IgE tests were done using the ACCESS Immunoassay System® (Beckman Coulter, Chaska, MN), according to manufacturer's specifications, with results reported as IU/ml (1 IU = 2.4 ng). Measures of Der p 1-specific IgG1 were done using modifications of 'reverse sandwich' ELISA assays previously described [7]. This assay gave a linear response on a log-log scale over the range of 0.05 to 100 ng/ml.

Genotyping

DNA was isolated from peripheral blood white cells by standard methods [9]. DNA samples were genotyped using Weber Screening Set 6 comprised of 360 markers whose positions spanned all chromosomes (≈ 10 cM spacing) by the Marshfield Center for Medical Genetics, Marshfield, WI (<http://research.marshfieldclinic.org/genetics>).

Genome Scans

Variance components linkage methods were employed accounting for pedigrees of arbitrary size and complexity [13]. This approach is based upon specifying the expected genetic covariances between relatives as a function of their identity-by-descent (IBD) relationships at a quantitative trait locus (QTL). The QTL mapping procedure was implemented in two phases: (1) estimation of IBD probabilities for all pairs of relatives in all pedigrees at all chromosomal positions conditional on their marker genotypes, followed by (2) estimation of effects of loci by correlating genetic IBD similarity at those loci for pairs of relatives with the phenotypic similarity of those relatives. IBD and variance components estimations were generated with the software program MERLIN [14] using supercomputer facilities of the Minnesota Supercomputing Institute (MSI).

RESULTS

Clinical Characteristics

A total of 367 individuals (176 female, 191 male) from 91 Caucasian nuclear families with history of atopic asthma were screened. Using a battery of 14 common aeroallergens, there was a high proportion of skin prick test positive (SPT [+]) response: 262/367 (71%). Among those with any SPT [+] result, 127/262 (48.5%) was SPT [+] for the house dust mite Der p allergen. The study population showed significant atopic sensitivity to common aeroallergens, and dust mite (Der p) allergen in particular. (Additional clinical characteristics have been previously reported [7].)

Allergen-Specific IgG1 as an 'Endophenotype'

As expected, there was a significant positive correlation between serum total IgE values and the number of SPT [+] results: $r = 0.56$; $t(366) = 12.8$; $p \ll 0.001$. However, as previously reported, total IgE values varied significantly due to age and gender [7]. Also, total IgE values were elevated among those with any SPT [+] result, and were highest among those with any SPT [+] who also had asthma (Table 1 in [7]). IgE levels are influenced both by allergen sensitivity and asthma-associated inflammation.

In contrast, allergen-specific IgG1 was not significantly influenced by these variables. Using the major dust mite allergen Der p 1 as a model, we have also reported for this study population that Der p 1-IgG1 values are not influenced by age, gender or asthmatic respiratory involvement [7]. Der p 1-IgG1 values are, however, elevated among those who are also sensitized to the Der p allergen extract (Figure 1 in [7]). As reported, for those who were SPT [+] for the Der p allergen extract, the average log-10 value for Der p 1-IgG1 was -0.64 (std. dev. = 0.51 ; $N = 127$). For those who were SPT [-] for the Der p extract, average log-10 Der p 1-IgG1 = -0.80

(std. dev. = 0.25; N = 240). By two-sided t-test, these values were significantly different: $t(365) = 3.17$; $p = 0.0008$.

There was no significant correlation between Log [Total IgE] and Log [Der p 1-IgG1] values: $r = 0.09$; $t(366) = 1.26$; $p = 0.21$. There was, however, a slight although statistically significant correlation between Log [Der p 1-IgG1] values and the number of SPT [+] results: $r = 0.15$; $t(366) = 2.91$; $p = 0.004$. This result may have been largely due to the high proportion of people who were SPT [+] for the Der p allergen in this study population.

Der p 1-IgG1 is a suitable 'endophenotype' for gene searches for atopy-associated immune responses to aeroallergens [8]; it is not influenced by age or gender effects, it is not elevated due to asthmatic respiratory involvement, but it is elevated among those with established atopic sensitivity to allergen(s).

Comparisons for Total IgE and Der p 1-IgG1 of Gene Scan Results for Quantitative Trait Loci (QTL)

Whole genome scans were done using 360 polymorphic markers comprising Weber Screening Set 6, with average inter-marker distance of ≈ 10 cM. Searches were done for quantitative trait loci (QTL) on all chromosomes for serum total IgE values and allergen-specific IgG1 using house dust mite Der p 1-IgG1 as a model. Screen results were conducted using the computer software MERLIN [14], which provided best estimates for the logarithm of the odds for association (LOD Score), its associated QTL-specific heritability (QTL-H2) and the p-value for these estimates. Table 1 shows the results of these scans.

Overall, 7 QTL's were identified with suggestive association to Der p 1-IgG1 production ($\text{LOD} \geq 1.23$; $p \leq 0.009$). Of interest, the QTL-specific heritability estimates for these loci were in the range of 73% to 80%. We have previously reported for this study population that the total narrow sense heritability for Der p 1-IgG1 is 61.5% [7].

Using total IgE values provided suggestive evidence at 12 QTL's ($\text{LOD} \geq 1.44$; $p \leq 0.004$). For comparison, the QTL LOD scores for total IgE were generally higher than that of Der p 1-IgG1. However, the QTL-specific heritability estimates for the IgE scans (30% – 35%) suggest that these results were less robust. This has been previously suggested by the total narrow sense heritability estimate for total IgE of 47.8% that was significantly influenced by age, gender and clinical status effects [7].

Although the results are not shown, we also did gene scans using the number of SPT [+] results as a quantitative trait. No LOD score higher than 0.74 ($p = 0.03$; $\text{QTL-H2} = 31\%$) was found. These results are consistent with previous findings that the number of SPT [+] results is a random variable whose quantitative expressions are not an inherited trait [11,12]. In contrast, Der p 1-IgG1 production is evident in all the subjects studied, and is a marker associated with an immune response to a well-defined allergen known to be associated with atopy.

DISCUSSION

Searches for chromosomal loci associated with the complex human atopic diseases have produced widely conflicting results, with numerous loci putatively associated with disease-related traits such as serum total IgE levels or skin prick test (SPT) results [1]. These studies assumed that the total phenotype expression was a simple summation of the contributions of multiple underlying causative elements (multiple gene effects). Yet, this strategy may overlook the behavior of the whole (complex) system that is influenced both by multiple genes as well as significant non-genetic, random environmental effects [15].

The choice of a particular phenotypic trait to be used in genome-wide scanning should be reflective of underlying physiological processes that are common to all, but are only abnormal in the advent of a disease [8]. In the case of the atopic diseases associated with immune system sensitivity to common, non-infectious environmental aeroallergens, it has been implicitly assumed that serum total IgE levels is such a trait. However, we have previously argued that serum total IgE may actually comprise at least two physiologically distinct components [16].

One is a basal component unrelated to atopic disorders, while a second arises from the production of allergen-specific IgE. A simple manifestation of increased levels of this fraction is reflected by an increased number of SPT [+] results [16]. Each SPT [+] result represents a unique 'clone' of lymphoid cells that respond specifically to and produce IgE against a particular allergen. But, the number and specific allergens to which people respond are random variables independent of any 'inherited propensity' to adversely respond to aeroallergens [11,12].

Further, not only is serum total IgE elevated among those who have any SPT [+] result [16], it is higher still among those who are SPT [+] and also have asthma [12]. This was due to the observation that among those with any SPT [+] result, the average number of SPT [+] results was significantly higher among those with asthma compared to those without asthma. The allergen-specific IgE fraction is influenced both by sensitivity to allergens and asthmatic inflammatory processes. Total IgE is not a robust phenotype to be used for gene scans related to atopic immune responses to allergens.

This is borne out in the results shown in Table 1. As with many previous studies [1], a number of quantitative trait loci (QTL) show suggestive associations to IgE production. Yet, on average, the estimates for QTL-specific heritability are, at best, modest (30% – 35%). This may have arisen from the inability to fully account for the several distributions of IgE that could arise.

There are numerous metabolic networks that can contribute to IgE production, although their complex interactions are not well understood [2]. From *in vivo* [17] and *in vitro* [18,19] studies, isotype class switching to IgE is far less common than a switch to IgG production. These results derived from IgG or IgE production under identical experimental conditions. The principle difference was the total time until the appearance of these isotypes, with IgG appearing far sooner than IgE. Both IgG and IgE production share common metabolic pathways leading to their production, but in probabilistic terms it is far more likely for an IgG response to be manifest in the presence of a defined antigen than is IgE.

Allergen-specific IgG production is probably more common than is specific IgE, as has been independently observed by our group [6] and others [4,5]. In particular, allergen-specific IgG1 is elevated among those who are also atopically sensitized to a particular allergen, like ragweed pollen [6] or house dust mite [7]. The production of allergen-specific IgG1 may not be atopic disease associated, but it is associated with immune system humoral response development to antigens known to be atopy-associated. Further, as a quantitative trait for use in genome scans, specific IgG1 is more robust than total or specific IgE as it is not significantly influenced by age, gender or overall atopic clinical status, and it is a reasonably well-defined immunological response to a well-defined source of antigenic stimulation [7].

Thus, allergen-specific IgG1 is an 'endophenotype' [8] that appears to provide more reliable information from gene scans than do the IgE traits. This is borne out by the results in Table 1. While we cannot say with certainty that the QTL's identified are linked to specific IgG1 production, it does appear that the results from the scan for this trait are more reliable than those for IgE, as the QTL-specific heritability estimates for specific IgG1 are significantly higher (73% – 80%) compared to those for total IgE (30% – 35%).

The complex atopic diseases of humans, like allergic rhinitis or bronchial asthma, are largely due to enhanced immune system responsiveness to otherwise biologically benign, non-infectious environmental allergens. Aside from disease manifestation, the atopic and non-atopic immune responses may have more in common than are disparate, including vigorous humoral responses to these antigens [6,20]. Before the genes associated with atopic disease can be found, it may be best to first unravel the genetic components involved with immune responses to these unique types of antigens.

Acknowledgements

Supported by NIH grant # RO1 HL049609.

References

1. Ober C, Hoffjan S. Asthma genetics 2006: the long and winding road to gene discovery. *Genes Immunity* 2006;7:95. [PubMed: 16395390]
2. Geha RS, Jabara HH, Brodeur SR. The regulation of immunoglobulin E class-switch recombination. *Nature Rev Immunol* 2003;3:721. [PubMed: 12949496]
3. ISAAC. Worldwide variation in the prevalence of asthma symptoms: the International Study of Asthma and Allergies in Childhood (ISAAC). *Eur Resp J* 1998;12:315.
4. Batard T, Basuyaux B, Lambin P, Bremard-Oury C, Hamilton RG, David B, Peltre G. Isotypic analysis of grass pollen-specific immunoglobulins in human plasma. 1. Specialization of certain classes and subclasses in the immune response. *Int Arch Allergy Immunol* 1993;100:68. [PubMed: 8428167]
5. Batard T, Basuyaux B, Laroze A, Lambin P, Bremard-Oury C, Aucouturier P, Hamilton RG, David B, Peltre G. Isotypic analysis of grass-pollen-specific immunoglobulins in human plasma. 2. Quantification of the IgE, IgM, IgA class and the IgG subclass antibodies. *Int Arch Allergy Immunol* 1993;102:279. [PubMed: 8219781]
6. Jackola DR, Pierson-Mullany LK, Liebler CL, Blumenthal MN, Rosenberg A. Variable binding affinities for allergen suggests a 'selective competition' among immunoglobulins in atopic and non-atopic humans. *Molec Immunol* 2002;39:367. [PubMed: 12220894]
7. Liebler CL, Basu S, Jackola DR. Allergen-specific IgG1 provides parsimonious heritability estimates for atopic immune responses to allergens. *Human Immunol* 2007;68:113. [PubMed: 17321901]
8. Rice, JP.; Saccone, NL.; Rasmussen, E. Definition of the phenotype. In: Rao, DC.; Province, MA., editors. *Genetic Dissection of Complex Traits*. San Diego: Academic Press; 2001.
9. CSGA (Collaborative Study on the Genetics of Asthma). A genome-wide search for asthma susceptibility loci in ethnically diverse populations. *Nature Genetics* 1997;15:389. [PubMed: 9090385]
10. Pierson L, Allauzen S, Blumenthal M, Rosenberg A. An automated method for determination of antibody affinity functions with nanogram quantities. *J Immunol Methods* 1998;211:97. [PubMed: 9617835]
11. Jackola DR, Liebler CL, Blumenthal MN, Rosenberg A. Random outcomes of allergen-specific responses in atopic families. *Clin Exp Allergy* 2004;34:540. [PubMed: 15080805]
12. Jackola DR. Random allergen-specific IgE expression in atopic families: evidence for inherited 'stochastic bias' in adverse immune response development to non-infectious antigens. *Molec Immunol* 2007;44:2549. [PubMed: 17250894]
13. Almasy L, Blangero J. Multipoint quantitative-trait linkage analysis in general pedigrees. *Am J Hum Genet* 1998;62:1198. [PubMed: 9545414]
14. Abecasis GR, Cherny SS, Cookson WO, Cardon LR. Merlin---rapid analysis of dense genetic maps using sparse gene flow trees. *Nature Genet* 2002;30:97. [PubMed: 11731797]
15. Marsh DG. Approaches toward the genetic analysis of complex traits. *Asthma and atopy. Am J Resp Crit Care Med* 1997;156:S133. [PubMed: 9351594]
16. Jackola DR, Blumenthal MN, Rosenberg A. Evidence for two independent distributions of serum immunoglobulin E in atopic families: Cognate and Non-Cognate IgE. *Human Immunol* 2004;65:20. [PubMed: 14700592]

17. Jung S, Siebenkotten G, Radbruch A. Frequency of immunoglobulin E switching is autonomously determined and independent of prior switching to other classes. *J Exp Med* 1994;179:2023. [PubMed: 8195724]
18. Tangye SG, Ferguson A, Avery DT, Ma CS, Hodgkin PD. Isotype switching by human B cells is division-associated and regulated by cytokines. *J Immunol* 2002;169:4298. [PubMed: 12370361]
19. Hasbold J, Corcoran LM, Tarlinton DM, Tangye SG, Hodgkin PD. Evidence from the generation of immunoglobulin G-secreting cells that stochastic mechanisms regulate lymphocyte differentiation. *Nature Immunol* 2004;5:55. [PubMed: 14647274]
20. Jackola DR, Liebeler CL, Lin C-Y, Chiu Y-K, Blumenthal MN, Rosenberg A. Evidence that negative feedback between antibody concentration and affinity regulates humoral response consolidation to a non-infectious antigen in infants. *Molec Immunol* 2005;42:19. [PubMed: 15488940]

Abbreviations

Der p	House dust mite allergen (<i>Dermataphagoides pteronissinus</i>)
LOD	Logarithm of the odds
SPT	(percutaneous) skin prick test
QTL	Quantitative trait locus
cM	Centi-Morgan

Table 1

Genome Scan Results

CHR	Position (cM)	LOD	Der p 1-IgG1 QTL-H2 ¹	p-value	LOD	Total IgE QTL-H2	p-value
1	13.4	---			2.46	0.35	0.0004
1	156.7	2.06	0.80	0.001	1.54	0.33	0.004
2	84.3	1.23	0.73	0.009	---		
2	252.1	---			2.05	0.34	0.0011
3	115.9	---			2.52	0.37	0.0003
4	51.2	1.31	0.78	0.007	---		
4	100.4	---			1.91	0.38	0.002
4	119.8	---			1.44	0.33	0.005
7	193.6	1.61	0.77	0.003	---		
8	72.5	1.43	0.73	0.005	1.56	0.35	0.004
10	113.6	---			2.65	0.37	0.0002
12	150.5	---			2.45	0.35	0.0004
13	97.9	---			2.37	0.35	0.0005
16	36.0	---			2.75	0.36	0.0002
17	112.1	1.31	0.79	0.007	---		
20	10.6	---			3.04	0.40	0.00009
20	90.4	1.39	0.74	0.006	---		

¹QTL-H2 = Quantitative Trait Locus-specific heritability estimate.