were palpable only. The distribution of the lesions was asymmetrical. There were none on the face, back, or neck, and there has been no evidence of suppuration. A low-grade pyrexia was present during each observed attack, but the general health has been moderately satisfactory.

The history, together with the typical lesions and characteristic microscopical appearances, confirms the diagnosis. Photographs were shown to Dr. Parkes Weber, who considered the appearances to be consistent with this diagnosis.

Histological examination revealed necrosis of the subcutaneous fat. The epidermis was intact and normal. The subcutaneous fat showed extensive foci of necrosis of the fat cells, these areas being replaced by groups of lipid-filled phagocytes. Some giant-cell reaction was seen, together with lymphocyte infiltration.

I am indebted to Dr. Alexander Smith, medical superintendent, Stonehouse Hospital, for permission to publish this case.

H. R. F. MACDONALD, M.B., F.R.F.P.S.,
Senior Registrar Physician,
Stonehouse and Cleland Hospitals, Lanarkshire.

REFERENCES

- Carleton, Alice (1949). In Parkes Weber's *Further Rare Diseases and Debatable Subjects*, p. 103. Staples Press, London.
Christian, H. A. (1928). *Arch. intern. Med.*, 42, 338.
Weber, F. Parkes (1925). *Brit. J. Derm.*, 37, 301.

Lymphosarcoma of the Appendix

Since Kundrat (1893) first described lymphosarcoma of the appendix many different names have been used to describe this condition. Ewing (1939) endeavoured to clarify the position by stating that lymphoid tissue contained three cellular elements—namely, lymphocytes, reticulum cells of the pulp and follicles, and endothelial cells of the pulp and cavernous sinuses. Thus he maintained that there were two possible types of sarcoma—a malignant lymphocytoma and a reticulum-cell sarcoma.

Lymphosarcoma can appear in the tonsils, lymph nodes, spleen, and thymus gland, and to a less extent in the lymph follicles of the gastro-intestinal tract. Geschichter (1935) stated that sarcoma comprises 3% of all malignant growths of the gastro-intestinal tract—mostly lymphosarcoma. In his series of 46 lymphosarcomata 13 were in the stomach, 20 in the ileum, 10 in the caecum, 2 in the rectum, and 1 in the duodenum.

Our patient had a lymphosarcoma of the appendix, and we record the case to draw attention to the possibility of misdiagnosis as chronic appendicitis, which is so frequent and which may give rise to a false sense of security.

CASE REPORT

A man aged 38 was seen on May 17, 1950. He had a 10-year history of intermittent diarrhoea and vague abdominal discomfort, which during the last 12 months had become a more definite pain localized to the right iliac fossa. Examination revealed no abnormal finding except tenderness localized to the lower pole of the distended caecum. A magnesium sulphate barium meal examination on May 24 showed that the appendix failed to fill.

He was admitted to hospital on August 30. Routine questioning revealed that the pain in the right iliac fossa was a sporadic dull ache which occasionally radiated to the left side of the abdomen. During the previous month his appetite had diminished and his abdomen had become "swollen and tender." His bowels were regular, with a slight tendency to loose motions, during which the pain was accentuated. The pain was relieved by rest and codeine. His only relevant past history was migraine since the age of 16 and glandular fever at 32. His weight was 8 st. 8 lb. (54.4 kg.) and his temperature and full blood count were normal. Chest radiography revealed only a calcified focus at the left apex. On the day before admission the symptoms of pain and nausea became prominent, and it was thought that an acute attack might be supervening.

An operation was performed on September 1. Through a gridiron incision a bulbous-ended pear-shaped appendix 2½ in. (5.7 cm.) long was found lying retrocaecally, with a

hyperaemic base and a similarly hyperaemic caecum. The appendicular mesentery was thick, contained much firm fat, and was extremely vascular. On the serous surface of the organ there was a purplish discoloration close to the base, which was attributed to the local necrosis in the appendix. Post-operative convalescence was uneventful, and the patient was discharged on September 10. Section of the appendix showed replacement of the mucosa and submucosa by masses of lymphoid cells with penetration of the muscle coat in one area. The appearances were those of lymphosarcoma.

During the following 10 months he was seen on eight occasions by both of us. In November, 1950, his weight was 9 st. 4 lb. (58.9 kg.). His appetite has been good, his bowel action normal, and he has been in full active employment. The only clinical finding has been a slightly tender boggy caecum, without any lymphatic glandular enlargement. A barium meal examination and blood count were normal at this time and also in February, 1951. It is significant that tenderness in the right iliac fossa over what might be described as a thickened caecum and a fatty mesentery was still present. This tenderness does not seem to alter. In November, 1953, the patient was still in good health.

COMMENT

Charache (1934) reported a case of primary leiomyosarcoma of the appendix. He summarized 22 reported cases of sarcoma of the appendix. Every one of these cases was diagnosed only on histological section. Clinically the diagnosis was an appendicitis of one kind or another, and the cases were reported only because of the rarity of the lesion. Most of these 23 patients were in the first or third decade of life. The symptoms in all instances were referable to the right iliac fossa, but their duration varied between four weeks and a year, although in two cases there had been symptoms for 10 and 13 years. The occurrence of "attacks" was noticeable in seven cases, whilst in all but two of the remainder the attack persisted for several weeks or presented as acute appendicitis. One case was an incidental finding at necropsy in a man aged 55 who had died of cerebral haemorrhage. Bizard *et al.* (1938) recorded a case of a 27-year-old man who had a lymphoblastic sarcoma; a three-year follow-up showed that the patient was still well. Knox (1945) described a further case in a boy aged 4, a 15-months follow-up recording that the patient was then essentially asymptomatic.

In our patient an interesting feature was the 10-year history of vague and repeated abdominal upsets. Is it that the repeated subacute attacks of appendicitis had slowly altered the structure of the lymphoid tissue and that a gradual change took place from an inflammatory reaction to a state of malignancy? The attack of glandular fever is of particular interest: had it any influence on the presumed development of the sarcomatous change? Other questions may be asked. How many appendices are removed and never sectioned? What is the true incidence of sarcoma in the organ? Is a right hemicolectomy the correct treatment, or local excision followed by radiotherapy?

The prognosis seems to be about seven months. Three years is the longest period of freedom from recurrence that has so far been recorded.

Our grateful thanks are due to Mr. R. Vaughan Hudson, senior surgeon, Middlesex Hospital, for permission to publish this case; to Professor R. Scarff, director of the Bland-Sutton Institute of Pathology, for the section report; and to Sir Harold Graham Hodgson, director of the X-ray Diagnostic Department.

F. AUSTIN HENLEY, M.B., B.S., F.R.C.S.,
Assistant Surgeon, Central Middlesex Hospital.

W. W. SLACK, B.M., B.Ch.,
Surgical Registrar, Middlesex Hospital.

REFERENCES

- Bizard, G., *et al.* (1938). *Rev. Chir., Paris*, 57, 195.
Charache, H. (1934). *Amer. J. Surg.*, 26, 357.
Ewing, J. (1939). *Bull. N.Y. Acad. Med.*, 15, 92.
Kundrat, H. (1893). *Wien. Klin. Wschr.*, 6, 211, 234.
Geschichter, C. F. (1935). *Amer. J. Cancer*, 25, 130.
Knox, G. (1945). *Arch. Surg.*, 50, 288.