laboratory staff for help with the eosinophil counts, and to Miss M. Hayes. Finally, the co-operation of Sister M. Nicholls and other members of the nursing staff is gratefully acknowledged.

REFERENCES

- Baker, R. D. (1945). *Amer. J. Path.*, **21**, 717.
 Bardeen, C. R. (1898). *Johns Hopk. Hosp. Rep.*, **7**, 137.
 de Groot, J., and Harris, G. W. (1950). *J. Physiol.*, **111**, 335.
 Dougherty, T. F., and White, A. (1947). *J. Lab. clin. Med.*, **32**, 584.
 Dunger, R. (1910). *Munch. med. Wschr.*, **57**, 1942.
 Evans, E. I., and Butterfield, W. J. H. (1951). *Ann. Surg.*, **134**, 588.
 Forsham, P. H., Thorn, G. W., Prunty, F. T. G., and Hills, A. G. (1948). *J. clin. Endocr.*, **8**, 15.
 Harris, G. W. (1951). *British Medical Journal*, **2**, 627.
 Hills, A. G., Forsham, P. H., and Finch, C. A. (1948). *Blood*, **3**, 755.
 Hume, D. M., and Wittenstein, G. J. (1950). *Proc. First Clin. A.C.T.H. Conf.*, edited by J. R. Mote. Blackiston, Philadelphia.
 King, E. J. (1951). *Micro-analysts in Medical Biochemistry*, p. 142. Churchill, London.
 Long, C. N. H., and Fry, E. G. (1945). *Proc. Soc. exp. Biol. N.Y.*, **59**, 67.
 McDermott, W. V., Fry, E. G., Brobeck, J. R., and Long, C. N. H. (1950). *Ibid.*, **73**, 609.
 Prunty, F. T. G. (1949). *Proc. roy. Soc. Med.*, **42**, 267.
 — (1950). *J. clin. Path.*, **3**, 87.
 Rich, A. R. (1944). *Bull. Johns Hopk. Hosp.*, **74**, 1.
 Sevvitt, S. (1951). *British Medical Journal*, **1**, 976.
 Thorn, G. W., Prunty, F. T. G., and Forsham, P. H. (1947). *Trans. Ass. Amer. Phys.*, **60**, 143.
 Vogt, M. (1944). *J. Physiol.*, **103**, 317.
 Wight, A., Raker, J. W., Merrington, W. R., and Cope, O. (1953). *Ann. Surg.*, **137**, 175.

THE ACHLORHYDRIC PATIENT WITH DYSPEPSIA

BY

B. F. SWYNNERTON, D.M., M.R.C.P.

Weir Research Assistant, Wandsworth Hospital Group

AND

NORMAN C. TANNER, F.R.C.S.

Senior Surgeon, St. James's Hospital, London

Among those patients who are referred to hospital with a complaint of dyspepsia there remain after routine investigation an appreciable number in whom no satisfactory diagnosis is reached despite the fact that they give the impression of having organic disease. Such cases are usually labelled functional or nervous dyspepsia, or, if there is no free hydrochloric acid secreted in response to a test meal, achlorhydric dyspepsia. Gastroscopic examination will provide a satisfactory explanation of the symptoms in a large proportion of cases. Not only may small ulcers and ulcer scars which have been missed at a barium meal x-ray examination be discovered by this means, but it is also the most satisfactory method of confirming the presence of and differentiating gastritis. Where gastritis is found it is, however, impossible to give a reliable prognosis regarding the future comfort of the patient or to assess the possibilities of complications such as bleeding, ulceration, or the development of gastric carcinoma. As many patients have gastric changes without dyspepsia it is also impossible to be dogmatic on how much the patients' symptoms are attributable to the gastric condition.

Schindler (1947) has given a good working classification of chronic gastritis. He divides cases into four types—hypertrophic, superficial, and atrophic gastritis, and gastritis in the post-operative stomach.

Chronic hypertrophic gastritis seems to be a very rare condition in this country and is not to be confused with the simple hypertrophy with thick folds and prominent areae gastricae which is a normal variant and is the usual finding in patients with hyperchlorhydria and duodenal ulcer. In hypertrophic gastritis intra-

mucosal scarring has led to unevenness of the mucosal pattern, and there may even be pseudo-polypoid changes.

In chronic superficial gastritis there is uneven congestion of the mucosa and excess adherent mucus. Simple diffuse reddening of the mucosa is a physiological state, whereas a stippled or variegated appearance is pathological. Non-adherent mucus is a normal exudate and mucopus is usually swallowed sputum.

The atrophic group is characterized by thinning of the gastric folds, inability to distinguish the areae gastricae, and, when atrophy is marked, by the ability to see the intramucosal and submucosal vessels through the thinned mucosa (Fig. 1). There is often an element of inflammation, shown by congestion along the tips of the folds and by stippling of the mucosa. The appearance of gastric atrophy may be generalized or patchy, and a localized variety is not infrequently seen in the vicinity of ulcer scars. This latter is in reality a new thin mucosa or a stretched mucosa around a scar (Fig. 2).

FIG. 1.—Gastroscopic view of atrophic gastritis, showing visible submucosal vessels.

FIG. 2.—Gastroscopic view, showing atrophic gastritis with vessels more prominent in the vicinity of a healed gastric ulcer. Either a true adenoma is present or there is pseudo-polypoid change as a result of scarring. The folds towards the greater curve are normal.

Present Investigation

The case summaries of all patients attending the Gastroenterological Clinic, St. James's Hospital, London, during the years 1940–6 inclusive have been studied. Where a finding of achlorhydria on a first fractional test meal using gruel as the stimulant was recorded and where some address for the patient was given, the case reports have been abstracted. Certain groups have been followed up during 1952, giving minimum and maximum observation periods of five and twelve years.

These patients are almost without exception those with gastro-intestinal symptoms. It is well known that the

absence of free hydrochloric acid need not produce symptoms. It occurs in a proportion of otherwise normal people, it is an invariable accompaniment of pernicious anaemia, and is common in such diverse conditions as rheumatoid arthritis, steatorrhoea, and various chronic skin diseases.

The records of 403 patients with achlorhydria have been found, and these have been grouped as follows:

Carcinoma of the stomach	63
Pernicious anaemia	15
Advanced atrophy without pernicious anaemia	113
Superficial gastritis and early atrophy	126
Chronic duodenal ulcer	13
Normal gastroscopy—no other cause found	21
No gastroscopy—no other cause found	52

These figures give no true indication of the incidence of the different conditions. Superficial gastritis and early atrophy are not necessarily associated with achlorhydria. Gruel is a comparatively weak stimulant, and had it been combined with an injection of histamine some cases would certainly have secreted hydrochloric acid. In some cases the finding was either false or only temporary. Thus, in the duodenal ulcer group, although the majority had pyloric stenosis with retention gastritis, there were some in whom the tube was evidently not in the stomach. Such cases can usually be eliminated, as no starch can be found in any sample of the gastric aspiration. In the present series, however, we have not been able to exclude them, since a number were referred from other hospitals merely for gastroscopy, and no detailed reports of the analysis are available.

Some of the groups can be disposed of rapidly. Active duodenal ulcers are never associated with achlorhydria, except when complicated by retention gastritis. In this paper we are not concerned with those patients who presented initially with a gastric carcinoma, and this group is not analysed.

The gastroscopic appearances of patients with pernicious anaemia are identical with those of atrophic gastritis. Weighty evidence suggesting that pernicious anaemia is a precursor of gastric carcinoma has been advanced by Kaplan and Rigler (1945, 1947) and Rigler, Kaplan, and Fink (1945), but confirmation of their conclusions must be awaited, since Wilkinson (1945), in a series of 1,600 patients with pernicious anaemia, was unable to find any such association. In the present series there are only 15 patients. The majority did not have gastric symptoms. Nine are still alive, four are dead (one from gastric carcinoma two and a half years after gastroscopic discovery of atrophic changes), and the remaining two have not been traced.

The bulk of the paper is devoted to the superficial gastritis, the early atrophic, and the advanced atrophic groups associated with achlorhydria. The remaining two—those with normal gastroscopic findings and those in whom no gastroscopy was performed—are referred to only where they have something of interest to offer.

Superficial Gastritis; Gastric Atrophy

Patients with advanced atrophy are easily separated from the remainder by reason of the presence of visible sub-mucosal vessels on gastroscopy. In the case of chronic superficial gastritis and early atrophy, however, there seems to be no sharp dividing line, and, at any rate in the presence of achlorhydria, the two conditions overlap. For this reason no attempt has been made to separate them, and they are referred to hereafter as the early atrophic group. A further complication in analysing these patients is that at any stage acute gastric erosions, acute, subacute, or chronic gastric ulcers, or old ulcer scars may occur. The presence of these alters the clinical findings to some extent, and it will often be necessary to separate the cases from their parent groups. There are a few cases of chronic gastric ulcer with achlorhydria in which the gastric mucosa was reported on as normal. Their inclusion in this analysis should not affect the results, as the chronic gastric ulcer cases are dealt with separately where they depart from the

general run of the findings. Patients in whom the only evidence of advanced atrophic changes was in the vicinity of ulcer scars are placed in the early group.

The Atrophic Groups

The data concerning the age and sex of the patients are outlined in Table I.

TABLE I.—*Atrophic Gastritis—Age and Sex Incidence*

Age Group	Early Atrophy			Advanced Atrophy		
	Men	Women	Total as %	Men	Women	Total as %
Under 30 ..	1	5	4.8	—	3	2.7
30- ..	13	3	12.7	8	3	9.7
40- ..	13	11	19.0	11	9	17.7
50- ..	20	17	29.4	21	18	34.5
60- ..	17	11	22.2	19	14	29.2
70-79 ..	6	7	10.3	4	2	5.3
Unknown ..	1	1	1.6	1	—	0.9
Total ..	71	55		64	49	

The ratio of men to women is about four to three in each group of the patients, but further division (Table II) suggests that this inequality may be due to the inclusion of patients with gastric ulcers.

TABLE II.—*Atrophic Gastritis—Sex Incidence*

	Early Atrophy		Advanced Atrophy		Combined	
	Men	Women	Men	Women	Men	Women
Acute erosions ..	7	10	6	13	13	23
Acute ulcers ..	6	3	6	3	12	6
Subacute ..	1	2	2	1	3	3
Chronic ..	27	7	7	4	34	11
Ulcer scars ..	9	9	4	2	13	11
Remainder ..	21	24	39	26	60	50
Total ..	71	55	64	49	135	104

The acute-erosions group has been taken to include not only cases in which an erosion was actually seen, but also those with manifest haemorrhage in which no lesion other than gastritis was found. Table II shows that these erosions were nearly twice as common in women as in men, while chronic gastric ulcers were three times more common in men than in women. If the erosion cases are kept with their parent groups and the ulcer cases are not included there is no great difference in the sex incidence.

The age distribution in the various subgroups is approximately the same, the highest incidence being in the fifties. As can be seen from Table I, there is a wide distribution, the youngest patient in the advanced atrophy group being a 17-year-old girl and the oldest a 79-year-old man. If the population at risk is taken into account there is probably an increasing incidence with increasing age.

TABLE III.—*Atrophic Gastritis—Age at Onset of Symptoms*

Age Group	Ulcers + Atrophy	Remainder	
		Early Atrophy	Advanced Atrophy
10- ..	1 (1.2%)	1 (1.7%)	4 (5.4%)
20- ..	9 (10.8%)	4 (6.9%)	6 (8.1%)
30- ..	15 (18.1%)	13 (22.4%)	15 (20.3%)
40- ..	26 (31.3%)	11 (19.0%)	14 (18.9%)
50- ..	21 (25.3%)	14 (24.1%)	20 (27.0%)
60- ..	9 (10.8%)	11 (19.0%)	10 (13.5%)
70-79 ..	2 (2.4%)	4 (6.9%)	5 (6.8%)
Total ..	83	58	74

Patients in whom the age at onset was unknown have been omitted.

The onset of symptoms is commonest between the ages of 30 and 60, with a maximum incidence in the fifties (Table III). There is little difference between the early and late groups. Where there is a concomitant ulcer the distri-

bution resembles more closely that of chronic gastric ulcer, with a maximum incidence in the forties.

The data concerning the length of history of symptoms are set out in Table IV, where it has been necessary to subdivide the groups. A greater number of patients with

TABLE IV.—*Atrophic Gastritis—Length of History of Symptoms*

Length of History (Years)	Combined Groups				Remainder	
	Erosions	Acute and Subacute Ulcers	Chronic Ulcers	Scars	Early	Advanced
Under 1 ..	13 (38.2)	9 (39.1)	11 (28.2)	6 (28.6)	16 (38.1)	17 (30.4)
1- ..	8 (23.5)	4 (17.4)	12 (30.8)	7 (33.3)	14 (33.3)	12 (21.4)
5- ..	4 (11.8)	3 (13.0)	2 (5.1)	1 (4.8)	6 (14.3)	10 (17.9)
10- ..	5 (14.7)	4 (17.4)	8 (20.5)	6 (28.6)	5 (11.9)	8 (14.3)
Over 20 ..	4 (11.8)	3 (13.0)	6 (15.4)	1 (4.8)	1 (2.4)	9 (16.1)
Total ..	34	23	39	21	42	56

Patients in whom the length of history was unknown have been omitted.

early atrophy present with a short history of symptoms than do those with advanced atrophy. This is not unexpected, since atrophy is presumably a progressive process. Indeed, it is probable that gastritis is a degenerative rather than a true inflammatory condition. However, nearly a third of those with advanced atrophy have a history of under one year's duration. Possibly some added factor such as inflammation, an exogenous irritant, or a transitory erosive process is required for the production of symptoms.

It has been suggested that the low acid values associated with chronic gastric ulcer are secondary to the presence of the ulcer. Table IV suggests that this is not necessarily so. In a series of 498 patients with chronic gastric ulcers previously analysed, about 40% gave a history of symptoms of under five years' duration, while in the present series the figure is 60%. This increase is explained by the inclusion of patients who were found at other hospitals to have a short history of symptoms combined with achlorhydria and an ulcer crater, and who were referred here for gastroscopic examination to exclude malignancy. This does not interfere with the finding that low acid values in chronic gastric ulcer are often associated with a long history of ulcer symptoms, and we again stress that chronic gastric ulcer is in most cases probably the outcome of degenerative changes in the gastric mucosa (Tanner, 1951).

Symptomatology of the Atrophic Groups

Epigastric Pain and Discomfort

Over 90% of patients complained of epigastric pain or discomfort, the most common form being pain related to meals (Table V).

TABLE V.—*Chronic Gastritis—Pain Related to Meals*

	Pain Related to Meals	Remainder
Early atrophy, including erosions ..	37 (59.7%)	25 (40.3%)
Late ..	47 (56.0%)	37 (44.0%)
Acute and subacute ulcers ..	14 (58.3%)	10 (41.7%)
Chronic ulcers ..	36 (81.8%)	8 (18.2%)
Ulcer scars ..	10 (43.5%)	13 (56.5%)

Patients in whom the symptomatology was unknown are not included.

The erosion cases have been kept with their parent group. If they are separated the findings are not materially altered. The patients with chronic gastric ulcer are more likely to complain of post-prandial pain than the remainder, but of those without a demonstrable break in the continuity of the gastric mucosa over half complain of this symptom. The mode of production of the pain is intriguing and obscure. It appears to differ little from that accompanying peptic ulceration, though in the late atrophic group it seems to come on more often directly after eating. This is not always so, but the details of the time relationship to meals are insufficient for statistical analysis.

In some patients the pain is relieved by alkalis (it is well known that many achlorhydric dyspeptics complain of "acidity"). In some cases vomiting, and in others taking food, gives relief; some find that nothing helps. Relief from taking food follows the general rule that the longer the interval between eating and the onset of pain the more likely is food to afford relief.

Comparatively few patients in whom ulcer scars in addition to atrophy were found complained of post-prandial pain, since these patients not infrequently present with manifest haemorrhage without a previous history of dyspepsia.

Only 8% of patients with chronic ulcers (compared with 18% in the early atrophic and 25% in the late atrophic groups) complained of pain unrelated to meals.

The symptoms of fullness or discomfort after meals were rarely prominent in the chronic ulcer patients, but were present in 18% of the early atrophic and 15% of the late atrophic parent groups.

Manifest Haemorrhage

It has been found in this clinic that 50% of patients with haematemesis or melaena in whom no gross cause is found for the bleeding have gastroscopic evidence of atrophic changes (Tanner, 1949). The data in the present series are set out in Table VI.

TABLE VI.—*Chronic Gastritis—Manifest Haemorrhage*

	History of Haemorrhage	No History of Haemorrhage
Early atrophy, including erosions ..	17 (27.4%)	45 (72.6%)
Late ..	18 (21.4%)	66 (78.6%)
Acute and subacute ulcers ..	13 (54.2%)	11 (45.8%)
Chronic ulcers ..	11 (25.0%)	33 (75.0%)
Ulcer scars ..	10 (43.5%)	13 (56.5%)

TABLE VII.—*Haemorrhage*

	Haemorrhage with Preceding Dyspepsia	Haemorrhage as Sole Symptom
Early atrophy, including erosions ..	13	4
Late ..	14	4
Acute and subacute ulcers ..	11	2
Chronic ulcers ..	11	—
Ulcer scars ..	6	4

There is a history of haemorrhage in about a quarter of the cases of early and late atrophy, but when these are complicated by acute or subacute ulcers or ulcer scars the figure is doubled (Table VI). The incidence where there is an additional chronic ulcer is not higher than for the parent groups, and is approximately the same as in patients with chronic ulcers without achlorhydria (Swynnerton and Tanner, 1953).

In an appreciable number of cases haemorrhage may be the only symptom (Table VII).

In some cases an obvious lesion may be found from which the bleeding comes, but in others there may be no obvious bleeding point even in those rare cases which continue to bleed and come to necropsy. However, one of us has seen acute erosions overlying visible submucosal vessels (Tanner, 1949), and it may be that these patients are more

FIG. 3.—Gastroscopic view of an acute erosion, overlying a visible submucosal vessel.

susceptible to mucosal erosion than normal and a shallow erosion into a submucous vessel may cause copious haemorrhage (Fig. 3).

Other Symptoms

Nausea, Vomiting, Anorexia, Weight Loss.—The symptoms of nausea and vomiting, anorexia, or weight loss are present in approximately a fifth of the patients (Table VIII).

TABLE VIII.—*Nausea and Vomiting, Anorexia, Weight Loss*

	Nausea and Vomiting	Anorexia	Weight Loss
Early atrophy, including erosions	12 (19.4%)	11 (17.7%)	14 (22.6%)
Late " "	16 (19.0%)	13 (15.5%)	19 (22.6%)
Acute and subacute ulcers	2 (8.3%)	4 (16.7%)	7 (29.2%)
Chronic ulcers	6 (13.6%)	6 (13.6%)	10 (22.7%)
Ulcer scars	1 (4.3%)	5 (21.7%)	6 (26.1%)

Note: The totals on which percentages are based may be obtained by reference to Tables V or VI.

The presence of an ulcer does not greatly affect the incidence except in the case of nausea and vomiting, which is less frequent when an ulcer is present. So far as possible those patients who vomited merely in an attempt to relieve their symptoms and those who were afraid to eat have been omitted.

Bowel Habit.—Complaints of diarrhoea or constipation as prominent symptoms were uncommon, the maximum incidence being in the parent late atrophic group, in which one in ten complained of diarrhoea and one in twenty of constipation. Bowel habit is dealt with more fully in the late analysis.

Lassitude and anaemic symptoms were complained of by less than one in ten of the patients. Other more uncommon main complaints included belching, regurgitation, and dysphagia.

Conclusions on Symptomatology

Over 90% of the patients complained of epigastric pain or discomfort, most often related to meals. One in five was troubled with vomiting, anorexia, or weight loss; and one in three had manifest haemorrhage at some stage during the course of the disease. To sum up, there is no definite clinical picture to distinguish chronic atrophic gastritis from peptic ulceration or gastric carcinoma, and it may at times mimic gall-bladder disease, hiatus hernia, or many other disorders.

Associated Conditions

Gastric Adenoma.—In five of the 113 patients with advanced atrophy there were coexistent gastric adenomata—in two a single adenoma was seen; in one there were two adenomata, and in two the adenomata were multiple. In the early atrophic group one woman had two adenomata, and in one man a doubtful adenoma was seen at a later date.

Gall-bladder Diseases.—Six of the 113 patients in the advanced atrophic and 7 of the 126 in the early atrophic groups had a past, concurrent, or subsequent history of gall-bladder disease. Since the two conditions may be indistinguishable clinically, mistakes concerning the causation of symptoms are bound to occur. Thus patients who have had a cholecystectomy may return to hospital with their symptoms unaltered, while others in whom both conditions are found at the same admission may be permanently relieved by cholecystectomy. It is even possible that the dyspeptic symptoms of gall-bladder disease (not, of course, colic) may be due to a coincident gastritis or duodenitis, for it is well known that flatulent dyspepsia may persist after cholecystectomy.

Chronic Respiratory Infections.—Fourteen patients in the late and 10 in the early groups suffered from chronic bronchitis. It is not possible to say whether the two conditions are associated, since chronic bronchitis is a common condition in South London, but an incidence of 10% would seem to be high. Four patients in the late group have been diagnosed as suffering from pulmonary tuberculosis, and

another gave a past history of tuberculous peritonitis. In the early atrophic group two gave a history of pulmonary tuberculosis.

Tertiary Syphilis.—Three patients in the late and two in the early groups had tertiary syphilis. Three of these had syphilitic aortitis and the other two had tabes dorsalis. In one of the former, submucous gastric swellings were seen on gastroscopy, and these disappeared after antiluetic treatment.

Other Associated Conditions.—These include psoriasis (2), chronic dermatitis (1), osteoarthritis (3), rheumatoid arthritis (1), haemophilia (1), cystinuria (1), and diabetes mellitus (1).

Late Results

The method of follow-up of patients was the same as that previously described (Swynnerton and Tanner, 1953). Results are rather less complete than in the follow-up of cases of chronic gastric ulcer, for two reasons: (1) that the records of addresses and next-of-kin were often less detailed; and (2) that four of the patients referred from other hospitals were war refugees from Malta and Gibraltar whom it has proved impossible to trace. The general results of the follow-up investigation are given in Table IX.

TABLE IX.—*Atrophic Gastritis—General Results of Follow-up*

	Early Atrophy	Late Atrophy
Alive—adequate information	78 (61.9%)	82 (72.6%)
" —inadequate "	2 (1.6%)	1 (0.9%)
Dead	38 (30.2%)	22 (19.5%)
No information	8 (6.3%)	8 (7.1%)
Total	126	113

Deaths

The causes of death in the early and late atrophic groups, in those with a gastroscopically normal mucosa, and in those in whom no gastroscopy was performed are shown in Table X.

TABLE X.—*Causes of Death*

	Late Atrophy	Early Atrophy	Normal Mucosa	No Gastroscopy
Gastro-intestinal:				
Gastric carcinoma	—	3	—	1
" ulcer	—	2	—	—
" haemorrhage	1	—	2	—
Carcinoma of large bowel	1	1	—	1
Sarcoma of ileum	—	—	—	1
Carcinoma of gall-bladder	—	2	—	—
" pancreas	—	—	1	—
Appendicitis	—	1	—	—
Mediastinitis	—	1	—	—
Cardiovascular:				
Heart failure	7	13	2	3
Syphilis	1	1	1	—
Cerebral vascular accidents	3	2	1	1
Respiratory:				
Chronic bronchitis	—	3	2	1
Acute infections	2	2	—	—
Tuberculosis	4	2	1	1
Carcinoma of bronchus	1	1	—	—
Renal:				
Pyelonephritis	1	1	—	1
Chronic nephritis	—	—	—	1
Other:				
Aplastic anaemia	—	—	1	—
Accidents	—	1	—	—
Suicide	1	—	—	—
Unknown	—	2	—	2
Total	22	38	11	13

The figures in the early atrophic group again draw attention to the fact that the dyspepsia may be due to causes other than the gastritis, and one should not be satisfied with a single diagnosis of chronic gastritis before excluding all other possibilities. Thus the two patients with carcinoma of the gall-bladder both died less than two months after gastritis had been found. It would seem possible, too,

that in a number of those dying of "heart failure" the achlorhydria and gastroscopic appearances may have been secondary to chronic cardiac disease.

The average age at death in the advanced atrophic group was 67.6 years, only two patients having died under the age of 60—one of tuberculosis and the other of syphilitic aortitis. Of those patients still alive at least one is over 90. In the early atrophic group the average age at death was 65.

In the advanced atrophic group, apart from the one patient who died of uncontrolled gastric haemorrhage, no patient appears to have died either as a direct or as an indirect result of atrophy.

Carcinoma of Stomach

In the late atrophic group none of the traced patients have developed a gastric carcinoma and only two of them have developed a malignant tumour elsewhere in the body. In some 3% of all people the cause of death is attributed to gastric carcinoma (Registrar-General, 1953), so it is reasonable to suppose that if atrophic gastritis is a precursor of gastric carcinoma one might find a certain number of cases in the present series. The essentially negative findings are in line with the only other series of gastroscopically confirmed cases found in the literature. Findley *et al.* (1950) followed for five years or more 100 patients, of whom 49 had "mild," 33 "moderate," and 18 "pronounced" atrophy, and found none who developed gastric carcinoma.

In the early atrophic group four patients have developed gastric carcinoma and seven a carcinoma elsewhere in the body. The former are analysed further:

Case 1.—A man aged 52 was admitted on December 5, 1941, with an eighteen-months history of epigastric pain immediately after meals. Fractional test meal analysis showed achlorhydria. Gastroscopic examination revealed no gross abnormality apart from fine congestion of the mucosa. Barium-meal examination revealed no lesion. He was reinvestigated in 1943 and 1944 on account of recurrent symptoms. On both occasions a barium-meal examination was normal. In 1943 gastroscopy showed thinning of the mucosa without visible vessels and a small adenoma. This latter was not seen in 1944. He was seen at the follow-up clinic in April, 1952, when he still complained of similar symptoms. On examination a large epigastric mass was palpated. He was admitted and died following removal of a large adenocarcinoma of the body of the stomach.

Case 2.—A man aged 70 was admitted on February 3, 1942, with a two-years history of epigastric pain two to three hours after meals associated with anorexia and vomiting. Fractional test meal analysis showed achlorhydria. Gastroscopy revealed marked congestion of the mucosa and a chronic gastric ulcer at the angulus. In view of his general condition a gastro-jejunostomy instead of a partial gastrectomy was performed after failure of the ulcer to heal on medical treatment. He continued to have recurrent dyspepsia, and the ulcer was again seen gastroscopically in August, 1944. He was readmitted in May, 1947, when he had an epigastric mass, a palpable irregular liver, and ascites. He died a few days later. Permission for a necropsy was refused.

Case 3.—A woman aged 73 was admitted on February 17, 1945, with a four-years history of vague epigastric pain and recent lassitude. A barium-meal examination showed no lesion. Fractional test meal analysis showed achlorhydria. Gastroscopy revealed a mucosa which was thin in patches and associated with speckled congestion. For technical reasons the examination was incomplete, and it was thought that there might be a lesion at the angulus. She was readmitted in May, 1948, with a six-months history of pallor, lassitude, and anaemic symptoms. There was no further mention of dyspepsia. Severe hypochromic anaemia was found. During her stay in hospital she developed sudden pain in the epigastrium and died on June 9, 1948. At necropsy a small flat carcinoma was found arising $\frac{1}{2}$ in. (1.3 cm.) proximal to the pylorus. In addition there was a subphrenic abscess.

Case 4.—A man aged 51 was admitted on August 25, 1941, with haematemesis preceded by a week of epigastric discomfort after meals. Fractional test meal analysis showed achlorhydria. Barium-meal examination showed an innocent ulcer on the lesser curve and this was confirmed gastroscopically. The gastric mucosa was reported to be otherwise normal. The ulcer was

shown to be soundly healed seven weeks later. He returned in May, 1946, with a history of recurrence of dyspepsia for the past year. On gastroscopy the ulcer was thought to be innocent, but at operation and on section it was found to be a diffuse infiltrating spheroidal-cell carcinoma.

Apart from the presence of achlorhydria there is no factor common to these four cases. In three there was mucosal congestion, in two an apparently innocent chronic gastric ulcer, and in two evidence of atrophy of the mucosa. It is not possible to reach any conclusion on the study of these cases.

Pernicious Anaemia

One patient, a 69-year-old man, was diagnosed as suffering from pernicious anaemia three years after the initial gastroscopic diagnosis of advanced atrophy. Two other patients gave family histories of pernicious anaemia.

Late Results in Patients Still Alive

In the advanced atrophic group eight patients have had partial gastrectomies performed—seven for chronic ulcer and one for gastric polyposis—leaving 74 for late assessment. The early atrophic group is loaded with patients with concurrent chronic gastric ulcers, and, since these ulcers are just as apt to recur as those in patients without achlorhydria, it has been decided to omit them from the late analysis. Fifty patients remain alive and were followed up. The subsequent course of events in the two series is shown in Table XI.

TABLE XI.—Chronic Gastritis—Late Results

	Early Atrophy Excluding Chronic Ulcers		Late Atrophy		Total	
	No.	%	No.	%	No.	%
No further trouble	13	(26.0)	18	(24.3)	31	(25.0)
Occasional mild symptoms ..	16	(32.0)	21	(28.4)	37	(29.8)
" severe	1	(2.0)	3	(4.1)	4	(3.2)
Frequent or constant symptoms	20	(40.0)	32	(43.2)	52	(41.9)

The results in the two series are very similar. Just over half the patients have had little or no further trouble from their stomach. Chronic superficial gastritis and the earlier stages of atrophy are conditions which may heal completely (Schindler, 1947), and, as we know from patients with pernicious anaemia, advanced atrophy need not cause symptoms.

Nearly all the patients in the advanced and many in the early atrophic group were advised to take a mixture containing hydrochloric acid, but only 11 have continued to do so, while 59 take alkalis more or less regularly.

The reason for this swing from acid to alkalis is three-fold. First, patients find hydrochloric acid unpleasant to take, no matter how much diluted or disguised; secondly, dyspepsia means to general practitioners a prescription for alkalis, and the recommendation to prescribe acid is soon forgotten; and, thirdly, many patients find that alkalis are effective for relieving their symptoms. A wide variety of alkalis were used with good effect.

The bowel habits of 39 patients with early atrophy and 59 with advanced atrophy are known (Table XII). Constipation is common in the advanced atrophic group, while diarrhoea, either in attacks or as a constant looseness of the motions, is present in one in ten.

TABLE XII.—Bowel Habit in Patients with Atrophic Gastritis

	Early Atrophy	Advanced Atrophy
Normal	22 (56.4%)	24 (40.7%)
Constipation	12 (30.8%)	27 (45.8%)
Diarrhoea	5 (12.8%)	6 (10.2%)
" or constipation ..	0 (0.0%)	2 (3.4%)

Discussion

The diagnosis of chronic gastritis should be accepted as the cause of the patient's symptoms only after exclusion of all other possible conditions. Confirmation of its presence will be obtained only by gastroscopic examination. Blind biopsy methods such as that described by Wood *et al.* (1949) may help to elucidate the process of gastritic changes, but will be unreliable for diagnosis, as the condition is often patchy.

The prognosis regarding symptoms is somewhat better than in the medical treatment of chronic gastric ulcer, since about half the patients have very little subsequent trouble. Amongst the other half there are a few who become disgruntled, finding that different regimes of treatment have little effect on their symptoms. A dietary regime similar to that used in patients with chronic gastric ulcer should be followed, and during exacerbations bed rest will undoubtedly hasten relief and may be followed by long remissions. A mixture containing hydrochloric acid should be prescribed, to be taken well diluted during meals, and the patient should be allowed to experiment until he finds the most effective flavouring agent, which is most likely to be one of the citrus fruit juices (Bockus, 1944). Hydrochloric acid therapy is usually sufficient to control attacks of achlorhydric diarrhoea, but in obstinate cases it may be supplemented with phthalyl or succinylsulphathiazole. A few patients are of the opinion that acid prevents or terminates attacks of dyspepsia, but this does not appear to be the general finding. If hydrochloric acid is badly tolerated or fails to relieve the symptoms when taken regularly over a period of time, one of the non-absorbable alkalis may be tried, or "autolavage" by taking 1 dr. (4 g.) of sodium bicarbonate in water before meals, as recommended by Spriggs and Marxer (1943).

Supplementary vitamin-B complex is often given to these patients, since there is evidence that a deficiency leads to a decrease in gastric acidity. It has been reported by several observers, notably by Jones, Benedict, and Hampton (1935), that in patients with pernicious anaemia there may be regeneration of the gastric mucosa with liver therapy, but this has been denied by others who regard advanced atrophy as an irreversible state. In this clinic one of us (N. C. T.) has seen a patient with pernicious anaemia whose gastric mucosa showed patchy regeneration on liver therapy. Until there is more evidence on the action of liver extract and vitamin B₁₂ on the stomach it would seem to be inadvisable to use them in simple atrophic gastritis for fear of masking the development of pernicious anaemia. Surgery appears to play little part in the treatment of atrophic gastritis. As Schindler (1947) remarks, "there is no reason to believe or to hope that the pain will be alleviated by surgical interference." Patients who have had partial gastrectomies performed for a coexistent ulcer or ulcer scar are not infrequently seen with their symptoms unaltered. In cases of uncontrolled haemorrhage a total gastrectomy would have to be contemplated to guarantee success, since the bleeding areas are often multiple and widely scattered, but in practice a partial resection usually leads to diminution and sometimes to cessation of bleeding.

Opinions concerning the role of chronic gastritis as a precursor of gastric carcinoma have been as much at variance as in the more familiar "ulcer-cancer" controversy. The histological examination of stomachs resected for cancer provides little evidence, for, even if there were general agreement on the degree of gastritic changes, it would still be impossible to decide whether these were the primary condition or whether they were secondary to the presence of the neoplasm. The solution of the problem is therefore a clinical one. The most important work done from the clinical aspect has been that of Comfort and his colleagues (1947) at the Mayo Clinic. They have studied a series of patients with gastric carcinoma who had had gastric analyses performed before the development of the carcinoma. They found that the proportion of low or

absent free hydrochloric acid values was greater than normal in these patients, suggesting that carcinoma arises in an already diseased stomach. *It must be borne in mind, however, that this was a selected series of patients, since they were those in whom the carcinoma was preceded by many years of dyspepsia, and it is not possible to estimate the importance of chronic gastric disease in the development of gastric carcinoma as a whole.*

In the present investigation none of the traced patients with advanced atrophic changes have developed gastric cancer. This is negative evidence which is suggestive but by no means conclusive, and, ideally, all these patients should be seen again in another five to ten years. The negative findings are to some extent supported by the sex incidence, which is not that of gastric carcinoma. It is now well established that gastric adenomata are prone to become malignant, and Schindler (1947) has reported that these adenomata are seven times more common in people with gastric atrophy than in those without. The adenomata are presumably secondary to or a variant of the atrophic changes and may form a connecting-link between the atrophy and the development of carcinoma. In the early atrophic series, in which four patients developed gastric carcinoma, it is not possible to estimate whether this is a significant incidence, and the factors involved were multiple. In only two was there macroscopic evidence of atrophy.

It will have been noticed that in 11 of the cases of advanced atrophy there were concomitant chronic benign gastric ulcers. It is usually taught that a diagnosis of gastric ulcer is incompatible with a finding of complete achlorhydria. In some of the present cases the atrophy was patchy, but in others it was generalized, and in these the capacity of the stomach to secrete acid in response to any stimulant must be negligible. It seems probable that hydrochloric acid is essential for the initial formation of the ulcer, but not for its maintenance. This latter may be due to poor healing powers in the degenerate atrophic mucosa and also, possibly, to superadded infection or trauma.

Summary

An analysis and late follow-up has been carried out on a series of patients who were found to secrete no free hydrochloric acid in response to a test meal using gruel as the stimulant. Two main groups of gastroscopically confirmed patients have been dealt with in some detail, the one comprising 113 cases of advanced atrophic gastritis, and the other 126 cases of chronic superficial gastritis or early atrophic gastritis.

The analysis is complicated by the frequent co-existence of gastric erosions or ulcers.

If patients with chronic gastric ulcers are excluded there is no great difference between the incidence in the two sexes.

There is a wide distribution in the age incidence, the majority of patients in the two groups presenting at hospital between the ages of 40 and 70. Information is given on the length of history and age at onset of symptoms.

Over 90% complain of epigastric pain or discomfort, which is usually related to eating.

Over a quarter give a history of manifest haemorrhage, and in one in five of these haemorrhage is the sole symptom. Complaints of nausea, vomiting, anorexia, and weight loss are frequent.

Less common main symptoms include diarrhoea, constipation, anaemia, belching, regurgitation, and dysphagia.

In the advanced atrophic group constipation is four times more common than diarrhoea.

Of associated conditions gall-bladder disease seems to be most important, since it may be clinically indistinguishable from atrophic gastritis, and since there may be direct association between the two. Other associated conditions which may be of aetiological significance include chronic bronchitis, pulmonary tuberculosis, tertiary syphilis, cardiovascular disease, and certain chronic skin and joint conditions.

In the late atrophic group 22 patients (19.5%) have died. In only one, a case of uncontrolled gastric haemorrhage, was there any connexion between the atrophic gastritis and the cause of death. No patient in this group has developed a gastric carcinoma.

In the early atrophic group 38 patients (30.2%) have died. Four of these have developed a gastric carcinoma, but, apart from the absence of free hydrochloric acid, no common factor could be found.

One patient with advanced atrophy was subsequently found to have developed pernicious anaemia.

Among those patients still alive the results in the two groups are very similar. About half have had very little subsequent trouble, but the course of the disease seems to have been little influenced by a variety of regimes in the remainder.

The treatment of atrophic gastritis is discussed.

Some of the evidence concerning the role of chronic gastritis as a precursor of gastric carcinoma is presented, and the apparently anomalous finding of peptic ulcer of the stomach in the absence of free hydrochloric acid is discussed.

We are greatly indebted to the many people, not least the patients themselves, who co-operated in the follow-up. We wish to thank Miss Temple and Mrs. Strong, of the Records Department, St. James's Hospital, and Miss Kirwin for the many hours spent unearthing ancient records, and the latter for her secretarial assistance. The cost of the investigation was met from a grant from the Weir Research Fund.

REFERENCES

- Bockus, H. L. (1944). *Gastroenterology*, vol. 1. Saunders, Philadelphia and London.
 Comfort, M. W., Kelsey, M. P., and Berkson, J. (1947). *J. nat. Cancer Inst.*, 7, 367.
 Findley, J. W., Kirsner, J. B., and Palmer, W. L. (1950). *Gastroenterology*, 16, 347.
 Jones, C. M., Benedict, E. B., and Hampton, A. O. (1935). *Amer. J. med. Sci.*, 190, 596.
 Kaplan, H. S., and Rigler, L. G. (1945). *Ibid.*, 209, 339.
 ——— (1947). *J. nat. Cancer Inst.*, 7, 327.
 Registrar-General (1953). Statistical Review for England and Wales for the Year 1951. Part 1. H.M.S.O., London.
 Rigler, L. G., Kaplan, H. S., and Fink, D. L. (1945). *J. Amer. med. Ass.*, 128, 426.
 Schindler, R. (1947). *Gastritis*. Heinemann, London.
 Spriggs, E. I., and Marxer, O. A. (1943). *Quart. J. Med.*, 36, 1.
 Swynnerton, B. F., and Tanner, N. C. (1953). *British Medical Journal*, 2, 841.
 Tanner, N. C. (1949). *Med. Press*, 221, 3.
 ——— (1951). *Med. Wld. Lond.*, 75, 455.
 Wilkinson, J. F. (1945). *British Medical Journal*, 2, 664.
 Wood, I. J., Doig, R. K., Motteram, R., and Hughes, A. (1949). *Lancet*, 1, 18.

The latest issue of *Scientific Film Review* includes appraisals of five medical films. "Coarctation of the Aorta" (E. R. Squibb & Sons), intended for students and society meetings. "Haemolytic Disease of the Newborn" (St. James's Film Unit) depicts the technique of replacement transfusion. "Marrow Puncture" (I.C.I.) surveys the techniques of marrow puncture in patients of various ages. It is intended for housemen and postgraduate students. "Venepuncture" (I.C.I.) demonstrates the technique in adults. These four films are in colour. A black-and-white film mentioned is "Prevention of Infection of Babies in Hospital" (British Film Institute), a comprehensive survey of modes of spread of infection in children's wards. It is for the training of nurses and medical students entering paediatric wards. *Scientific Film Review* is published quarterly by the British Film Institute at an annual subscription of 5s.

RESULTS OF TREATMENT OF CHRONIC VASOMOTOR RHINITIS

PART I. RESPONSE TO PHYSICAL TREATMENT

BY

PHILIP READING, M.S., F.R.C.S.

Surgeon, Throat and Ear Department, Guy's Hospital

AND

KENNETH MALCOMSON, F.R.C.S.

Registrar, Throat and Ear Department, Guy's Hospital

Sufferers from chronic vasomotor rhinitis form a large proportion of all those attending the throat clinics of most hospitals, and we have been impressed not only by the frequency of the disorder but also by the unsatisfactory response to treatment. In addition it is apparent that many of these patients are emotionally upset, and that their disabilities are in part caused and in part magnified by psychological imbalance.

Results of Physical Investigation and Treatment

A series of 127 consecutive patients were studied. Symptoms, rhinological appearances, allergic responses, and the results of treatment were noted, and a superficial assessment of the emotional background was attempted in the throat clinic (by K. M.). As many as possible were invited to participate in more detailed psychological investigation, and those who did attend may fairly be regarded as a representative sample of sufferers from vasomotor rhinitis.

The main symptom in this disease is either paroxysmal rhinorrhoea or persistent nasal blockage. While well-defined examples of each may be recognized, yet in a great number the two clinical pictures are merged, to give symptoms and signs of both. In this series, there were 46 cases of paroxysmal sneezing and rhinorrhoea and 81 of chronic nasal oedema (49 with polypi and 32 without). A brief recapitulation of the two syndromes is appended.

1. Non-specific Paroxysmal Sneezing and Rhinorrhoea

Sufferers often refer to this as a "constant cold" or hay-fever, because of the similarity of the symptoms.

In non-specific rhinorrhoea the characteristic attacks usually occur in the morning, at or soon after waking; or they may be precipitated by some change in body temperature, as in getting out of bed or walking barefoot on a cold floor, by exposure to bright sunshine, by the inhalation of airborne irritants, especially dust, or by emotional stress. There is a feeling of irritation at the root of the nose, followed by sneezing, lacrimation, and a profuse watery discharge. The rhinorrhoea may last for only a few minutes or may continue for several hours, and as many as 50 large handkerchiefs may be soaked in the course of a day. Nasal obstruction, headaches, and anosmia are not usually features of the condition.

The attacks are distinguished from common colds by the nature of the discharge, which is clear and watery and not purulent as in coryza. Secondary infection can follow these attacks, whereupon the discharge becomes thick and yellow, and the infection is then identical with the later stages of a coryza.

Examination of the nose shows that the mucous membrane has a normal appearance, and during an attack the nasal passages are found flooded with thin mucus. Radiological examination of the sinuses indicates no change beyond a slight and occasional mucosal thickening. Repeated