CLINICAL PATHOLOGY IN GENERAL PRACTICE

TESTS FOR VENEREAL DISEASE—I

BY

I. N. ORPWOOD PRICE, M.R.C.S., D.P.H.

Pathologist, Whitechapel Clinic, the London Hospital

Laboratory tests for venereal diseases are mostly concerned with gonorrhoea, non-gonococcal urethritis, chancroid (soft sore), and syphilis. Occasionally laboratory investigation of material from patients suspected of suffering from lymphogranuloma inguinale (lymphogranuloma venereum) or granuloma inguinale is required.

At the outset it must be stressed that before instituting treatment for any of these diseases laboratory assistance should always be invoked in order to confirm the diagnosis, and when occasion demands the practitioner should telephone the laboratory for the necessary glass slides, culture media, or screw-capped bottles, with containers for transport. If he is uncertain about which test should be employed, the pathologist will give the necessary advice.

Before taking material from the patient it is essential to mark each specimen with an identification name or number, the date, and the source of the material—for example, urethra, vagina, cervix.

Specimens

No laboratory, whatever resources it may have or however experienced its personnel, can give the best advice when supplied with badly harvested specimens.

Smears.—These should be made with a platinum loop, thinly and evenly spread on the glass slides (mix a drop of sterile water or saline with a drop of the discharge, if necessary), and fixed by moderate heat—that is, the slide should not feel uncomfortably hot when placed on the back of the hand.

Cultures.—Specimens for culture should be taken with a platinum loop or the sterile wool swabs supplied by the laboratory.

Blood.—With adults at least 8 ml. should be obtained by venepuncture. In order to avoid haemolysis, which may vitiate the tests, care should be taken that both needle and syringe are free from water or spirit (wash out with sterile normal saline if necessary). The needle should be removed before transferring the blood to the sterile screw-capped blood bottle, care being taken to use no undue force. In the case of adults or small children with “difficult veins” a guarded safety-razor blade can be used. All except the last $\frac{1}{4}$ in. (3 mm.) of one cutting edge is covered with adhesive plaster; a clean-cut nick is inflicted into the surgically cleaned and warmed heel of the patient. If the skin is stretched by the use of the thumb and forefinger of the left hand and the cut made at right angles to the “stretch,” the small wound will tend to gape and sufficient blood will usually run into the container (5-ml. bijou screw-capped bottle) held against the skin below the cut. Umbilical cord blood, as will be explained in Part II of this article, is not a suitable specimen to collect.

Cerebrospinal Fluid.—It is, of course, essential that cerebrospinal fluid should not contain any blood.

Serum from Primary Sore.—A capillary tube closed at one end should be used to obtain serum from a suspected syphilitic sore if the presence of *Treponema pallidum* is to be demonstrated. If the sore is covered with a scab, this should be removed, the lesion washed thoroughly with plain tap-water or normal salt solution (soap should not be used), and then dried. The dried surface is gently scari-

fied so as to provoke *slight* bleeding and the exudation of serum. Wipe away the first few drops of exudation, especially if they contain blood, and wait for a clear drop of serum to appear from the depth of the lesion. Gently heat the closed end of the capillary tube and immediately apply the open end to the drop of clear serum, and as the closed heated end cools allow sufficient serum to be drawn up into the tube so that none is present $\frac{1}{2}$ in. (13 mm.) from either end of the tube. The open end is then sealed by heat.

Clinical Details.—Finally, all specimens should be accompanied by a brief note containing the name and address of the doctor, the patient's name or identification mark, the essentials of the clinical history, the probable diagnosis, and the examination required. It cannot be too strongly stated that “slot-machine” laboratory reports are largely a waste of time and money. If the pathologist and clinician do not collaborate, the latter may well be deprived of vital information.

Transport.—The speedy transport of specimens from the patient to the laboratory is of paramount importance. Time is often the essence of a useful investigation. Fluid specimens may deteriorate rapidly after being taken, and thus the examination of such material would be fruitless. For example, the cells in the cerebrospinal fluid begin to disappear within two hours of lumbar puncture, and in 24 hours few, if any, cells can be seen.

Reports.—Reports on the findings of tests for venereal diseases are intended to be regarded only as part of the evidence required in order to make a diagnosis. Thus a report that no gonococci can be found in the smear from a urethral discharge is no proof that the patient is not suffering from gonorrhoea. Further investigation may be required and the final decision must rest with the clinician.

Identification of Gonococci

When the patient is suspected of suffering from gonorrhoea, three types of test should be employed—namely, smears, cultures, and blood tests.

In acute gonorrhoea in the male smears should be made with a platinum loop from urethral discharge after carefully cleaning up the glans penis and meatus with sterile water or saline. In the female, with the patient in the lithotomy position, smears should be made from the urethra, posterior fornix of the vagina, and the cervical canal. In the case of suspected chronic gonorrhoea in the male smears should be made from the prostatic secretion after massage as well as from the early morning discharge obtained from the urethra after the patient has held his water all night. In the female, examination of smears from the urethra, cervical canal, and the rectum should be made. Satisfactory rectal smears cannot be obtained without the aid of a proctoscope, and unless the practitioner is skilled in the use of this instrument he should obtain the advice of an expert. Strands of mucus on the mucosal wall can often be found (30% cases, C. S. Nicol, personal communication), which on examination may reveal the presence of gonococci.

Most bacteriologists will report that the smears contain Gram-negative intracellular diplococci and/or other organisms such as coliform bacilli, diphtheroid bacilli, or monilia. Further information on the cellular content of the smear will also be given. Polymorphonuclear leucocytes predominate in acute inflammatory conditions, whilst epithelial cells suggest a more chronic infection. Two points must be remembered: (1) that the absence of the relevant organism

is no proof that the patient is not suffering from gonorrhoea, and an examination of another specimen may be indicated ; and (2) a report that Gram-negative intracellular organisms morphologically resembling gonococci have been seen will not be accepted in a Court of Law as proof that the patient from whom the specimen came is suffering from gonorrhoea, because other organisms of the *Neisseria* group may occasionally be present in the genital tract.

In suspected gonorrhoea in the female smears and cultures should be taken just after the menstrual period.

Cultures should be taken immediately after the smears and from the same anatomical sites. Direct inoculation of discharges on to solid media has been largely abandoned in favour of transport media. The sterile cotton-wool swab impregnated with carbon provided by the laboratory should be well rubbed into the discharge and then plunged straight into a bijou bottle containing Stuart's transport medium. The wooden stick is snapped off by pressure against the side of the bottle so that the screw cap can be replaced and tightly screwed up again. Since the success of this transport medium depends on anaerobic conditions, this manoeuvre should be carried out with speed and great care taken to check that the screw cap is not loose when the operation is completed. After receipt at the laboratory, the swab is used to inoculate suitable media, incubated in an atmosphere of carbon dioxide and examined after 24 to 48 hours, depending on the amount of growth obtained. Typical pure gonococcal colonies, if present, are picked off, some examined under the microscope, some tested with the oxidase reactions, and some subcultured for testing with sugar reactions. If gonococci are grown the organism can be isolated in pure culture and "sugared" in 80% of such specimens. These are reported as morphologically and culturally gonococci. The 20% which cannot be obtained in pure culture are reported as "presumptive" gonococci, and warrant the institution of anti-gonococcal treatment, provided the history and clinical condition indicate a strong probability that the patient is suffering from gonorrhoea. If there is any doubt consultation with the bacteriologist is indicated.

Gonococcal Complement-fixation Test

The only blood test at present in routine use is the complement-fixation test for gonorrhoea (G.C.F.T.) and reports on this test are given as negative or positive. Positive reactions are quantitated and reported as positive with neat serum or a dilution thereof. A positive reaction is evidence of a gonococcal focus of infection, but in the absence of any supporting evidence the test should be repeated. A negative reaction, however, does not exclude gonorrhoea, because in the early stages of the disease the body may not have had time to elaborate the antibodies detected by the test. The table shows how soon after exposure to infection one might expect a positive G.C.F.T. in untreated gonorrhoea.

Time after Exposure	No. of Cases	Percentage with Positive G.C.F.T.
1st week	663	9
2nd "	491	16
3rd "	160	34
4th-6th weeks	90	39
7th week	26	50

During the course of treatment of acute gonorrhoea with penicillin the fate of the reaction is as follows : 68% fail to record a positive reaction throughout the course of their illness ; 19.2% are positive at first testing before treatment is begun ; 6.4% at the end of the first week of treatment ; 2.5% at the end of the second week ; 0.8% at the end of the third week ; and 0.5% at the end of the fourth to eighth weeks. The remaining 2.1% show "oscillating" reactions—that is, on serial testing at intervals the reaction results recorded may be negative, positive, negative, or alternatively positive, negative, positive. This type of serological pattern probably means that the antibody content of the

patient's serum is sometimes over and sometimes under the threshold of recognition by the test. In gonorrhoea this process goes on for a relatively short time, until antibody production recedes, and thereafter negative results are recorded.

Of patients who record a positive reaction at the first testing before treatment, approximately 64% may be expected to become negative after treatment within three months, leaving 36% who will become "oscillators" and revert to negative in due course. Of patients who have a negative test when first tested, and subsequently become positive, 60% will have a positive G.C.F.T. before the end of the first week, when in all probability the treatment has not had time to take effect ; 24% will become positive before the end of the second week ; and the remaining 16% will become positive during the third to eighth weeks. Complications of gonorrhoea are not common, but at least 66% can be expected to show a positive G.C.F.T.

The value of the G.C.F.T. may be summed up as follows : in early acute gonorrhoea, its use is limited ; it is of value in subacute or chronic gonorrhoea, particularly in the female, when smears may give little or no help initially ; it is of definite value in the diagnosis of complications ; it is useful in the management of gonorrhoea ; it is unwise to ignore a positive test which has been confirmed by retesting.

Non-gonococcal Urethritis

Non-gonococcal urethritis is a most troublesome condition. According to the latest Ministry of Health returns 11,552 men attended public V.D. clinics with this condition in 1952, as compared with 15,510 new cases of gonorrhoea in the male. The cause of the condition is unknown, though it is thought that it may be due to a virus infection or one of the organisms of the *Pleuropneumonia* group. Laboratory help is therefore entirely confined to negative findings in excluding a diagnosis of gonorrhoea. In the acute stage of the disease smears from the urethral discharge usually reveal the presence of masses of leucocytes but no organisms. Later many lymphocytic cells and contaminants such as staphylococci and diphtheroid bacilli may also be found. In like manner cultures will either fail to grow any organisms or grow only the contaminants. Blood tests for gonorrhoea are always negative.

Chancroid

Laboratory tests in the diagnosis of chancroid are of some value. Smears taken from the sore usually contain a mass of leucocytes plus a variety of contaminating organisms. Occasionally organisms resembling Ducrey's bacilli (small Gram-negative bacilli, characteristically arranged in chains) may be seen. Cultures can be taken into clotted sterile rabbit's blood which has been heated at 55° C. for five minutes. Fresh medium should be used and a bead of discharge is obtained from the base of the ulcer under its undermined edge by means of a platinum loop. The medium is inoculated by the wire being passed several times round the clot so as to ensure even distribution of the inoculum. Two media tubes should be inoculated and examined after 24 to 48 hours. According to Teague and Deibert* if the characteristic chains and tangles of small Gram-negative bacilli are found even in admixture with contaminants the culture may be regarded with confidence as being positive for Ducrey's bacillus. In all cases a cutaneous skin test (dmelcos) should be performed to confirm the diagnosis and standard serum tests should be requested with a view to excluding syphilis. Always remember that the first result of the latter may be negative if the blood is taken during the incubation period of syphilis.

[Part II will appear in our next issue.]

*System of Bacteriology, 2, 415. Medical Research Council, 1929.