CLINICAL PATHOLOGY IN GENERAL PRACTICE

SEROLOGICAL TESTS IN DIAGNOSIS

BY

E. T. C. SPOONER, M.D.

Professor of Bacteriology and Immunology, London School of Hygiene and Tropical Medicine

In this country serological tests—that is to say, the detection and measurement of antibodies in the patient's serum—aid diagnosis in enteric fever (typhoid and paratyphoid B), brucellosis (undulant fever), leptospirosis (Weil's disease and canicola fever), infective mononucleosis, pneumonia, influenza, mumps, psittacosis, various other virus infections, Q fever, and toxoplasmosis; also in typhus fever and smallpox, cases of which may arrive from overseas by ship or aeroplane. Each of these conditions is considered separately below. The serological diagnosis of venereal disease has already been dealt with in this series.*

Serological tests can seldom, if ever, establish a diagnosis without the aid of clinical information and usually of other laboratory investigations. The occurrence of "normal" antibodies and the variability of antibody production in humans influence antibody titres unpredictably and necessitate caution and experience in interpreting their meaning. For this reason the final interpretation of an antibody titre must be made by the clinician who knows the full case history. The bacteriologist can indicate probable interpretations of his tests, but he can do this to the best advantage only if he knows something of the circumstances of the case. A bald request for "Widal test, please" deprives the bacteriologist of much of his power to help: he should always be put in the picture as much as is reasonably possible. Apart from name, age, and tentative diagnosis, he should know how long the patient has been ill, what the main signs and symptoms are, the patient's occupation, whether he has been recently overseas, details of relevant vaccination history, and so on. Very often a telephone conversation between clinician and bacteriologist is well worth while: it may lead the bacteriologist to make tests which the clinician had not thought of, and is in any case likely to increase the value of the laboratory report.

The evidence given by a serological test is less conclusive than that of a positive blood culture or virus isolation, especially in the first few weeks of "pyrexia of unknown origin." The bacteriologist can always give advice on the choice of appropriate tests as well as on their interpretation. He will usually also supply outfits for the collection of specimens. In difficult cases, therefore, he should be consulted early.

Collection and Dispatch of Specimens

Most diagnostic laboratories will supply on request a "blood collecting outfit," comprising a dry sterile syringe with needle and a dry sterile 15- or 25-ml. screw-capped bottle or "universal container," in which the blood sample can be sent to the laboratory. Most laboratories also supply bottles of culture medium for blood cultures, should it be necessary to take these at the same time.

It is wiser to use one of these laboratory-prepared collecting outfits rather than to boil up one's own syringe, partly on account of convenience and safety, but also be-

cause a blood sample can be spoiled by a little water or spirit in the syringe. The barrel and plunger of the syringe must fit really well, or else leakage of air along the plunger may prevent the withdrawal of sufficient blood.

Before venepuncture the skin over the antecubital area should be cleaned with spirit or other disinfectant, and must be allowed to dry completely before the needle is inserted. At least 5 ml. of blood should be taken for a serological test. If a blood culture or other blood test is to be made at the same time, a sample of 10 ml. allows 5 ml. for this and 5 ml. for the serum test.

As soon as the sample of blood for the serum test has been placed in its sterile screw-capped bottle, the syringe and needle should be washed out with 2% lysol solution before residual blood has had time to clot in them. They should then be well rinsed under the cold-water tap, and returned, with the sample, to the laboratory. Unless the laboratory is in the same building, the bottle of blood should be allowed to stand at room temperature for a few minutes, until the blood has clotted, and should then be packed and sent at once, by hand if possible, to the laboratory.

If the specimen is sent by bus or by passenger train the laboratory should be notified by telephone, so that collection from the station may not be delayed. It is best not to send blood samples by post: in this country it is hardly ever necessary to do so. Postal travel may render blood quite unsuitable for some tests. If the examination required is a special one, such as a virus complement-fixation test, which can be made only at some central laboratory many miles away, the best procedure is to send the blood by hand, bus, or passenger train to the local laboratory, where the serum can be separated and forwarded to the central laboratory. Serum travels very much better than does whole blood, and may be sent by post.

Tests for Antibody in Serum

Four kinds of test are used to detect and measure antibodies for routine diagnostic purposes. These are agglutination tests, complement-fixation tests, haemagglutinin inhibition tests, and precipitation tests. Various other tests are necessary in the exploration of new or obscure diseases.

The only commonly used precipitation tests are the Kahn test and its various derivatives, which are employed in the diagnosis of syphilis. In agglutination tests, dilutions of the serum are mixed with suspensions of bacteria (as in the Widal and Weil-Felix tests) or of red blood corpuscles (as in the Paul-Bunnell test) and agglutination or clumping is recorded. Complement-fixation tests follow the pattern of the Wassermann test: antibody in the patient's serum when mixed with its specific antigen can deprive fresh normal guinea-pig serum of its ability to haemolyse sensitized sheep red corpuscles. Complement fixation is occasionally of value in the diagnosis of chronic gonorrhoea and non-gonococcal urethritis, but is chiefly used in the investigation of virus infections. Haemagglutinin inhibition tests are used as alternatives to complement-fixation tests to measure antibodies to influenza and mumps viruses.

The result of a serological test is expressed as a titre or the highest dilution of serum giving a positive reaction.

Because many antibodies are present at low titre in normal sera, owing either to past exposure to infection or to prophylactic vaccination, as with T.A.B. vaccine, it is

*March 6, p. 575, and March 13, p. 637.

usually difficult to assess the significance of any titre unless it is a very high one; and high titres are unusual in the early stages of the infection. For these reasons, *demonstration of a rise in titre when a serum sample is compared with another taken several days earlier has a much higher diagnostic significance than has any single serum titre.* It is customary to accept a fourfold rise in titre as significant, because smaller rises may sometimes be due to non-specific causes. The stage of the disease and the time interval between paired serum samples will obviously affect interpretation. The bacteriologist should be consulted about the interval between samples, because the optimum interval is different in different circumstances.

Pyrexia of Unknown Origin

This is not the place for a discussion of the diagnosis of obscure pyrexia. The most important laboratory tests are those for the isolation and recognition of a pathogenic micro-organism or virus; but at the same time a sufficient sample of blood should always be taken for all relevant serological tests, because a rising antibody titre of one kind or another gives the best and sometimes the only clue to correct diagnosis. In addition to bacterial infections, the more probable non-bacterial infections, such as infective mononucleosis, psittacosis, Q fever, and leptospirosis, must be remembered.

Enteric Fever

In Britain typhoid and paratyphoid B are the usual enteric fevers. Paratyphoid A and C are rare, but should be remembered when the patient has recently returned from overseas. In enteric fever blood culture may be positive during the first three or four weeks, especially during the first and second. A positive blood culture gives a much surer guide to diagnosis than does any serological test, and therefore blood culture should never be neglected. Serum should, however, be tested for antibodies, because blood cultures are not always successful. Antibodies specific to the enteric salmonellae usually begin to appear around the end of the first week. They are measured in terms of their ability to agglutinate standard suspensions of enteric bacilli, in a test called the Widal or Widal-Grünbaum test. The suspensions against which the serum is tested in this country are usually *Salm. typhi* and *Salm. paratyphi B* O, *Salm. typhi* and *Salm. paratyphi B* H (specific), and a non-specific group H suspension which will detect antibodies to various other salmonellae without disclosing which they are.

Antibody to the *Salm. typhi* O, or somatic, antigen may react also with the somatic antigens of *Salm. paratyphi B* and vice versa, but the H, or flagellar, antigens and their antibodies are more specific. Antibodies to O and H antigens do not always develop *pari passu*, and therefore both must be measured.

Serum titres depend on past exposure to infection, on the duration of illness, on vaccination history, and on individual ability to produce antibody. It is therefore impossible to name titres which definitely indicate infection. It is usually safe, in *uninoculated patients*, to regard an H titre of 1/50 or an O titre of 1/100 observed during the first ten days of the disease as strong presumptive evidence of enteric infection, but lower titres by no means exclude the diagnosis of enteric, nor are higher titres certain evidence of infection, especially in those who have been vaccinated. Later in the disease titres of 1/1,000 or more are not uncommon, but they may be much lower.

The strongest serological evidence of enteric infection is a rising O titre. A rising H titre, though strongly suggestive, is less convincing, especially in the previously vaccinated, in whom pyrexial diseases may produce some non-specific rise in salmonella H titre.

In subjects who have previously received T.A.B. vaccine, the Widal test can provide little useful evidence, except when a pronounced rise in O titre is revealed by examination of

paired sera. Antibodies to the Vi antigen of *Salm. typhi* are not as a rule helpful in the diagnosis of acute cases, but may be valuable in detecting chronic carriers of the typhoid bacillus.

Food-poisoning

During the acute phase serological diagnosis is useless, and all efforts should be made to cultivate the responsible organism. A retrospective diagnosis may sometimes be made in the weeks following the attack by detecting a rise in antibodies which agglutinate the commoner food-poisoning salmonellae, such as *Salm. typhi-murium*, *Salm. dublin*, etc.

Brucella Infection : Undulant Fever

Practically all brucella infections in man in Britain are due to *Br. abortus*.

As with enteric fever, blood culture is the more valuable test when it is positive, but a search should always be made for agglutinins to *Br. abortus*. The test is essentially the same as for enteric infections, and its interpretation presents parallel difficulties. Normal brucella titres up to about 1/80, or even higher, may occur in healthy persons, such as farmers or dairy workers, whose work exposes them to the risk of infection, or who are drinkers of raw milk. Titres of less than 1/80 may be very suggestive when they occur in pyrexial patients whose disease resembles undulant fever clinically. Most clinical cases show titres of 1/500 to 1/2,500 at some stage of the disease. The higher the titre, the more suggestive it is. A negative test does not, however, exclude brucella infection. Great importance should be attached to a rising titre.

Typhus Fever : The Weil-Felix Test

No typhus fever is endemic in Great Britain now, though cases from abroad may be encountered. The sera of patients with either epidemic or murine typhus, which are the only kinds to appear here, will usually at some time after the fifth day of the disease agglutinate a suspension of *Proteus* OX 19 (the Weil-Felix reaction). The test depends on the fortuitous sharing of an antigen by the rickettsiae which cause typhus and the quite unrelated *Proteus* OX 19.

The Weil-Felix test is the most generally useful laboratory test for typhus fever. The interpretation of low titres is again difficult; two or more separate samples of serum should always be tested to detect a rising titre, and the significance of any titre measured must be assessed in conjunction with the clinical features of the case. Titres of 1/50 are suggestive; titres of 1/100 are more so. But the significance of a rising titre is greater. Titres of 1/1,000 or more are strong evidence, but seldom occur early in the disease.

Infective Mononucleosis (Glandular Fever) : The Paul-Bunnell Test

The aetiology of glandular fever is unknown: there may be more than one cause. The sera of some, but not all, patients with this disease agglutinate sheep red blood corpuscles. Heterophil antibodies which also agglutinate sheep red blood cells may appear in human sera for other reasons, one of which is past injection of horse serum; little reliance therefore can be placed on a simple sheep cell agglutination test. However, Barrett has shown that the antibodies produced in infective mononucleosis differ from other heterophil antibody in being absorbable with ox red cells but not, or by no means completely, with a suspension of guinea-pig kidney. A titre of 1/128 or more given by the patient's serum after absorption with guinea-pig kidney suspension, as well as by the unabsorbed serum, is very suggestive of infective mononucleosis.

An explanation of this quite empirical test will probably have to await the discovery of the agent which causes the disease. To avoid bleeding the patient repeatedly, blood for a complete blood count should be taken at the same time as the sample for serum.

Virus Infections

Virus diseases, in the diagnosis of which serological tests are useful, present clinically as pyrexia with few localizing signs, or as pneumonia, or with meningitic syndrome.

Psittacosis and *Q fever* are best revealed by isolation of the virus or rickettsia from blood or sputum. Complement-fixing antibodies to a titre of 1/30 or more are suggestive, but a rising titre is more so, and paired sera should always be examined.

Lymphogranuloma venereum is likely to be diagnosed on clinical grounds: the serum may give complement fixation with L.G.V. or psittacosis antigen.

Pneumonia

The terms "virus pneumonia" and "primary atypical pneumonia" are applied somewhat loosely to pneumonias which are not caused by bacteria and which are not evidently secondary to some other infection. Pneumonia, often mild, may be caused by a psittacosis virus or by the *Coxiella burnetii* of Q fever. In both of these infections the responsible virus may sometimes be isolated early in the disease, or the diagnosis may be revealed by rising antibody titres.

Influenza viruses A and B may occasionally produce a true pneumonia, although more usually they cause a bronchitis or bronchiolitis, on which a secondary bacterial infection may be superimposed. Influenza antibodies may be revealed by complement-fixation or haemagglutinin inhibition tests on serum. Serological diagnosis rests chiefly on demonstration of a rising titre in paired sera.

Of the cases of pneumonia not included in one of the categories so far mentioned, about one-half yield no evidence of their causes; the other half constitute a group, probably due to a virus which is not yet clearly defined, recognizable by the appearance in the patient's serum of "cold agglutinins" and/or agglutinins to the MG streptococcus, an apparently non-pathogenic throat streptococcus the aetiological relationship of which to the disease is doubtful. Cold agglutinins are recognized by their ability to agglutinate human group O red corpuscles at cold room temperature but not at 37° C. or higher.

Serological diagnosis is not very helpful in the management of virus pneumonia, since the various antibodies that may be measured do not appear until the second or third week of the disease, by which time antibiotic treatment is likely to have been tried if circumstances so dictate. An approximate diagnosis is usually made on clinical and bacteriological grounds; the mortality is not high.

Virus Meningitis and Meningo-Encephalitis

So-called "benign lymphocytic meningitis," or meningitis in which examination of the cerebrospinal fluid reveals no bacterial cause and a lymphocytic response, may be due to the viruses of poliomyelitis, lymphocytic choriomeningitis, mumps, or herpes. A similar syndrome may occur in infective mononucleosis, infective hepatitis, and, rarely, lymphogranuloma venereum. Leptospirosis and toxoplasmosis may have to be considered. In the second or third week of disease, or retrospectively, antibodies may appear which were absent from, or present only in low titre in, acute phase sera. Antibodies for lymphocytic choriomeningitis, mumps, and lymphogranuloma venereum can be detected by complement-fixation tests; tests for infective mononucleosis have already been discussed.

Antibodies for herpes are too widespread to be of great diagnostic value. Those for poliomyelitis can be detected by slow and costly neutralization tests in the monkey, or by complement-fixation tests with antigens prepared from suckling mouse brain or tissue-culture preparations of the virus. Neutralization tests on tissue-culture virus may also be useful. At present, laboratory facilities for serological diagnosis of poliomyelitis are limited and chiefly experimental; moreover, the information that they can give is of more use in epidemiological research than in the manage-

ment of cases. The Central Public Health Laboratory, Colindale Avenue, London, N.W.9, should be consulted before serological evidence of poliomyelitis is sought.

The most useful serological tests in virus meningitis or meningo-encephalitis are those for lymphocytic choriomeningitis and mumps. Isolation of virus from the cerebrospinal fluid may be attempted if the necessary facilities are available.

Mumps antibodies can be measured either by complement-fixation or by haemagglutinin inhibition tests. Their detection may facilitate diagnosis of other atypical forms of mumps, in which parotitis is not a dominant sign. Generally speaking, in obscure cases of meningitis and meningo-encephalitis, cultural, cytological, and biochemical examination of cerebrospinal fluid gives more useful information than do serological tests.

There are as yet no laboratory tests for the virus of infective hepatitis and its antibody.

Smallpox

Smallpox diagnosis rests chiefly on the examination of the vesicle or pustule contents and scabs, by complement fixation and by isolation of virus in eggs. Serological tests are relatively unimportant, but complement-fixation and virus-neutralization tests on patients' sera may be useful in establishing a retrospective diagnosis.

Leptospirosis: Weil's Disease and Canicola Fever

During the first week blood should be examined for the presence in it of leptospirae. From the end of the first week onward antibodies usually appear. These are detected by agglutination tests on serum collected in the usual way already described. Agglutination can distinguish between the antibody to *L. icterohaemorrhagiae* and to *L. canicola*, the only two leptospirae known to infect man in Britain. It is important to seek for a rising titre. Low titres may have considerable significance, depending on the technique of the test used; the bacteriologist must be consulted.

Toxoplasmosis

The two most useful serological tests for the detection of toxoplasma antibodies are the complement-fixation test and the cytoplasm-modifying test, or "dye test," which depends on the power of the antibody to modify the staining reaction of the parasite. Antibodies demonstrable by the dye test appear earlier and persist longer than do those which fix complement. Since infants may be infected from their mothers, the serum of the mother should always be examined when the disease is suspected in her infant. A rising titre is significant early in the infection.

Next article on Clinical Pathology.—"The Assessment of Gonadal Function," by Dr. G. I. M. Swyer.

The risk of premature detonation of blasting charges by radio and television waves is among the subjects which a Swedish safety engineer is going to study on a four-nation tour arranged by the International Labour Organization. He is Yngve Hagerman, Chief of the Mining Department of the Swedish Workers' Protection Board. He will visit France, Germany, Canada, and the United States. Among the problems facing his department is the protection of workers against premature detonations in blasting operations as a result of lightning and thunderstorms. "Television has not yet come to Sweden," Mr. Hagerman said at an interview, "but it is coming. We have had no detonations from radio waves, but we have read about United States precautions against this risk and I would like to study these." He also wants to find out whether radio waves sent out by police cars and road maintenance and snow clearance squads could detonate a charge just as they could bring down avalanches in the mountains.