CLINICAL PATHOLOGY IN GENERAL PRACTICE

BIOPSIES

BY

RAYMOND WHITEHEAD, M.D., D.Sc.

Reader in Pathology, University of Manchester

A biopsy is an examination of tissue removed surgically. Since the term "biopsy" implies an examination, it is superfluous to speak of "biopsy examination." When the surgeon says he is "taking a biopsy," he is in fact "taking a specimen for biopsy" or "taking a biopsy specimen"; biopsies are performed by the pathologist, not by the surgeon, who simply obtains the material. Another name for this branch of pathological work is "surgical histology," which indicates clearly both the source of the material and the characteristic feature of the examination. The term "histopathology" has no advantage over "biopsy" and "surgical histology," and does not indicate whether the material was obtained from the living or the dead.

Biopsy may be performed before, during, or after treatment of the patient. When the lesion is both large and readily accessible—for example, on the surface of the body or in the mouth or rectum—the surgeon may remove a small piece of it and await the pathologist's report before deciding on the treatment. A lesion that is both small and readily accessible should be excised completely; there is no point in subjecting the patient to two operations; one, an excision of a minute fragment of the lesion for pathological diagnosis, and the other, excision of the rest of the lesion. Surgical excision of a lesion is preferable to therapeutic destruction of it *in situ*, since excision allows of a microscopical examination. When the diagnosis is reasonably certain on clinical grounds—for example, if the patient has the usual signs of a mammary scirrhus—the surgeon treats the patient radically without preliminary biopsy in the confident expectation that the diagnosis will be subsequently confirmed by the pathologist. When the lesion is internal, preliminary biopsy is often impracticable, and the biopsy can be made only on the specimen obtained after treatment on the strength of the clinical findings.

Value of Biopsy

All material removed surgically is worth examining, no matter how sure of his diagnosis the surgeon may feel. Although the clinical diagnosis is often confirmed, surprises are not rare, and the prognosis or treatment may be radically revised as the result of the pathologist's report. Malignancy may be simulated exactly by benign conditions; familiar examples are certain inflammations of the large intestine; and giant hypertrophic gastritis, though rare, may be clinically indistinguishable from gastric cancer. Conversely, lesions that the surgeon thinks benign—in the breast, for instance—may prove on microscopical examination to be malignant.

The certainty of pathological diagnosis varies as widely as that of clinical diagnosis. Many diseases are identifiable with certainty on examination of a section; these include most surgically removed cancers and tuberculosis and other well-known infections. Sometimes only the general nature of the disease is determinable; for example, a malignant growth may be so undifferentiated as to be unclassifiable even broadly as carcinoma or sarcoma, but the fact of malignancy is nevertheless an important item of informa-

tion. Sometimes the same histological picture is given by different diseases; in some cases of Hodgkin's disease the lesion is microscopically identical with that of mixed-cell sarcoma, and the two conditions have probably been confused. Similarly, early Hodgkin's disease in a lymph node may be microscopically indistinguishable from lymphatic leukaemia or lymphosarcoma. Some clinically malignant tumours may show little or no histological evidence of malignancy; these include certain "myxomas," "lipomas," and tumours of the thyroid; in such cases treatment must obviously be based on the clinical findings, supplemented when practicable by examinations other than biopsy.

The Specimen

The sooner a specimen is fixed, the better the section will be; the specimen should therefore be put in a fixing fluid immediately after its removal from the body; if the fresh specimen is put aside in a dish to be dealt with at the end of the day's operating, it may easily be thrown away or be left to dry until the hatching of maggots reminds someone of its presence.

The best practice is for the laboratory to keep the operating theatres supplied with vessels of various sizes already containing the fixative. The most convenient fixative is 10% formol saline; a litre of this consists of 100 ml. of commercial formalin, 900 ml. of tap-water, and 8.5 g. of sodium chloride. Since commercial formalin is a 40% solution of formaldehyde, the fixative contains 4% formaldehyde. Owing to the similarity of their names, formol saline and normal saline could be confused, with serious results; it would therefore be best if each container bore a large coloured label with the words "Pathological Fixative—Poison."

The whole of the tissue removed should be sent to the laboratory. This enables the pathologist to see the lesion as a whole, to note its relation to surrounding tissues, and to choose the best points for sampling. The whole of an amputated limb is not often sent, but if the condition is serious enough to require amputation the pathologist ought to have the whole limb so that he can study the extent of the tumour or the state of the blood vessels. To deal satisfactorily with large specimens such as limbs or massive tumours, the pathologist needs the help of the surgeon; by examining such specimens together, they can decide on the most suitable parts for microscopical examination, and so obviate the necessity for cutting many blocks. It is not unknown for the laboratory to receive only a minute fragment of a large excised tumour, or even only tissue adjoining an excised tumour, because the tumour itself was wanted for somebody's private collection. The commonest mistake of the surgeon is to send a part when he should send the whole.

When the specimen is more than a few inches in diameter, the surgeon should bisect it through the part containing the lesion. This gives the fixative immediate access to the lesion and incidentally allows the surgeon to see at once what he has removed. The worst fault is to make multiple incisions into such specimens as lymph nodes and small tumours of the breast; in such cases it may be impossible to get a satisfactory section for microscopical examination. Another good reason for not incising lymph nodes is that they may be heavily infected; they should be allowed to fix for some time before being opened.

Urgent Specimens and Dispatch

When a request form is marked "urgent" the time normally needed for reporting is reduced by (1) processing faster than usual if the specimen is small, (2) examining the section as soon as it is ready, and (3) telephoning the report. Abnormally rapid processing may result in poor sections, and consequently the diagnosis may be difficult or impossible; extra tissue may then have to be processed, the final result being a longer delay in reporting than if the request had not been marked "urgent." It is not desirable to process lymph nodes faster than normal, although time can be saved by immediate examination and telephoning the report.

There may be a variety of good reasons for "urgent" requests, but such requests should be made with discretion, and preferably with an explanation of the nature of the urgency. The laboratory has equal obligations to many surgeons, and every "urgent" request from one surgeon means delay in reporting for others. Most surgeons in fact make "urgent" requests only at long intervals, and the requests are accordingly accepted without question; when "urgent" requests come in large numbers from a single firm, the pathologist may well suspect that the word "urgent" is being used with insufficient discrimination.

A specimen for examination deserves at least the same care as a registered letter. There is little or no risk of loss once the specimen has been collected from the operating theatre by the messenger; but loss and delay are familiar enough in the theatre. The trouble is that usually no one is responsible for seeing that specimens are dispatched promptly and properly. The surgeon may ask the theatre sister to send them, but she may well pass on the instruction to another nurse or orderly—and either may have gone off duty when the messenger arrives to collect the day's specimens.

One person should be made responsible for the dispatch of specimens, and the most suitable person is the surgeon's assistant, who knows the importance of pathological examinations and what the requirements are; no member of the nursing staff is properly qualified for this function. The assistant in charge of specimens should never delegate his responsibility, but see to it personally that the specimens are collected by the messenger. There should be no difficulty in fixing regular times for the collection of specimens.

Data Needed by Pathologist

The meaning of a histological picture sometimes depends on the patient's age and the history of the lesion; if the pathologist is not informed of these, he can easily make a wrong diagnosis. A familiar example is the pigmented mole of the skin; this may look particularly malignant under the microscope and a report in this sense might result in an amputation. In childhood such moles are usually benign, however malignant they may look, and inquiry may show that the lesion has been present for years without any increase in size.

The accumulated experience of pathologists finds expression in the blank form commonly supplied to the surgeon who wants a specimen examined. This form requests such obvious particulars as the patient's name, sex, age, and ward; the site of the lesion; the duration of the illness; the clinical diagnosis; and the treatment given, especially if it is likely to affect the microscopical appearances, as in toxic goitre. This represents the bare minimum of information needed by the pathologist before he can form a sound opinion. All the information asked for by the laboratory should be supplied, preferably by the surgeon himself and not by an assistant or nurse.

The printed form cannot provide for all the information sometimes required by the pathologist. In specially difficult or interesting cases the surgeon may realize this and anticipate the need by sending unusually detailed information, but the pathologist often has to extract the information himself from the surgeon or his staff. When full information has

been supplied, it is sometimes possible for the pathologist to suggest a likely diagnosis from this alone.

Whenever there is doubt or difficulty on either side, surgeon and pathologist should meet for discussion. Such consultations usually result in complete agreement on the most probable diagnosis and the most suitable treatment and are of immense benefit to the patient; some surgeons are well aware of this and consult regularly. At the other extreme we have the surgeon who goes in for "penny-in-the-slot" pathology; consulting, usually with the reporter's superior, only when a mistake has been made through the surgeon's failure to supply essential information; or seeking a second opinion on a section without the knowledge of the reporter. A competent pathologist knows when he ought to consult a fellow-pathologist, and, the more able he is, the more often he sees the need and consults. Any query about a report should of course be raised with the reporter, not with a third party.

Paraffin and Frozen Sections

Paraffin sections are normally used for diagnostic purposes; the sections are stained with haemalum and eosin or some other routine combination of dyes. The introduction of automatic tissue processors has greatly reduced the minimum time interval between receipt of the specimen and issue of the report. The normal interval is now from two to three days, but very small specimens, such as those from a bronchus, can be processed and reported on within a day of receipt. These periods are attainable only if the section is of satisfactory quality and the microscopical diagnosis obvious. The chief cause of delay in reporting is difficulty in diagnosis; this may call for staining of further sections by special methods, the making of extra blocks, consultation with the surgeon or another pathologist, or work in the library. In such cases it may be possible to issue a provisional report.

The advantage of frozen sections is that they can be made within a few minutes of receiving the material, and where special arrangements exist the surgeon can interrupt the operation until he receives the report. The disadvantage is that frozen sections are much less satisfactory for diagnostic purposes than paraffin sections. When the surgeon is reasonably certain of the diagnosis on clinical grounds, he sends the specimen for examination by the paraffin method after the operation; it is when he is in serious doubt, having begun the operation, that he would most like a report before proceeding. But when the surgeon finds it difficult to make a diagnosis, the pathologist often does so too, and finds it more than ever necessary to study a paraffin section before expressing any opinion. When a biopsy is not feasible before operation, the surgeon should normally act solely on the clinical data, without interrupting the operation for a frozen section to be made, and send the specimen afterwards for study in a paraffin section. An exception to this rule may be made by the neurosurgeon, since frozen sections of tissue (or squashes of tissue) obtained by needling during an operation may enable the pathologist to say whether a lesion is in fact a tumour. Work of this kind presupposes both special arrangements and special training in neuropathology and is therefore outside the scope of the general pathologist.

Cytodiagnosis

This means diagnosis from the appearance of isolated cells in such material as sputum, gastric contents, or uterine smears. The method is used chiefly for the diagnosis of malignancy. Its use has been advocated enthusiastically and perhaps uncritically, with the result that experienced pathologists often refuse to use the method at all or use it with considerable reservations. It is true that no preparation can compare with a good paraffin section of a piece of tissue for quality and reliability for diagnostic purposes. The pathologist who decides to use only such material ought not to be labelled conservative or out of date; he is, after all, thinking only of the best interests of the patient, and if he himself were the patient he would wish to be examined in the same way.

The one field in which cytodiagnosis is of proved value is that of gynaecology. Smears critically studied may be helpful in detecting early cancers of the cervix. It is not so much isolated atypical cells that are significant as the occurrence of atypical cells in large numbers; the preparation can then be regarded almost as a teased tissue, and the method does not differ radically from the examination of a section.

Cytodiagnosis has also been much used for the detection of bronchial carcinoma. I have examined many specimens of sputum for this purpose but have never been able to make a confident diagnosis of malignancy. Since the thoracic surgeon can often obtain a piece of tissue without much difficulty, and since malignancy can often be diagnosed with certainty even in a minute fragment of material, it is uneconomic to consume the time of technicians and pathologists in innumerable sputum examinations yielding negative or inconclusive results. The average pathological laboratory is already fully occupied with ordinary biopsies; if a routine cytodiagnosis service is required in addition, the hospitals will have to provide extra facilities, technicians, and pathologists, and to pay for the special training required—expenses that are certainly not warranted by the results obtainable.

Grading of Tumours

The pathologist is sometimes asked to grade tumours numerically according to the degree of their structural difference from the corresponding normal tissue. There is undoubtedly an infinite series of gradations in appearance between a normal tissue—epithelial or connective—and the most undifferentiated tumour derived from it, but so many factors other than tumour pattern may affect prognosis that the value of such grading is limited. Another difficulty is that we can usually examine only a small part of a tumour; the more tissue we examine, the clearer it becomes that there are often wide variations in the appearance of a tumour from point to point, even transcending the limits of recognized types. For example, a bronchial carcinoma, if thoroughly examined, may show features of several distinct varieties of this growth; estimates of the frequency of different varieties may thus depend on the extent of the examination. These remarks are equally true of carcinomas of the breast, and malignant tumours in general may be justifiably classified only according to the predominant pattern. While formal grading is not worth doing, the information of interest to the surgeon can be conveyed by skilful description in the pathologist's report, and more usefully, since the meaning of the report will be clear to all who may read it, and not only to those acquainted with a particular classification. A description of a tumour implies an assessment of the degree of malignancy, so far as this can be judged histologically; in this sense, tumours always have been graded by all competent pathologists for generations past.

Biopsy Reports

The object of a report is to help the surgeon, a fact that cannot be too often remembered by the pathologist. The writing of reports is an art that can be acquired only by practice, but some general principles are worth stating. The surgeon is busy: therefore reports should be brief. With many reports about many patients converging on him, he is not going to read a technical dissertation, without a heading, filling a page with single-spaced type. Every report should be headed by the diagnosis in two or three underlined words, and the text should normally require only a few lines of type, which should be double-spaced. The three or four salient points about a section can be got into such a report. When the result of the examination bears on prognosis or treatment, the report should make this clear; in particular, the pathologist ought not to leave the surgeon guessing whether he should perform a radical or conservative operation or no operation at all. When the pathologist himself is doubtful, he should discuss the case with the surgeon before writing the report; a short conversa-

tion often solves such problems. While a report must be truthful, it should also be tactful, so that an intelligent patient would not be unduly alarmed.

A rare or interesting lesion may call for an unusually full description, but long reports issued as a routine on all cases mean that the pathologist is incapable of selecting the salient points or of writing concise English. The worst possible reports are those that wobble helplessly between "benign" and "malignant," or consist of academic descriptions of phenomena from which no practical conclusion can be drawn. Most of us have probably written reports of this kind, but we learn in time to resolve our difficulties beforehand by consulting a fellow-pathologist.

Next article on Clinical Pathology.—"Tests of Gastric Function," by Dr. F. Avery Jones.

Refresher Course Book.—The first collection of articles in the Refresher Course for General Practitioners (fully revised) are available as a book containing 55 chapters, price 25s. Copies can be obtained either direct from the Publishing Manager, B.M.A. House, Tavistock Square, London, W.C.1, or from booksellers.

AMERICAN SURGEONS IN LONDON

CONCLUSION OF SECTIONAL MEETING

The sectional meeting in London of the American College of Surgeons concluded on May 19. The report of the first day's proceedings was published last week (p. 1203). During the three days of the meeting 64 papers were read, 32 by American and 32 by British authors. The meeting on May 18, presided over by Dr. F. A. COLLIER (Michigan), took the form of a symposium on cancer.

Cancer of the Lung and Cigarette Smoking

Professor E. A. GRAHAM (St. Louis) presented the results of a survey of 600 patients with cancer of the lung. A questionnaire was devised for the purpose of finding out how far cigarette smoking was implicated, and each patient was asked the questions at an interview. The results were nearly identical with those brought forward by Professor Bradford Hill and Dr. Richard Doll in this country. A remarkable relationship between excessive cigarette smoking and lung cancer was demonstrated, and the same was true of six additional studies carried out in the States. Atmospheric pollution had been advanced as an alternative cause, but this would not explain the sex incidence. The disease was five or six times as prevalent in men as in women. Any carcinogenic agent in the atmosphere would affect both sexes equally. On the other hand, it took 20 or 30 years of cigarette smoking to establish cancer, and large numbers of women had not been smoking for that length of time.

Mr. HOLMES SELLORS (London) described the results of surgical treatment of cancer of the lung. There were two methods of treatment available: the surgery of excision and radiotherapy. Results with the latter had been unfavourable, though the picture might be different with improved technique and apparatus. Age was not necessarily a contraindication to surgical excision; he had operated on many patients in their seventies, and his oldest patient was 84. He showed a table of results of 700 cases from three hospitals. At the end of two years after operation the proportion of survivors was 45%; after three years, 30%; after four years, 22%; and after five years, 20%. The most favourable type of cancer for operation was the medium-sized central squamous-celled carcinoma; 72% of the patients were alive after three years, whereas with "oat-celled" cancer the figure was 40%.