SURGICAL PROCEDURES IN GENERAL PRACTICE

This is one of a series of articles contributed by invitation

SALINE INFUSIONS

BY

HAMILTON BAILEY, F.R.C.S.

Rectal Saline Infusion

Rectal saline infusion has the advantage of simplicity; it requires neither special apparatus nor asepsis. A drachm of salt to a pint of water at a temperature of 115° F. is allowed to gravitate into the rectum through a funnel and tube connected to a catheter. The patient's hips are elevated, and the funnel is held at a height of two feet. The usual amount of infusion is half a pint every four hours.

The celebrated John B. Murphy of Chicago introduced continuous rectal saline infusion (proctoclysis), which was a distinct advance, especially in cases of peritonitis. Slow continuous delivery of the saline aided its absorption, and with less disturbance to a patient in a critical condition. Murphy emphasized that the height of the reservoir should not be more than one foot above the rectum (Fig. 1), that the delivery tube should be of comparatively wide calibre, and that his glass bulb with many perforations (Fig. 1, inset) should always be employed. *It is to be noted especially that in Murphy's apparatus there is no valve or dripper in the circuit; thus flatus can be expelled up the wide rubber tube into the reservoir.* These simple principles have not been fully appreciated, and need further diffusion among the medical and nursing professions.

FIG. 1.—Diagram embodying the essential principles of proctoclysis as laid down by Murphy. Inset: Murphy's glass cannula; the part to the right of the dotted line lies within the rectum.

Many consider that the administration of saline by the rectal route satisfies almost every requirement. I have heard it said that if a mariner with "water, water everywhere, but not a drop to drink" were equipped with a funnel and tube he could absorb enough sea-water via the rectal mucosa to avert death for days or even weeks. Whether or not this is true, a close observer soon realizes

that rectal saline, even when given by an experienced and painstaking nurse, is sometimes returned, particularly in cases where it is most needed. I am compelled to agree with Willard Bartlett of St. Louis that whoever wishes to adopt saline by the rectal route exclusively should first administer it a few times in person, and watch the results.

Summarizing, rectal saline is simple, safe, but slow and not satisfyingly certain.

Subcutaneous Saline Infusion

The apparatus for administering saline subcutaneously consists of two sharp-pointed aspirating needles which are attached to some form of infusion apparatus by a Y-shaped glass connexion. The subcutis beneath the breasts is usually chosen for the injection. The skin is sterilized and the sterile needles are inserted with the saline already flowing. They are kept in place with adhesive tape. The fluid should be at a temperature of about 112° F.; it cools considerably while passing through the tubes. The container is raised to a height of three feet, and the saline must gravitate at such a rate that there is no undue swelling of the subcutaneous tissues. It is often possible to give two or more pints of saline by this method without removing the needles.

This mode of administration of saline is inclined to be painful; it cannot be continued for more than a few hours; and the tendency is to replace it by the more certain method of continuous intravenous infusion, which can be administered without discomfort and with great precision.

Intravenous Infusion

Given in a massive single dose (a pint or more) intravenous infusion has many times saved life. True, saline and other isotonic solutions are but a poor substitute for

FIG. 2.—Funnel and tube in readiness for an intravenous infusion. Some fluid has been allowed to escape from the nozzle in order to be certain that there is no air in the tube.

whole blood, but occasions arise when urgency forbids the delay inseparable from blood transfusion. When a patient is collapsed his veins will be in a like condition, and it is necessary to insert a cannula into the vessel.

Technique.—Before the vein is exposed the funnel and tubes should be filled with the solution. To ensure the absence of air in the tube and nozzle some fluid is permitted to escape. As soon as air-locks and bubbles are eliminated the tube is clipped (Fig. 2). A rubber catheter

is placed round the arm. If the veins in front of the elbow are not visible the venous blood is milked upwards from the wrist. A folded towel is placed beneath the elbow to hyperextend the joint. If the patient is conscious a little local anaesthetic is injected into the sterilized skin over the vein. A short transverse incision is made through the skin, and the beak of a small haemostat is introduced into the wound and its jaws opened widely (Fig. 3). If this manoeuvre is carried out once or twice the

FIG. 3.—A rapid and efficient method of displaying a subcutaneous vein through a small transverse cutaneous incision.

vein will be cleared from the subcutaneous tissues better than by painstaking dissection, and there is no fear of tearing even a delicate vein. Two catgut ligatures are passed beneath the vein; the lower is tied and the ends are caught in the haemostat. Using gentle traction on the latter, a small transverse nick is made in the vein wall. The anterior wall of the vein is held up (Fig. 4) until the nozzle (through which the saline is now running) is within the lumen; then the proximal ligature around the vein and the nozzle within it is tied. After the administration, as the last ounce of fluid is gravitating into the vein, the cannula is withdrawn and the ligature surrounding the vein is tied quickly. Two skin sutures are required to complete the operation.

FIG. 4.—A vein exposed at the fold of the elbow ready for the insertion of the cannula.

Continuous Intravenous Infusion (Venoclysis)

Useful as is the massive dose of intravenous saline the indications for its use are limited. It must be borne in mind that unless there has been a corresponding loss it is manifestly unphysiological to put into the circulation a pint of fluid more or less suddenly. In shock, dehydration from vomiting, oliguria, and many forms of toxæmia, a slow continuous flow of saline intravenously approaches the ideal, for we know exactly how much fluid the patient is receiving. The fluid can be given over a period of

days, and the rate of flow and the amount of fluid can be regulated with mathematical precision.

Continuous intravenous saline is valuable in the treatment of shock. In these cases an ampoule of coramine may be added to each pint of the saline solution with advantage. French observers have reported favourably on the results of adding adrenaline in doses of 5 minims of 1 in 1,000 solution to each pint.

Preparation of the Solution.—Obviously a solution which is introduced directly into the circulation must be absolutely sterile. But this is not enough: it must also be free from dead bacteria and other extraneous matter, such as fur from the sterilizer. Boiled tap-water does not fulfil these requirements, and reactions are bound to occur if it is used. Triply distilled water not more than a week old must be employed in all cases. It is, of course, equally important to have the sodium chloride sterile and pure. Tablets sold in tubes for the preparation of normal saline are not manufactured with a view to intravenous therapy. Sterile ampoules for making the solution are obtainable.* Leading manufacturing chemists now put up sterile normal saline in sealed litre bottles. When there is difficulty in obtaining fresh distilled water, and the facilities for sterilizing the somewhat bulky apparatus are not perfect (as in a private house), saline in this form, though comparatively expensive, has obvious advantages.

Saline or Saline and Glucose?—Normal saline solution alone meets most requirements. When glucose is employed we are introducing a substance into the circulation which requires katabolism, and if glucose is to be used there should be adequate reasons for its introduction: the patient must be in need of glucose, as one who has been starved, and only enough glucose for his immediate requirements should be introduced by the intravenous route. As regards insulin, if there is no reason to think that there is a deficiency of this hormone there is no need to give more.

Apparatus.—Special needles and cannulae for intravenous infusion are available. The ones shown in Fig. 5

FIG. 5.—Author's needle and cannulae. A, Needle for insertion into a vein by puncture; B, Cannula; C, Cannula, child's size. All the above are gold-plated to prevent corrosion. D, An interceptor for continuous intravenous saline.

will be found to be satisfactory. Every portion of the apparatus (see Fig. 7) must be freshly boiled and washed through with sterile water or saline before the reservoir is filled. New rubber tubing should be boiled, stretched, and washed through to remove chemicals, and boiled again before the apparatus is assembled. An interceptor—Fig. 5 (D)—is an integral part of the armamentarium; it should be freed of contained water from the sterilizer before the reservoir is filled. The whole apparatus must

* The Crookes Laboratories, Park Royal, London, N.W.10.

be seen to be in working order and the filled reservoir suspended suitably before any attempt is made to prepare the patient for the reception of the saline.

TECHNIQUE

Choice of the Vein.—In general it is best not to use a large vein; one just a little larger than the cannula or needle is most suitable. A vein of the forearm is usually chosen. If no veins can be seen an incision should be made over the beginning of the internal saphenous vein; even in newborn infants one never fails to find this vessel just anterior to the internal malleolus. In patients who are restless, or may become so, the leg as a site for the infusion is preferable, for the lower limb can be splinted securely.

Inserting a Cannula.—In many emergency conditions the veins are collapsed; consequently it is necessary to expose the vessel. This is done through a small transverse incision, as already described. Full aseptic ritual must be observed, in the preparation both of the patient's skin and of the operator's hands; it cannot be stressed too often how vital it is for everything to be aseptic in this procedure. For subsequent smooth running it is recommended to flood the wound and moisten the cannula with citrate solution immediately before inserting the cannula into the vein. Once the cannula is satisfactorily in position the wound is closed about it. The cannula is now fixed to the skin by means of fine stitches passed through the slots near its base (Fig. 6).

Inserting a Needle.—If a needle, as opposed to a cannula, is chosen, it is a mistake to take the most prominent vein. One that is easily palpable but well obscured and supported by subcutaneous fat will be found to be less mobile and thus more readily entered. The large prominent veins of the aged are notoriously difficult to enter, and in these cases it is as well to transfix the vein transversely with a fine hypodermic needle before carrying out the puncture. The special needle is unfastened, the base being left attached to the tubing of the apparatus (Fig. 7). The needle is then placed on a Record syringe and a small amount of citrate solution is drawn up through the needle into the barrel. After a tiny nick has been made in the anaesthetized skin with a scalpel puncture of a vein is carried out, and this will be found easier with the syringe attached to the needle. In order to be certain that the vein has been entered blood is drawn up into the syringe, and a little citrate solution is then injected into the vein. The syringe is removed and the base of the needle connected to the adaptor at the end of the tubing. The needle is kept in position by suitable strips of adhesive plaster passed over the wings of the needle.

FIG. 6.

Keeping the Limb at Rest.—For the arm a posterior plaster strip well padded with Gamgee tissue with a turn around the wrist forms an ideal splint readily adapted to each individual patient. For the leg a posterior splint with a footpiece is very serviceable (Fig. 8). Alternatively, a Thomas's knee-splint can be used; this has the advantage of ensuring rigidity of the limb if the patient is inclined to be restless.

The Temperature of the Fluid.—In spite of the many ingenious devices that have been put forward for keeping the fluid warm the simplest appears to be the best—namely, the patient's leg or arm into which the saline solution is flowing is kept warm with a hot-water bottle

FIG. 7.

or an electric heating pad. In addition to simplicity this leaves the reservoir and the delivery tube in full view.

Rate of Flow.—The average rate of flow for an adult in need of fluid should be fifty drops per minute—that is, a quarter of a pint per hour, or six pints in the twenty-four hours. In urgent cases, for the first hour the rate of flow is often accelerated to 100 drops per minute to see how the patient responds. If the blood pressure is increased satisfactorily we are encouraged to proceed with the measure. When in doubt as to how much fluid the patient should receive the flow should be cut down to thirty drops a minute—that is, approximately three and three-quarter pints in the twenty-four hours—a dose which, at all events, is unlikely to harm him.

The Administration.—The actual administration is largely in the hands of the nursing staff. It is of paramount importance to be sure that the nurse appreciates the simple yet vital responsibilities connected with the care of the apparatus and the maintenance of strict asepsis. The nursing instructions are simple, and can be carried out by any conscientious State-registered nurse providing that she receives special instructions if she has not been trained in the method.

FIG. 8.—The apparatus in use. In very ill patients who are restless or may become so the limb can with advantage be placed in a Thomas's splint. (This apparatus is made by the Genito-Urinary Manufacturing Co. Ltd.)

NURSING INSTRUCTIONS

1. Every utensil used in the transfer of the saline to the reservoir must be absolutely sterile.
2. To prevent the entry of bacteria from the air the top of the reservoir must be covered with sterile gauze.
3. The reservoir must always be kept more than half full of saline.

4. Veins are easily obstructed. There should be no bandage over the line of the vein.

5. The rate of flow should be timed and recorded at frequent intervals.

6. If the flow stops: (a) do not pinch the tubing; (b) see if the tubing is kinked; (c) if the flow is not restarted by some simple adjustment of the limb or the tubing report the matter at once.

7. Watch for and report immediately: (a) rigors; (b) redness along the vein; (c) oedema of the feet, face, or arms; and (d) any sign of respiratory distress.

8. Measure the amount of urine the patient has passed. If the output is less than the intake report the matter. Also report if the specific gravity becomes low.

Making up a Balance Sheet.—Unless a balance sheet is made up every twenty-four hours there is no check on the patient's requirements. I am definitely of the opinion that without a balance sheet intravenous saline should never be continued for more than twenty-four hours. To run in pints of fluid without keeping a check on its disposal is extremely dangerous. The balance sheet reproduced here is one that is extensively used.

CONTINUOUS INTRAVENOUS SALINE BALANCE SHEET

Patient's Name.....

24 hours ending	INTAKE	OUTPUT
..... 19		
INTRAVENOUSLY:		URINE: (S.G. .... a.m.) .. pts.
* @ ..... drops per min. pts.		(S.G. .... p.m.) .. pts.
* @ ..... " " pts.		Vomitus .. .. pts.
* @ ..... " " pts.		Normally 1½ pints—
BY MOUTH		Faeces .. .. pts.
Total		Sweating and Lungs—say .. pts.
		Total .. ..
GLUCOSE:—		
5% Solution (allow 30		
grams = 120 calories		
for each pint) ..... calories		
* 50 drops per minute =		
6 pints in 24 hours		
* 30 drops per minute =		
3½ pints in 24 hours		

CONTRAINDICATIONS

Continuous intravenous saline, employed intelligently, is an extremely valuable addition to the treatment of many urgent surgical conditions. The contraindications to its use are few, but they are definite and should be noted carefully.

1. *The Failing Heart.*—A history of dyspnoea on exertion, uncompensated valvular disease, or any possibility of cardiac weakness should call for hesitation in increasing the bulk of circulating fluid. When in doubt as to whether the poor pulse is due to cardiac weakness or to another cause, I have many times placed the patient on continuous intravenous saline and coramine, and waited for an hour or more to see if the blood pressure improves. If it does so, the flow is reduced to thirty drops per minute and the patient is again visited after a lapse of an hour or so.

2. *Pulmonary Congestion.*—In all conditions where there are signs of oedema or consolidation of the bases of the lungs the method is inadvisable.

3. *Hypertension.*—If the blood pressure is high it is obviously courting danger to burden the circulation still further.

4. *Bright's Disease* in all its stages is a grave contraindication. On the other hand, in obstructive anuria and

oliguria, after the obstruction has been removed, and in many of the surgical uraemias, continuous intravenous saline or, better, continuous intravenous sodium sulphate (made by dissolving 42.85 grammes of Glauber's salt in one litre of water), used with discretion, is a real therapeutic advance.

RESEARCH IN LUNACY AND MENTAL DEFICIENCY

REPORTS TO THE BOARD OF CONTROL

The Board of Control issues its annual report in two parts. The first of these, for 1936, dealing mainly with administration and statistics, was reviewed in our issue of November 6, 1937 (p. 914). The second part,¹ a volume of 540 pages, which would be improved by an index, embodies brief records of research from mental hospitals. In appendices the Commissioners give an account of their visits to the various institutions. These accounts turn mostly on small details of administration, but this makes them none the less useful to those concerned. A list of institutions and approved nursing homes completes the volume.

A slight note of disappointment is struck because the scientific reports come from only forty-eight out of the seventy-five county mental hospitals, and from only fifteen of the twenty-five county borough mental hospitals, and in a number of these cases the communication is limited to a list of routine tests and examinations. But it is recognized that in several institutions the medical staff is numerically small, and it may happen that all available time outside daily duty in hospital has to be spent at out-patient centres. Eight hospitals, by the way, report on their out-patient work, one of them—the Isle of Wight county mental hospital—commenting at length upon its four mental welfare clinics, which 211 patients attended during the year.

But even lists of tests have an intrinsic value of their own, and their scope appears to increase. Attention is drawn to a very complete series of tests at Northampton county mental hospital which have been arranged for every newly admitted patient. In addition to the ordinary clinical examination and inspection by the dental surgeon and ophthalmologist, laboratory tests are made of the blood, urine, the cerebrospinal fluid if necessary, and the faeces, and an examination of swabs and sputum. This routine has proved of real therapeutic value by indicating lines of treatment, and has been the starting-point of other special investigations. It is mentioned that it is impossible to gather from the comparatively few reports on the subject whether or not the Wassermann test is applied to every new admission. At St. Andrew's, Northampton, reliance is placed upon the Meinicke reaction for syphilis; the Wassermann is not used at all, and in the opinion of the reporters there it should have been abandoned years ago.

Chronic Infective Processes

In the forefront of the report stands a long and important communication from the Joint Board of Research for Mental Disease (City and University of Birmingham). It is devoted mainly to the question of chronic septic foci in the head and neck as related to disturbance of the capillary circulation in the brain. A description is given of twenty-eight cases illustrating the relationship of mental disorder and oronasopharyngeal sepsis. The series generally illustrates the value of the removal of septic or irritative areas and the conversion of "closed" into "open" sepsis.

Parallel with this is an account from the Warwickshire and Coventry mental hospital of the treatment of 200

¹ H.M. Stationery Office. (8s. net.)