hospital treatment, may result in cure even in such apparently dangerous cases (Plate, Fig. 2). Prognosis depends entirely on the stage at which diagnosis is arrived at, and thus every effort must be made to search out these cases and to recognize at a very early date the fact that they have been infected. The primary infection will show no clinical signs typical of the condition, and diagnosis can only be made on the grounds already mentioned.

Treatment

First and mainly treatment consists in removal of infants from all possible tuberculous contact. They should not be put in sanatoria where there are cases of open phthisis. They should be taken to infant hospitals or preventoria, and the isolation must be maintained until healing is evidenced by calcification or by fibrosis. Calcification is the less usual end-result, because if infants have caseating foci there is nearly always a fatal termination. Fibrosis is the commoner end-result; and here in later radiographs nothing abnormal is seen, and the only proof of past infection is the serial radiographs and a positive tuberculin test. Healing is established in mild cases in six months, but other cases may take up to two years. The falling of the sedimentation rate to normal also indicates healing. The second essential of treatment is rest in bed. Movements in the cot have to be permitted, but exercise out of the cot must be postponed until serial radiographs and normal sedimentation rate and weight gain show that the lung is healed. Fresh air is essential throughout treatment, but infants must not be exposed beyond their limits of endurance, and throughout treatment they must be kept warm. Digestive disturbances must be attended to and a suitable diet rich in vitamins, with added calcium, must be given. Ultra-violet rays and strong sunshine are absolutely contraindicated in all pulmonary lesions in infants and children; however, in these climates outdoor shade in summer and exposure to sun in spring and autumn are safe. Temperature charts, the sedimentation rate, and radiographs all demonstrate how dangerous is sun-bathing to a small child with a pulmonary lesion, be this only in the primary stage. It will be found that casualties occurring during the primary stage of pulmonary tuberculosis will be among those infants whose parents are not convinced, and who do not allow this treatment to be carried on until complete healing has been ensured.

Summary

1. Seventy-eight cases of tuberculosis of the lungs are reported in infants under 1 year old: 23 per cent. recovered.
2. Healing can only be achieved by early diagnosis—that is, in the primary stage.
3. Early diagnosis can only be made by tuberculin skin test, radiography, and history of contact.
4. Treatment consists mainly in immediate removal from contact and in rest.

D. Bobeff (*Mösch. Kinderheilk.*, October 8, 1937) records the case of a Bulgarian girl, aged 10, who on the sixteenth day of a severe attack of faucial diphtheria, six days after the onset of palatal palsy, developed gangrene of both legs, most marked on the right side. The necropsy three days later showed thrombosis of the abdominal aorta, multiple infarcts in the left kidney, gangrene of both feet, and myocarditis. Bobeff has found only two other cases on record of occlusion of the abdominal aorta following diphtheria.

AN UNUSUAL TUBERCULOUS APPENDIX

BY

D. J. HARRIES, M.D., D.Sc., F.R.C.S.

AND

EMRYS WILLIAMS, M.D., M.R.C.P.

Cardiff

(WITH SPECIAL PLATE)

Although tuberculosis of the appendix is not an uncommon disease it is still the exception for it to be mentioned in standard textbooks of medicine or surgery. It has been stated that from 1 to 3 per cent. of all appendices removed at operation are tuberculous. It is obvious that an investigation into this problem at a large hospital would entail an enormous increase in the microscopical work of the institution, and this might be regarded by the pathologists as out of proportion to its value. An investigation carried out by Pagel and Weichherz (1936) at Papworth Village Settlement, based on eighty-four necropsies on tuberculous subjects, showed that roughly one-third had no intestinal tuberculosis; another third had intestinal tuberculosis with no involvement of the appendix; the remaining third had involvement of both the intestine and the appendix. In three cases the appendix was the only part of the gut to be involved, and in each there was extensive tuberculosis in the lungs.

The appendix is almost always involved in ileo-caecal tuberculosis. It is generally recognized that tuberculosis may primarily involve the intestines, but it is not so certain that the appendix alone can be the site of tuberculosis. In view of the above remarks it requires some courage to suggest that the case described below is one of primary tuberculous appendicitis, successfully treated by operative removal, with the patient alive and perfectly well nearly two years later.

Case Record

On March 18, 1936, a marine engineer, aged 19 years, had a slight attack of abdominal pain lasting about an hour. It was so mild that the patient did not trouble to obtain medical attention, and in fact went on with his work. On March 23 he had a recurrence of the pain, and when seen by one of us (E. W.) was sent into a nursing home with the diagnosis of acute appendicitis. On admission the temperature was 104° and the pulse 90. There was very little rigidity but some tenderness in the right lower abdomen. On account of the comparative absence of rigidity it was possible to palpate a large tubular structure, about an inch in diameter, running from the right iliac fossa into the pelvis. It felt like an indurated coil of intestine or an intussusception. The patient and his mother were quite definite that he had never been ill before.

Operation.—This was carried out within three hours of the onset of symptoms by D. J. H., the anaesthetic being given by E. W. On opening the abdomen it was seen that the tumour was an enormously enlarged appendix, its peritoneal coat being dark red. It felt perfectly solid and hard, and there was no constriction at its junction with the caecum. After ligating its mesentery the appendix was removed intact and the raw area left on the caecum covered over by making use of the ileo-caecal fold of peritoneum. There was some induration of the wall of the caecum for about half an inch beyond the actual area to which the root of the appendix was attached, and if the tuberculous nature of the appendix had then been known this area would have been resected.

Subsequent Progress.—The patient made a rapid recovery, the temperature falling two degrees a day, and was normal on the third day. It is curious that the pulse never exceeded 90 at any time during the illness. The incision healed by

first intention and the patient went home on April 8, a fortnight after the operation. He has been seen and examined several times since. In February, 1937, he served as second engineer on a trip to the East, and returned in June. He was then carefully examined by us, but nothing abnormal could be found in the chest or abdomen. In fact he felt and looked perfectly well.

The Specimen.—Before hardening in formalin this was six inches long and one inch in diameter, and slightly curved. It was quite solid and hard, with a small black spot representing the original lumen. The cross-section at the root was greyish pink, with no evidence of necrosis or suppuration—in fact, the only sign of inflammation was in the peritoneal coat, which was dark red. The specimen, which has been placed in the museum of the Royal College of Surgeons, is now smaller and the colour of the peritoneum has faded. Fig. 1 (see Plate) shows the size and shape after hardening and mounting. Fig. 2 gives the microscopical appearance of a section taken from the cut surface at the root. A few giant cells can be seen in a mass of lymphoid tissue. The various layers normally found in the wall of the appendix have been completely replaced by tuberculous tissue, the peritoneum alone having survived.

Comments

It is unnecessary to state that the exact nature of the change in the organ was not realized until the microscopical slides were examined. The specimen was unlike any appendix we had seen previously. It is obvious that the disease must have started months, if not years, before it was discovered. The absence of symptoms until within three hours of operation allowed us to remove the specimen before it had been damaged by inflammatory processes. We are of the opinion that the acute symptoms were caused by interference with the blood supply, as there was no visible lumen which could have been infected. At the operation the other abdominal organs were looked at in the ordinary way, but nothing abnormal was found. It must be admitted that small tuberculous lesions were not specially sought, as the diagnosis of tuberculosis had not then been suspected. But in view of the subsequent history of the patient it is reasonable to assume that there could not have been any well-developed tuberculous changes in the intestines or lungs. He returned to work in a few weeks, and has since enjoyed perfect health.

We are indebted to Mr. W. J. Salter, technician to Llandough Hospital, for the photographs.

REFERENCE

Pagel, W., and Weichherz, E. (1936). *British Medical Journal*, 2, 1305.

La Migraine Ophtalmique, by Drs. G. Renard and A. P. Mekdjian (the publishers are Masson of Paris and the price is 22 francs) gives a convenient summary of what is known of migraine, well set out under carefully designed subheadings. No unifying conception emerges, the authors stressing the multiplicity of associated disturbances, and dealing in particular with the allergic basis of the affection. One point of interest which they bring out is the disturbance in blood pressure, which they studied both from the general aspect and by ophthalmodynamometry of the retinal vessels. In general they found a tendency to hypotension during the attacks. Their thesis is illustrated by some case reports. They argue that hypotension can be taken as a diagnostic feature distinguishing true migraine from the transient blotting out of vision seen in elderly people with hypertension. They regard migraine as a condition which, though distinct from epilepsy, has in common with it the fact that there is a widespread, though ill-defined, underlying general departure from the normal, both psychological and somatic.

PELVIMETRY AND THE TRUE CONJUGATE

BY

JOHN HOWKINS, M.D., M.S., F.R.C.S., M.C.O.G.

Obstetric Registrar, the Middlesex Hospital

Every obstetrician must from time to time have been asked to see ante-natal cases in which a practitioner or midwife has felt on vaginal examination the sacral promontory; and there must be a number of women who are, because of this genuine misconception, submitted to unnecessary obstetric operations, with a diagnosis of gross pelvic contraction.

While not decrying the academic importance of attempting to measure the true conjugate clinically, it is the purpose of this brief communication to draw attention to the extreme rarity with which the sacral promontory can be palpated per vaginam; and my excuse for burdening an already adequate literature is that in the course of the last twelve months' ante-natal work three cases have been referred to the Middlesex Hospital Ante-natal Clinic labelled with such a phrase as: "Sacral promontory felt per vaginam: ? Caesarean section for pelvic contraction." All three were subsequently delivered by the natural passages. The only explanation that answers this fallacy is that the "sacral promontory" is in reality one of the lower bodies of the sacral vertebrae.

Three Factors to be Considered

Whether or no the sacral promontory is palpable depends on three factors: first, the length of the true conjugate; secondly, the effective length of the examining finger; and, thirdly, the rigidity of the vaginal introitus, which largely depends on the co-operative relaxation of the patient and her parity.

The anatomical true conjugate on the average female dry bony pelvis measures between 4.25 and 4.5 inches: the distance of the point behind the symphysis, from which the true conjugate is measured, to the subpubic angle is 1 inch on the bone, and must be nearly 50 per cent. greater than this when measured to the anterior lip of the vaginal introitus, in order to take into consideration the triangular ligament and its associated superficial small muscles and soft structures. This latter is greatest in the primipara and least in the multipara, for obvious reasons. Applying the well-known principles of geometry, the total effective length of an examining finger must be roughly 4.75 inches to feel the promontory in a normal pelvis.

The effective length of the examining finger, assuming that the index and middle fingers are used together, varies slightly with different surgeons, but surprisingly little. The determining factor here is largely the resistance encountered by the ring finger knuckle against the perineum, and to increase this useful length the perineum must be pushed upwards and backwards. Discounting this last factor, the uncompensated effective length of one's fingers is about three and a half inches. In a normal pelvis, therefore, with a true conjugate of 4.5 inches, the finger must be lengthened by depressing the perineum at least 1.25 inches before the promontory is reached, or, conversely, the pelvis must be foreshortened down to a true conjugate of 3.5 inches before the tip of the middle finger is likely to meet the posterior bony landmark. The exact location of the promontory is, moreover, a matter of difficulty, the manoeuvre of sliding the finger up the

D. J. HARRIES AND EMRYS WILLIAMS: AN UNUSUAL TUBERCULOUS APPENDIX

FIG. 1.—Photograph showing size and shape of appendix after hardening and mounting.

FIG. 2.—Microscopical appearance of a section taken from the cut surface of the base of the appendix.

H. BARSOUM: A CASE OF CHLOROMA

Showing the dura, seen from the inside with the tumour-like deposits.

W. A. BELLAMY: ABNORMALITY OF RIBS

Antero-posterior view showing ribs to be thinner and less calcareous on the right than on the left.