syndrome or, to use the transatlantic term, from neuro-circulatory asthenia. But from the all-important point of view of treatment the problems are not so essentially different. A short rest will be suitable for both; reassurance will help both, as also will suitable exercise and retraining. When it is admitted, as most people will, that in the effort syndrome group a man must sometimes be warned that he is not suited for intense or prolonged effort the practical considerations are identical. The divergence arises on account of compensation litigation on the one hand and of theoretical prejudgment as to whether the normal heart can be strained on the other.

The American teaching (White, 1937) seems to be that heart symptoms or signs that follow industrial strain or accidents are usually of the neuro-circulatory asthenia type, or are due to aggravation of heart trouble already existing. But Donahue (1927), in an article entitled "Accidents and Heart Disease from the Court's Point of View," writes: "We do have a great many cases of dilatation of the heart occurring at work while the employee is under some particular strain. He is lifting some unusually heavy load, or is in a rather awkward position doing heavy lifting, when he has dilatation and becomes sick, sometimes collapsing immediately." If we leave out the word "dilatation" the events are as described. Probably some are due to rupture of pectoral muscle fibres, with a cardiac neurosis superimposed. Probably some are due to the effort syndrome, to which the event has called attention. But the probability still remains that some are due to strain of the heart muscle or of the pulmonary or aortic ring. It would be in the best interests of medical science if such cases came before a medical board. We could study them with a mind more open. In those industries where there is a medical adviser to the firm he should include judicious management of the case in his decision.

We can remember that a generation ago sometimes the strained heart was diagnosed, quite mistakenly, on the evidence of a negligible systolic bruit in a growing boy. Invalidism on this account is a thing of the past. The profession has learned that the effort syndrome requires guidance and retraining, lest a selfconscious heart become permanent. Much semi-invalidism has been prevented by the accepted teaching that the heart is the part least likely to break down from exhaustion in some athletic event.

Our knowledge of physical signs and of their interpretation has advanced. There remain, however, some problems which cannot be solved by these means, and the last word must come from reliable clinical histories. By such histories, from doctors who have made notes, I believe enough evidence may be collected to prove that the healthy heart can be strained.

Summary

The forms of heart disease which have been recorded as the result of direct violence to the chest wall or as the result of strain are discussed. The conclusion is drawn that the physical signs on examination are of little assistance in assessing the diagnosis of trauma. Reliable clinical histories, in the widest sense of those terms, must be assessed with judicious care. This includes such evidence as the patient's own doctor can supply.

The clinical diagnosis of a contusion of the heart is discussed.

It is claimed that there is clinical evidence that the normal heart may become diseased as the direct result of overstrain from effort.

With regard to these two conditions, contusion of the heart and primary cardiac overstrain, it is difficult to obtain the proof that the symptoms in question are due to a genuine heart disability, but the probabilities in some cases are sufficient to justify the diagnosis.

BIBLIOGRAPHY

- Allbutt, T. Clifford (1870). *St. George's Hosp. Rep.*, **5**, 23.
 — (1873). *Trans. Clin. Soc. Lond.*, **6**, 101.
 Barber, H. (1912). *Practitioner*, **89**, 230.
 — and Osborn, G. R. (1937). *Guy's Hosp. Rep.*, **87**, 510.
 Brockbank, E. M. (1926). *Incapacity or Disablement in its Medical Aspects*, London.
 Coffen, T. H. (1930). *Amer. Heart J.*, **5**, 667.
 Fraser, A. J. (1929). *Trauma, Disease, Compensation*, p. 100, London.
 Gibson, G. A. (1909). *System of Medicine*, Allbutt and Rolleston, **6**, 381.
 Goodall, J. S., and Weir, H. B. (1927). *British Medical Journal*, **1**, 834.
 Gordon, W. (1922). *Practitioner*, **109**, 223.
 Groom, W. (1897). *Lancet*, **1**, 1202.
 Gunewardene, H. O. (1934). *British Medical Journal*, **2**, 942.
 Hamilton, J. A. (1934). *Ibid.*, **2**, 1101.
 Hay, J., and Jones, H. W. (1927). *Ibid.*, **1**, 559.
 Horton-Smith, P. (1902). *Clin. J.*, **19**, 254.
 Kahn, M. H., and Kahn, S. (1928). *Ann. intern. Med.*, **2**, 1013.
 Lewis, T. (1932 ed. and 1937 ed.) *Diseases of the Heart*, London.
 O'Neill, B. J. (1914). *J. Amer. med. Ass.*, **62**, 697.
 Peacock, T. B. (1865a). *Croonian Lectures*, London.
 — (1865b). *Trans. Path. Soc.*, **16**, 67.
 Price, F. W. (1937). *Text-book of the Practice of Medicine*, London.
 Smith, K. Shirley (1932). *British Medical Journal*, **1**, 129.
 Smith, S. A. (1931). *Med. J. Austral.*, **2**, 575.
 Walker, G. F. (1933). *The Injured Workman*, Bristol.
 White, P. D. (1937). *Heart Disease*, New York.
 Wilkinson, K. D. (1930). *British Medical Journal*, **1**, 360.
 Wilks, S. (1865). *Trans. Path. Soc.*, **16**, 77.
 Wilson, C. (1930). *British Medical Journal*, **1**, 189.

CHANGES IN THE LIVER OF MALE RATS AFTER CASTRATION AND INJECTIONS OF SEXUAL HORMONES

BY

KATHLEEN HALL, Ph.D.

AND

VLADIMIR KORENCHEVSKY, M.D.

(From the Lister Institute, London)

Normal life of the organism is dependent to such a large extent upon the various functions of the liver that any changes observed in this organ need to be very carefully investigated. The manifold effects of castration on the different organs, including the liver, of male rats have been described in a previous paper (Korenchevsky and Dennison, 1934a). In most cases about seventy days or more after castration the actual weight of the liver was on the average 18 to 22 per cent. less than that of the control normal litter mates. When the weights were calculated per unit of body weight this decrease was not so great—namely, 11 to 18 per cent.—owing to the fact that castration also caused a decrease in the body weight of the rats. Histological investigation of paraffin sections did not show any considerable changes, although in some cases the cells of the "castration" liver were more vacuolated than those of the normal control (Korenchevsky and Dennison, 1935).

The injections of purified extracts of sexual hormones from male urine (Korenchevsky, Dennison, and Kohn-Speyer, 1933) into castrated rats were followed by an increase in the weight of the liver up to 32 per cent. (actual) and 25 per cent. (per unit of body weight). Histologically (Korenchevsky and Dennison, 1935) no difference

was found as compared with the controls. Oestrone (Korenchevsky and Dennison (1934b, 1934c) gave more uniform results when injected into normal than into castrated rats. In most cases oestrone caused a slight decrease in the actual weight of the liver, but no change, or sometimes even a slight increase, in that per unit of body weight. The fall in actual weight of the liver was due to the decreased appetite of oestrone-injected rats, with resulting decrease of body weight and of most of the organs, and not to any specific action of the hormone on the organ. Histologically no changes were found, except some increase in vacuolation of the liver cells (Korenchevsky and Dennison, 1935).

Ito and Kon (1935a and 1935b) injected purified extract of sexual hormones from male urine into infantile and adult male rats, and obtained hypertrophy of the liver in the immature rats up to 100 per cent. Bates, Riddle, and Lahr (1937) observed an increase of about 60 per cent. in weight of liver after androsterone injections, and a decrease of about 26 per cent. after oestrone injections, into adult pigeons. It is also important to note that the sexual hormones are accumulated in the liver and excreted in the bile (Gsell-Busse, 1929; Frank and Goldberger, 1930).

Present Experiments

The present paper is a description of the histology of the liver of those male rats in which the effect of injections of pure crystalline sexual hormones on the weights of this organ has already been described previously (see below). At the same time the uninjected normal and castrated control animals used in these experiments gave an opportunity for further study of the effects of castration. The hormones investigated were:

1. Androsterone, androstanediol, and their water-soluble esters (Korenchevsky, Dennison, and Levy Simpson, 1935).
2. Transdehydroandrosterone (Korenchevsky and Dennison, 1936).
3. Δ^4 -androstenedione and Δ^5 -androstenediol (Korenchevsky, Dennison, and Eldridge (1937a).
4. Testosterone (Korenchevsky, Dennison, and Brovsin, 1936; Korenchevsky, Dennison, and Hall, 1937).
5. Testosterone propionate (Korenchevsky, Dennison, and Eldridge, 1937b; Korenchevsky, Dennison, and Hall, 1937).

Technique

One hundred and eighty-one rats (fifty-one normal and 130 castrated) were investigated. Castration was usually performed at the age of 21 to 24 days, never later than 28 days—that is, always before sexual maturity. In this series of experiments, on the special mixed diet used, the changes in the liver produced by castration and injections became noticeable earlier than in the previous experiments—that is, at the age of about 70 days, about forty-five days after castration. Therefore in Table II the data for rats 70 to 150 days old only are summarized. The doses injected are shown in the same table. The hormones were dissolved in sesame oil and were injected daily, except the oestrogens, which were injected three times a week. Androsterone and androstanediol were injected in some experiments for periods up to fifty-three days, oestrone and oestradiol up to 120 days, and all the other hormones for twenty-one to twenty-three days.

Diet has a considerable influence upon the liver. In the previous experiments (Korenchevsky and Dennison, 1934a) the diet used was chiefly synthetic, while in the present

experiment the following diet of natural food products mixed into a paste was given:

Steamed herring, with bones (four times weekly)	1,020 grammes
White bread	1,350 "
Milk	1,700 c.cm.
Wheat	230 grammes
Oats (rolled)	230 "
Hemp	240 "
Wheat germ	75 "
Dried yeast	90 "
Salt mixture (McCullum's 185)	30 "

In addition each rat received twice a week 5 drops of cod-liver oil, 2 grammes of freshly minced liver, and 5 grammes of fresh carrot.

For histological investigation the liver was fixed in Bouin's solution, embedded in paraffin, and stained with haematoxylin and eosin. For fat and lipoids frozen sections were cut and stained with scharlach red. In order to compare the size of the cells, outlines of about seventy-five cells from one lobule of each liver were drawn with the aid of a camera lucida (magnification $\times 530$) from the central vein towards the periphery. The size of these outlines was compared directly, but because of their variability the number of the cells contained in a square measuring 7×7 cm. was also calculated, the

TABLE I.—Average Weights of Liver, actual and per 200 grammes of Body Weight, and Average Comparative Numbers of Lobules and of Cells of Normal ("N") and Castrated ("Ca") Rats.

Column :		I	II	III	IV	V	VI	VII	VIII	IX	X
Group	Age in Days	Number of Rats in Groups .		Weight of Liver (grammes)				Number of Lobules		Number of Cells	
				Actual		Per 200 gm. of Body Weight					
		N	Ca	N	Ca	N	Ca	N	Ca	N	Ca
1	40-49	6	—	8.3	—	10.1	—	6	—	53	—
2	60-69	5	6	10.5	10.9	8.6	8.5	6.6	6.4	52	60
3	70-79	4	10	12.3	9.5	8.7	6.6	6.1	7.1	58	62
4	80-89	7	9	12.6	10.1	7.8	6.9	5.7	7.1	60	61
5	90-99	12	11	13.1	10.5	7.4	6.5	5.1	6.4	54	57
6	100-150	3	4	13.1	10.4	6.9	6.8	5.2	6.3	57	52

TABLE II.—Average Weights of Liver and Comparative Average Numbers of Lobules and of Cells of Rats aged 70 to 150 Days.

Column :		I	II	III	IV	V
Group	Hormones Injected and Daily Dose (mg.)	No. of Rats	Liver			
			Weight in grammes		No. of Lobules	No. of Cells
			Actual	Per 200 gm. of Body Weight		
1	(A) Normal Rats : Control, uninjected	26	12.8	7.7	5.5	57
2	Testosterone 0.5-1.5	5	13.0	7.5	5.1	56
3	Testosterone propionate 0.5-1.5	9	13.1	7.9	5.4	56
4	(B) Castrated Rats : Control, uninjected	34	10.1	6.7	6.7	58
5	Androsterone 0.45, androstenedione 0.5-3	15	10.2	7.4	6.3	57
6	Androsterone 0.9-3.6, androstanediol 0.175-0.7, transdehydroandrosterone 1-4, testosterone 0.167-1.4, testosterone propionate 0.167-1.4	51	12.3	7.8	5.9	55
7	Oestrone 0.006-0.018, oestradiol 0.02	12	8.2	7.6	6.8	62
8	Oestrone 0.006-0.018 in combination with the hormones of group 6	12	9.5	7.2	6.9	62

Note.—The greater the number of lobules or cells in the measured area (columns IV and V) the smaller they are, and vice versa.

greater the number of the cells obtained the smaller being the size of the cells. These figures, being more reliable, are given in columns IX and X of Table I and column V of Table II. Actual measurements of the size of the lobules were practically impossible owing to their irregular shape and indistinct boundaries. Therefore the number of central veins of the lobules seen in six to eight microscopical fields at $\times 90$ was calculated, the average figure obtained being given in columns VII and VIII of Table I, and column IV of Table II; the greater the number the smaller the size of the lobules, and vice versa.

Liver of Normal Rats

The data are summarized in columns I, III, V, VII, and IX of Table I. Group 6 contains only a few rats, but the data are similar to those of our previous experiments (Korenchevsky and Dennison, 1934a). Because of the difference in the diet, and because of the different method of killing the rats at the end of the experiment (gas in the previous and bleeding in the present experiments), the liver weight was greater in the former case (Korenchevsky and Dennison, 1934a, p. 239).

Effect of Age.—As was found previously and confirmed in the present experiments, the liver belongs to that group of organs of which the actual weight (column III, Table I) slowly increases with age, while the weight per unit of body weight (column V) gradually decreases after sexual maturity. This seems to indicate a particularly active role of the organ in the young animal. The decline in weight of liver per unit of body weight after sexual maturity might be explained only partly by the accumulation of inactive tissue such as fat, and probably chiefly by the lowering of the metabolism. While the size and the histological structure of the cells show no definite changes with age (column IX), the lobules appear to become slightly larger (column VII), which may partly explain the mechanism of increase in the actual weight of the liver.

Liver After Castration

The data are summarized in columns II, IV, VI, VIII, and X of Table I, and should be compared with the corresponding data for normal animals in the other columns of the same table.

Effect of Age.—In agreement with our previous results (Korenchevsky and Dennison, 1934a) the actual weight of the liver (column IV) about four weeks after castration, in contrast with the normal animals (column III), ceases to increase with age. Moreover, the weight per unit of body weight considerably decreases at the age of about 70 days (column VI) and then remains more or less stationary (Groups 3 to 6). This, as in the case of the liver of normal animals, probably indicates a more active role also of the "castration" liver in the young animal. Histologically no definite alteration was found in the liver at different ages.

Effect of Castration.—Up to the age of 70 days no definite changes were noticed in the livers of the castrated rats, but in most rats older than 70 days the weight of the liver was less than that of normal rats (columns III and IV; V and VI), the difference being on the average 21 per cent. in actual weight and 13 per cent. in relative weight. This difference in weight could in most animals be at least partly explained histologically by the smaller size (22 per cent. on the average) of the lobules in the livers of most castrated animals (columns VII and VIII). The cells themselves, however, were not smaller (columns IX and X), which suggests their hypoplasia rather than

atrophy. No mitoses, however, were found in either normal or castrated livers, and the number of binucleated cells was about the same in both groups. No degenerative or any other considerable changes were found in the cells. In about 50 per cent. of the castrated rats the protoplasm of the liver cells in paraffin sections showed an increase of vacuolation (usually only slight) as compared with the normal animals (see also Korenchevsky and Dennison, 1934a, p. 333), but staining with scharlach red showed a definite increase in lipid granules in only a few cases. The absence of definite pathological changes in the "castration" liver, considered together with the decreased weight of the organ, suggests more a physiological than a pathological decrease in function of the liver after castration.

Effect of the Injections

The data are given in Table II—all were averaged after dividing them into eight groups. The figures in Group 1 (normal) and Group 4 (castrated rats) are the averages for the control uninjected rats aged 70 to 150 days.

Normal Rats.—Testosterone and testosterone propionate injected into normal rats (Groups 2 and 3) did not change appreciably either the weight of the liver or its histological structure (including content of lipid granules).

Castrated Rats.—Small doses of androsterone produced no definite effect, and Δ^4 -androstenedione, in the doses used, only a slight effect (Group 5). Larger doses of androsterone and all the doses indicated of androstanediol, transdehydroandrosterone, testosterone, and testosterone propionate (Group 6) caused in most cases a return towards or actually to normal of the weight (columns II and III) and histological structure (columns IV and V) of the liver. Since the increase was not only in the actual weight of the liver but also in that per unit of body weight, it is unlikely that this change was due to increased appetite and body weight, which usually occur in rats after these injections. Moreover, it is typical that in normal untreated animals greater appetite leads to increased gain in body weight with increased fat deposition. In the rats injected with the above hormones this was usually not the case, since an enlarged liver was present in rats with a decreased fat deposition and sometimes even with decreased gain in body weight (Korenchevsky and co-workers, 1935-7). It is probable, therefore, that these hormones exercise a special stimulating effect on the liver. Oestrogens, on the contrary, whether alone (Group 7) or in combination with the other hormones (Group 8), caused a decrease in the actual weight of the liver, while that per unit of body weight was slightly increased. As already described, this change may be explained by the loss in appetite caused by injections of the oestrogens. In "oestrogen" rats the gain in body weight and in some organs and fat deposition were usually decreased. The histological structure was similar to that of castrated uninjected rats. After injections of any of the sexual hormones used the liver cells, in some cases only, appeared to contain more lipid granules.

Summary

1. In normal uninjected rats the weight of the liver per unit of body weight decreases with age, which probably indicates an especially active condition of the liver in the young animal.

2. In rats castrated before sexual maturity the weight of the liver (actual and per unit of body weight) at the age of 70 days or more is less than in normal rats, although

histologically the only definite change is a slight decrease in the size of the lobules.

3. Injections of androsterone, androstanediol, trans-dehydroandrosterone, testosterone, and testosterone propionate returned the liver of most castrated rats towards or to normal weight and structure, while testosterone and testosterone propionate had no appreciable effect upon the liver of normal rats.

4. The absence of pathological changes in the small "castration" liver and in this organ enlarged after injections of the above sexual hormones, as well as the nature of the changes observed, suggests a natural stimulating action of these hormones on the liver.

5. This might have a practical significance in therapy of the liver and some metabolic disturbances if the same effect of these hormones be found in the case of human liver.

6. Injections of oestrone or oestradiol into castrated rats in the doses used caused a decrease in weight of the liver to below that of the control rats but no definite histological changes. The changes in weight observed are probably due chiefly to the depressing effect of oestrogens on the appetite, with resulting decrease in gain of body weight and weight of different organs, including the liver.

Grants from the Medical Research Council and from the Lister Institute have enabled us to carry out this work, and to them our thanks are due. We are also indebted to Professor A. Girard for oestrone and oestradiol, and to Messrs. Ciba, Ltd., in particular to Dr. K. Miescher, for all the other hormones used.

REFERENCES

- Bates, R. W., Riddle, O., and Lahr, E. L. (1937). *Amer. J. Physiol.*, **119**, 610 (Table II).
- Frank, R. T., and Goldberger, M. A. (1930). *Proc. II Intern. Congr. for Sex. Research*, p. 378.
- Gsell-Busse, M. A. (1929). *Arch. exp. Path. Pharmacol.*, **139**, 328.
- Ito, M., and Kon, T. (1935a). *Jap. J. Derm. Urol.*, **37**, 87.
- (1935b). *Ibid.*, **37**, 131.
- Korenchevsky, V., and Dennison, M. (1934a). *J. Path. Bact.*, **38**, 231.
- (1934b). *Biochem. J.*, **28**, 1474.
- (1934c). *Ibid.*, **28**, 1486.
- (1935). *J. Path. Bact.*, **41**, 323.
- (1936). *Biochem. J.*, **30**, 1514.
- and Brovsin, I. (1936). *Ibid.*, **30**, 558.
- and Eldridge, M. (1937a). *Ibid.*, **31**, 467.
- (1937b). *Ibid.*, **31**, 475.
- and Hall, K. (1937). *Ibid.*, **31**, 1434.
- and Kohn-Speyer, A. (1933). *Ibid.*, **27**, 557.
- and Simpson, S. Levy (1935). *Ibid.*, **29**, 2534.

The Royal Prince Alfred Hospital, Sydney, N.S.W., derives its name from H.R.H. Prince Alfred, Duke of Edinburgh, who, when visiting Sydney as Captain of H.M.S. *Galatea* in 1868, was shot and severely wounded while attending a charitable function. As a mark of their gratitude for his recovery the people of New South Wales raised a fund, which was applied to the establishment of the hospital. The fifty-third annual report covers the year ended June 30, 1937. During this period the number of beds was increased from 568 to 762 by the opening of Gloucester House, an additional building which makes available to all classes of the community, at reasonable rates, the resources of a large university hospital. In the same year of expansion the nurses' home has been doubled, a complete psychiatry pavilion with its own operating theatre, x-ray room, and special investigation facilities, including a neurosurgical unit of 15 beds, has been constructed, and a new x-ray department with the most modern equipment has been completed. In the course of the year under review 10,352 in-patients and 222,272 out-patients were treated in the public hospital, in addition to 1,724 in-patients at Gloucester House.

SCAPULO-HUMERAL PERIARTHRITIS (DUPLAY'S DISEASE)

BY

A. H. DOUTHWAITE, M.D., F.R.C.P.

Physician, Guy's Hospital

During the last three years I have seen thirty-seven patients (twenty-nine in private, eight in hospital practice) suffering from unilateral painful and stiff shoulder-joint associated with normal x-ray appearances. Although, as will be seen later, the antecedent factors leading to this state of affairs differ to some extent, yet the clinical picture is so constant and the treatment so well defined as to justify its description as a morbid entity. In searching the literature for evidence of earlier recognition of the disease, the first and most complete account was found to have been written by Duplay (1872), who recognized, however, only the post-traumatic forms. Mention of the rheumatic types has been made by Chaumet (1934), Douthwaite (1936), Tiegel (1936), and Sievers (1926).

Aetiology

Sex and Age.—There were fifteen males and twenty-two females. The average age was 60 years, and with the exception of two patients aged 33 and 34 they all fell into the fifth, sixth, and seventh decades.

Injury.—Trauma appeared to be the causative factor in twelve patients. In nearly every case it was the effect of a fall either upon the outstretched hands or upon the point of the shoulder. In one person a backward fall from a step-ladder resulted in bilateral disease. The lapse of time between the injury and the onset of typical symptoms—that is, those not simply due to bruising—varied from five days to three weeks.

Rheumatism.—Fibrositis either as simple stiff neck or more often in a widespread involvement of the shoulder-girdle preceded, or merged into, the symptoms of peri-arthritis in fourteen cases. Apart from the history it is quite impossible to distinguish between the two classes. Two patients had had recurrent gout, but whether this was aetiologicaly related to the peri-arthritis is doubtful. In an elderly hemiplegic the disease developed in the shoulder of the paralysed side three months after the cerebral haemorrhage. In eight persons stiff, painful shoulder developed without any apparent cause.

Associated Diseases.—Two patients had diabetes mellitus, and two were subject to gout.

Symptoms and Signs

The outstanding symptom is pain, which with one exception was strictly unilateral. It is often spread diffusely over the shoulder-joint, but sometimes it is referred to the anterior aspect over the long head of the biceps or lower down over the belly of this muscle. In long-standing cases the pain may be much more widespread, even extending to the forearm and fingers. It is of the nature of a dull ache if the arm is kept still, but passes into acute agony in attempted shoulder movement. Often it is of neuritic type, shooting down the limb and partaking of a boring and burning character, and made worse by getting hot in bed at night. Limitation of movement as a symptom is of secondary importance owing to the mobility of the scapula masking the actual fixation. In women, however, it is often a source of complaint that