We know, further, that cancer cells may remain latent, presumably encapsulated by a fibrous tissue reaction. This is the only possible explanation of the late recurrences sometimes seen. I would instance a patient whom I saw recently who had a radical removal of the breast done for carcinoma, and who came up twenty years later for a recurrence in the lumbar spine. In this particular case it is certain that the deposit was there at the time of the operation. I am suggesting that operation, by cutting down the amount of cancer present, allows it to be successfully dealt with and shut off. Then at a later stage some other factor comes into play, possibly trauma, possibly a lowering of the general health of the patient, and the cancer cells resume their interrupted activity.

I do not wish to imply that partial operations should be undertaken, or that radical removals of a primary source should be done in the presence of gross secondary deposits. I am only placing this case on record to show that a patient who might have been regarded as hopelessly inoperable is still alive and well after an interval of ten years.

Summary

A case of carcinoma of the descending colon associated with a nodule in the liver is reported in which the patient is alive and well ten years after a partial colectomy.

REFERENCES

- Gould, A. Pearce (1910). *Lancet*, 2, 1665.
Teacher, J. H. (1908). *J. Path. Bact.*, 12, 131, 487.

THE PLANTAR WART

VERRUCA PLANTARIS: VERRUCA PEDIS: PLANTAR PAPILLOMA

BY

R. M. B. MacKENNA, M.A., M.D., M.R.C.P.

Honorary Dermatologist, Royal Southern Hospital, Liverpool

(WITH SPECIAL PLATE)

At the present time there is a widespread epidemic of plantar warts among school children. So far as my experience goes the epidemic is most severe among adolescent girls, although I have seen several boys whose ages ranged from 8 to 17 years suffering from the complaint. In South-West Lancashire the epidemic scarcely affects the so-called hospital classes, and the majority of cases are in children who attend boarding-schools; it would seem to be unduly optimistic to believe that all classes of the community will not be soon infected. Until recent years this type of wart has not been prevalent, and it is difficult to discover why the present epidemic has developed.

Plantar warts (or "verrucae," as they are called by many patients and some chiropodists) are a minor malady, but when they have reached an average size they considerably curtail the activities of the person affected, owing to the pain caused by pressure on the warts whenever he attempts to walk or run. The warts develop insidiously. The first sign is the formation of a small semi-translucent macule, 1 to 2 mm. in diameter. There are no signs of inflammation. The macule is slightly painful on pressure, but is usually disregarded by the patient, who, if he considers the matter at all, believes that he has a tiny piece of grit embedded in the sole of his foot. The lesion, as a rule, develops on the heel or the ball of the foot, but may appear on any part of the plantar surface,

including the toes. The macule increases slowly in size until it reaches a diameter of approximately 0.5 cm., when the wart may be regarded as being fully formed. On examination one finds a roughened, hyperkeratotic type of lesion, greyish yellow in colour, and its papillomatous nature is usually not apparent until the wart is pared with a knife. Satellite warts develop, and it is not uncommon to find as many as ten warts on one foot. The subjective symptoms vary with the site of the warts and the stoicism of the patient. One can generally obtain a history of throbbing pain, which is most severe when the patient gets out of bed in the morning or when he puts on his boots and starts to walk. The pain usually diminishes after he has been standing for some little time. He may complain that the warts throb when he has a hot bath.

Plantar warts have to be differentiated from callosities due to pressure; but medical men are so accustomed to recognizing callosities due, for example, to dropping of the anterior arch of the foot or to faulty shoes that the differential diagnosis is not often a matter of difficulty. In a doubtful case one should remember that whereas little or no pain is experienced when a callosity is pressed, the patient always complains of severe pain if a plantar wart is treated similarly; further, if the lesion is pared down the papillomatous nature of the wart becomes apparent.

Aetiology and Pathology

There is little doubt that plantar warts are caused by a filter-passing virus. It is probable that individual susceptibility to this virus varies considerably, but the virus is virulent and appears to be able to obtain a secure foothold on skin which to the naked eye reveals no sign of abrasion. Whilst the virus probably cannot develop or multiply except on a medium of living cells, it is reasonable to presume that it can remain alive for some minutes, if not for a few hours, on areas where an affected foot has pressed. If the superficial layers of a wart are shaved off the closely packed papillae are revealed. The papillae are in most cases yellowish white in colour, but may be stippled with red or brownish stains where haemorrhage has occurred. They are encircled by a tough horny collar. The lesion is broader at its base than at the opening on the surface of the skin.

The histology has been described by MacLeod (1933) in the following terms: "Microscopical sections show hyperkeratosis at the periphery, and marked acanthosis and elongation of the interpapillary processes in the centre. Towards the centre of the processes the prickle cells tend to be vacuolated." Inclusion bodies may be found in these vacuolated prickle cells.

Treatment

Treatment must be divided into two categories, preventive and active. In my opinion preventive treatment is of the utmost importance at the present time, and I would urge that all doctors who are in charge of schools should institute regular inspection of their pupils' feet, at least until the present epidemic has passed. School children suffering from plantar warts should be segregated so far as dormitory accommodation is concerned. They should have their own bathrooms and changing-rooms, and should not be permitted to use the school swimming-bath. Special care should be taken that their boot lockers are far removed from their colleagues', whilst the bootblack should have instructions always to clean the patients' boots in a batch after he has finished with those belonging to the rest of the school.

In private houses similar precautions should be taken. The infected person must never tread with bare feet where an unaffected person will step. On leaving his bath he should step on to his own bath mat, which should be removed from the bathroom as soon as he leaves. No one else should use his bath towel, and probably it is safer to swill down the bath with a weak antiseptic after he has used it. The evidence is not conclusive, but it is probable that the infection is more usually contracted in bathrooms than elsewhere. So little is known about the dissemination of the wart virus that it is better to be overcautious rather than risk dissemination of the virus by lack of caution.

In the earliest stages the warts can in most cases be destroyed by the use of a wart paint—for example:

R. Zinci chloridi 3 ss
Collodium acid. salicyl. (1 in 10) ad 3 i
Misce.

A spicule of wood should be dipped into this paint and then bored gently into the wart. The treatment should be carried out twice daily, and if the patient is placid the results are usually satisfactory. Many persons complain bitterly that they cannot stand the pain, and other measures have to be employed.

Most dermatologists prefer to use radium or x rays, and these physico-therapeutic agents are the least painful methods of treatment and the most certain in their results. In my experience, using x rays filtered through 0.5 mm. aluminium, a plantar wart requires an average of 900 r units of x rays before it succumbs. While receiving treatment it is advisable for the patient to wear white cotton socks underneath his usual socks throughout the day. These socks should be boiled every night. The feet may be powdered with an antiseptic dusting powder.

If physico-therapeutic measures are not available excision may be attempted. In my hands local anaesthesia has not been very successful: I have found it to be almost impossible to infiltrate the part with novocain without causing much pain. I prefer to use a general anaesthetic (nitrous oxide or gas-and-oxygen). The warts should be scraped out with a sharp curette. It is surprising how deeply the roots are situated. I have found it best to remove the horny collar which surrounds each wart, as recurrence is frequent if this is left. Bleeding should be controlled and the area disinfected, so far as is possible, by the application of the silver nitrate pencil. Dry dressings are usually adequate, and healing is complete in about fourteen days. Diathermy or refrigeration with carbon dioxide snow are alternative measures. Goldsmith (1936) recommends that the "snow" should be applied for three to five minutes: a large bleb will develop, allowing the wart to be painlessly excised about three days later. Good results have been claimed for treatment by cataphoresis with zinc sulphate.

Chiropodists often treat these cases successfully. They generally aim at destruction of the wart by escharotics or keratolytics. Runting (1935) states that their object is "to destroy the growth so thoroughly that such a reaction may be created in the healthy tissue as to bring about a line of cleavage between the destroyed growth and the normal healthy tissue. The culmination of this process is usually accompanied by effusion of serum or sero-pus. . . . The whole art of the successful destruction of verrucae depends upon inducing this condition effectually without causing more disturbance to the normal tissue than is actually essential to the cure." "The majority of chiropodists bring about the desired result with acids, such as nitric, acetic, monochloracetic,

trichloracetic, salicylic, chromic, pyrogalllic, or acid nitrate of mercury; or with such powerful alkalis as potassium hydroxide or sodium hydroxide."

In the majority of cases the surrounding tissue is protected by an application of vaseline, and the acid or alkali is applied with an orange stick and "worked into the growth." A protective dressing is employed and a few days later the treatment is repeated, the tissue which has been destroyed by the previous treatment being removed before fresh acid or alkali is applied. By proceeding slowly and giving regular treatments the warts are destroyed with a minimum of pain. The use of pyrogalllic acid is effective and is not associated with much pain unless the application is left *in situ* too long. The acid is applied as an ointment (30 to 50 per cent. in ung. resinae) once or twice weekly, the usual occlusive precautions being observed.

Summary

Attention is drawn to an epidemic of plantar warts among school children.

A short description of the signs, symptoms, aetiology, pathology, and treatment of the warts is given.

[The photograph in the Special Plate is reproduced from *Diseases of the Skin* (fourth edition), published by Messrs. Baillière, Tindall and Cox.]

REFERENCES

- Goldsmith, W. N. (1936). *Recent Advances in Dermatology*, p. 387, London.
MacLeod, J. M. H. (1933). *Diseases of the Skin*, p. 1096, London.
Runting, E. G. V. (1935). *Practical Chiropody*, p. 35, London.

INTRATHECAL ALCOHOL INJECTION FOR RELIEF OF PAIN: A REVIEW

BY

CONSTANCE OTTLEY, F.R.C.S.

Surgeon, New Sussex Hospital for Women; Surgeon,
Marie Curie Hospital

Dogliotti in 1931 first suggested the use of subarachnoid injections of absolute alcohol as a simple and comparatively safe means of alleviating severe chronic pain of which the cause is not amenable to treatment. He pointed out that the various surgical measures designed to relieve pain (posterior rhizotomy, chordotomy, sympathectomy) are operations to which many patients suffering from such conditions as advanced malignant disease cannot justifiably be submitted; nor are they always successful. Paravertebral nerve-blocking is an alternative, but it is difficult to carry out and is uncertain in its effects.

Dogliotti's aim in devising this method was to bring about a partial destruction of the appropriate posterior nerve roots close to their entry into the spinal cord. Accepting the assumption that pain is proportional in severity to the numerical strength of the sensory impulses reaching the centres concerned, he argued that this "decimation" of the afferent nerve fibres should reduce the pain-provoking stimulus below threshold value. From animal experiments he satisfied himself that it is possible to limit the action of the alcohol with sufficient precision to the posterior roots; only when the doses used were excessive did symptoms of motor paralysis follow the injections.

Principles of Treatment

The principle underlying the method is that absolute alcohol possesses a specific gravity considerably lower

DIGBY CHAMBERLAIN: SPONTANEOUS DISAPPEARANCE OF CARCINOMA

FIG. I.—Showing the proliferative character of the growth. ($\times 45$.)

FIG. II.—Showing how the growth is invading the muscle. ($\times 45$.)

R. M. B. MacKENNA: THE PLANTAR WART

Plantar warts.

PHYLLIS F. L. DAPLYN: BLOCK ANAESTHESIA FOR
INSERTION OF RADIUM TO EPITHELIOMA OF LIP

Condition of lower lip at operation.