thoroughly cleared, pus being found. Convalescence was uneventful, and she was discharged on May 15 with the wound and the tympanum healed and hearing nearly restored to normal. Complete recovery ensued, and the patient had no complaints to make when last seen on September 25.

Summary

1. A simple method is described in which Linzemeier's tube is employed.

2. The figure 10 is selected as the usual maximum reading attained in health.

3. A chart method is employed. By taking weekly readings throughout the year it is shown (a) that the B.S.R. varies in health within moderately wide limits; (b) that each individual has an optimum reading; (c) that moderate deviations from this can usually be attributed to minor ailments.

4. During pregnancy the B.S.R. is undisturbed until about the fourth month, when a steady rise takes place till term.

5. General principles are deduced from the behaviour of the B.S.R. in local infection. The rise of the B.S.R. to an appreciable degree takes about three days. Its decline is comparatively slow, and is (a) continuous and satisfactory; (b) interrupted, due to complications; or (c) incomplete and unsatisfactory. Cases should be followed up until the B.S.R. returns to normal.

I owe my thanks to Dr. T. C. Hunt, who first aroused my interest in the subject, and to Professor J. R. Fraser for suggestions and criticism.

Towards the expense of this research a grant was made by the British Medical Association.

ASCORBIC ACID IN BRONCHIAL ASTHMA

REPORT OF A THERAPEUTIC TRIAL ON TWENTY-FIVE CASES

BY

H. B. HUNT, M.D.

There is a certain amount of evidence that ascorbic acid may play a part in allergic conditions in general and asthma in particular which, although suggestive, is difficult to correlate exactly, and is perhaps best classified in three groups as follows:

Theoretical Considerations

1. Hypersensitivity to foreign proteins results after adrenalectomy in dogs (Simpson, 1937). In asthma a condition of protein hypersensitivity also exists, and hypoplasia of the adrenal glands has been noted in asthmatics coming to necropsy (Adam, 1931; Waldbott, 1933-4). An eosinophilia is not uncommon in either asthma or Addison's disease. At least one case has been recorded of the association of these two conditions (Wilkinson, 1937).

2. Adrenaline, the most potent anti-allergic remedy yet known, is a secretion of the adrenal medulla. The fact that cortical extracts have been employed in asthma without any marked success (Cohen and Rudolph, 1934-5; Wilmer and Miller, 1936) tends to show that if the suprarenal glands are affected in this condition, then the medulla rather than the cortex is involved.

3. Ascorbic acid is normally present in the adrenals. Indeed, it was from this source that Szent-Györgyi originally isolated vitamin C.

From these disjointed observations it appears probable that (1) asthma or allergy may be associated with adrenal hypofunction; (2) the medulla is chiefly at fault; (3) some especial relation exists between ascorbic acid and the adrenals. Dr. L. J. Harris has suggested (in a personal communication) that as, in the light of recent observations, one of the functions of ascorbic acid appears to be to prevent the oxidation of adrenaline, it may be possible, by giving suitable doses, to conserve sufficient adrenaline to prevent the occurrence of asthmatic symptoms.

Vitamin C in Anaphylaxis and Allergy

Claims have been made that ascorbic acid will protect guinea-pigs against anaphylaxis (Hochwald, 1935; Salomonica, 1936), and in human beings will diminish the cutaneous reactions to tuberculin, in positive reactors when given intravenously (Heise *et al.*, 1937). Hochwald (1936) has employed ascorbic acid in cases of bronchial asthma, and found it of some value in preventing symptoms when given regularly, or aborting symptoms when injected intravenously in massive doses (500 to 1,500 mg.). Epstein (1936), while unable to confirm the work of Hochwald when he used ascorbic acid in either asthma or hay fever, did find that it was distinctly valuable in the treatment of asthma if employed in conjunction with injections of gold salts.

Investigations

In view of what has been said above there is reason to expect that ascorbic acid may prove useful in the treatment of bronchial asthma. It is not suggested that asthma is a deficiency disease or even that the majority of asthmatics suffer from a subclinical C avitaminosis. Ascorbic acid was given to exert any anti-allergic properties which it may possess rather than to correct a specific deficiency. As all of the cases were out-patients it was not practicable to employ the usual laboratory procedure of estimating any possible avitaminosis. However, among the whole of the patients attending the asthma clinic of the Birmingham General Hospital during the last year I detected only one who showed any clinical signs of vitamin C deficiency. In this case the symptoms—bleeding of the gums and lassitude—were very mild, and cleared up after suitable treatment with ascorbic acid.

Clinical Material.—Twenty-five patients were investigated—sixteen females and nine males; five of them were children. Other allergic symptoms were present in seventeen cases, and chronic bronchitis in nine. Thus, although small, the series was fairly representative.

Dosage and Method of Administration

The ascorbic acid was dispensed in tablets of 50 mg. (the synthetic preparation of Roche Products Ltd., redoxon, was employed throughout), two of these being given for each day, one to be taken in the morning and one at night. Each patient received one or two weeks' supply of tablets when attending the clinic. The course was designed to last for eight consecutive weeks, but as attendances were not as regular as could be desired all the patients did not complete eight weeks' treatment, and, of those who did, not every one covered the course in eight consecutive weeks. During this treatment no medicine or injections such as vaccines were given, apart from antispasmodics—for example, ephedrine or "felsol" powders—or medicine for a troublesome cough. Five patients (quite apart from the group of twenty-five mentioned above) seen when symptoms were present were

given massive doses of ascorbic acid by injection. One of them received 500 mg. intramuscularly, another 800 mg. intravenously, and the remaining three 500 mg. by the same route.

Results

In reviewing the twenty-five cases after several weeks' treatment by oral medication with 100 mg. of ascorbic acid daily no evidence could be found of any marked improvement in the amount of wheezing, the incidence of attacks, or the general condition. None of the five patients who received injections of ascorbic acid in doses of 500 mg. or over experienced any relief in their symptoms after a period of twenty to thirty minutes, and the amount of adrenaline subsequently required to relieve the spasm was in no way diminished. Thus it was impossible to confirm the statement of Hochwald (1936) that asthmatic symptoms could be relieved by large intravenous injections of ascorbic acid. It may be mentioned that in these five cases symptoms were severe in two instances and slighter in three.

Conclusions

The investigations described above have not shown ascorbic acid to be of any value in the treatment of bronchial asthma when given in comparatively large doses either by injection or by mouth.

I wish to thank Professor W. H. Wynn for the facilities he offered me in the investigation; and also Roche Products Ltd. for the supplies of redoxon which they kindly placed at my disposal.

REFERENCES

- Adam, J. (1931). *Glas. med. J.*, **116**, 81.
Cohen, M. B., and Rudolph, J. A. (1934-5). *J. Allergy*, **6**, 279.
Epstein, A. (1936). *Schweiz. med. Wschr.*, **66**, 1087.
Heise, F. H., Martin, G. J., and Schwartz, S. (1937). *Brit. J. Tuberc.*, **31**, 23.
Hochwald, A. (1935). *Zbl. inn. Med.*, **56**, 769.
— (1936). *Klin. Wschr.*, **15**, 894.
Salomonica, B. (1936). *J. Immunol.*, **31**, 209.
Simpson, S. L. (1937). *British Medical Journal*, **1**, 229.
Waldbott, G. L. (1933-4). *J. Allergy*, **5**, 323.
Wilkinson, J. F. (1937). *Lancet*, **2**, 61.
Wilmer, H. B., and Miller, M. M. (1936). *J. Allergy*, **8**, 77.

R. Benda (*Prag. med. Z.*, January 25, 1938, p. 67) does not agree that a retroverted uterus discovered during a routine examination and producing no symptoms should be left untreated. He recommends replacement and insertion of a pessary in all cases. Operative treatment is indicated if after one year's conservative treatment the uterus still becomes displaced after removal of the pessary. Replacement of a gravid retroflexed uterus should only be undertaken in hospital. Although spontaneous replacement may take place, and although it may be aided by lying in the prone position, it must not be left to chance. When retroversion is due to pelvic peritonitis conservative treatment of the pelvic inflammation is indicated. The insertion of a ring pessary in the treatment of uterine prolapse is only of value when the levatores ani are still intact. It is valueless when the neck of the cervix is elongated and when cystocele or rectocele is present. Of the various operative procedures Benda advises Wertheim-Schauta's method of interposition of the uterus between the vagina and bladder, or Halban's fixation of the uterus to the bladder. He attacks anterior and posterior colporrhaphy as an operation attended by an appalling proportion of relapses. Measures for the prevention of uterine prolapse include careful forceps deliveries, more frequent use of episiotomies, and exercises in the puerperium. Benda advocates that women should be encouraged to get up five days after a normal confinement, but they should not be allowed to do their ordinary work for some time afterwards.

THE FEMALE CLIMACTERIC AND THE MENOPAUSE

BY

H. R. DONALD, M.A., B.M., M.R.C.P.

Honorary Physician, Manchester Northern Hospital, Stretford
Memorial Hospital, and Royal Eye Hospital;
Visiting Physician, Crumpsall Hospital

The recent advances made in the physiology of the sex hormones have tended to stimulate rather than to satisfy our curiosity. The shafts of light that have been shed on a few isolated points have by contrast served merely to accentuate the darkness of surrounding ignorance. This is particularly true of the existing knowledge of the female climacteric. The common employment of the terms "menopause" and "climacteric" as though they were synonymous advertises the looseness of present conceptions, for although both these conditions possess certain similar features there is no possible justification for regarding them as identical. Some of this confusion may be due to the tendency to make deductions from the similarity of the symptoms of the artificial and of the physiological menopause. Such argument by analogy is always risky (as in the case of the experimental diabetes of pancreatectomy and its supposed clinical counterpart), and in view of the unknown causation of the physiological menopause any inferences should be drawn with extreme caution. Since a clear differentiation between the menopause and the climacteric is becoming increasingly important, the two subjects are here reviewed from both the theoretical and the clinical standpoint.

The Menopause

The use of the term "menopause" should be restricted to its literal meaning of the cessation of the menses, which should also imply the termination of the menstrual life. The cause of this permanent amenorrhoea is not known, but it is commonly accompanied by two principal endocrine changes. First, there is a relative diminution in circulating oestrogens, which are almost or completely absent from the blood stream three years after the cessation of the menses in probably at least 50 per cent. of women. It is doubtful, however, if the menopause is due directly or solely to a decline in the ovarian secretion, since full endometrial hyperplasia, which may be regarded as a specific oestrogenic function, may persist for many years after the establishment of amenorrhoea. Its occurrence has been variously reported in forty out of 804 and fifteen out of 130 post-menopausal subjects (Novak and Yui, 1936). The site of formation of these oestrogens is not known, but some at least may originate in the ovary. In the second place, there is an increased content of prolactin in both the pituitary gland and the urine, but the significance of even this finding is not appreciated. The increased content of prolactin in the pituitary is probably more characteristic of a resting than an actively secreting state, and it may be of interest to quote the work of Engelhart and Häusler (1936-7), who reactivated the senile ovaries of rats by the implantation of pituitary tissue.

The Climacteric

The term "climacteric" is derived from the Greek, meaning "a rung of a ladder," and originally was used to describe certain critical epochs, in the life of both men and women, at which the body was believed to undergo radical changes. These epochs were supposed to recur at intervals of nine years, the two important ones being at adolescence and at the sixty-third year, the