improvement has been effected? I believe that it can. There has been no change in the methods of reference of patients to the Institute over the whole seven years under review, and cases referred as carcinoma of the antrum remain few. The average number of missed antrum cases per year is ten. During these seven years the Institute staff failed to recognize fifteen out of the seventy cases until recurrence took place. Of these fifteen only three have occurred during the last two years—that is, since the high incidence of antral involvement in palate malignancy was realized. Each of these three cases was missed because again a negative x-ray report was accepted as being conclusive.

Conclusions

It may be inferred from the report of the National Radium Commission that the frequent involvement of the maxillary antrum in certain palatal and upper alveolar malignant lesions is not generally appreciated.

Investigation in the Manchester Radium Institute of 167 cases of malignant disease of the palate showed that 94 per cent. of the lesions of the middle third of the palate, spreading on to the alveolus and into the alveololabial sulcus, involved the maxillary antrum. In the Institute this is known as the typical antral site. Lesions in the other usual palate site—that is, the posterior part of the hard palate, spreading on to the soft palate and fauces—did so in only 1.4 per cent. of the cases.

Typical antral-site lesions should be regarded as involving the antrum unless it is proved that they do not. If definite proof is not forthcoming they should be assumed to do so, and be treated accordingly. X-ray examination alone is conclusive only when positive. A negative report should not be accepted as being conclusive. Thirteen cases out of fifteen with antral involvement, missed until recurrence followed treatment, escaped notice through the acceptance of negative x-ray reports.

The needle test described is useful, but should only be accepted when positive.

Histological examination does not help to clear up the diagnosis, because of the high proportion of squamous carcinomata in both the simple palate lesions and those involving the antrum, taken in conjunction with the fact that all malignant tumours of the palate are not epitheliomata. As treatment must include both mouth and antrum, anatomical differentiation of site of origin is not considered to be of great clinical importance.

The cases seen in the Institute suggest that intra-oral ulceration is by far the commonest evidence of antral neoplasm. This finding is contrary to accepted teaching, but is not due to the fact that usually it is the more advanced cases which are referred for radiation therapy, as cheek-bulging is rare even in the obviously late cases.

It is shown that recognition of the palate-antrum relationship has been followed by a great reduction in the number of cases entirely missed until local palate treatment was followed by recurrence.

From April 1 the importation into this country of all meat, including bacon and ham, and also meat products such as canned meats, sausages, and sausage casings, will be permitted only if they are accompanied by an official certificate recognized by the Minister of Health as evidence of satisfactory inspection and hygienic preparation. The importation of certain defined classes of meat (such as scrap meat) continues to be prohibited.

TINEA OF THE FOOT

BY

P. K. FRASER, M.B., Ch.B.

Surgeon Lieutenant, R.N.

Since the discovery of pedal tinea by Whitfield in 1908 the condition has become increasingly prevalent (C. J. White, 1927a), and according to American authorities presents to-day one of the major problems of dermatological practice. The many statistical surveys that have been made in America by Hulsey and Jordan (1925), Legge, Bonar, and Templeton (1929), Andrews and Birkman (1931), and Gilman (1935), to mention only a few, appear to confirm this, and it seems strange that no comparable figures have been produced in this country. It may be as Gray (1934) said, that "the profession as a whole is still barely aware that the problem exists," and should the disease achieve the lay popularity it has achieved elsewhere there is every danger of a too free diagnosis on inadequate evidence.

Difficulties in Diagnosis

The frequent and early spontaneous disappearance of fungal mycelia from the infected area is one of the main difficulties, and in late cases there may be nothing to distinguish the condition from dermatitis from other causes (Whitfield, 1908, 1911). The reason for this disappearance is hard to understand in view of the viability of the pathogenic fungi (Farley, 1921; Weidman, 1927a; Bruhns, Alexander, and Kadisch, 1929), but the secondary bacterial contamination which sometimes occurs may in some way exert a lethal influence on the moulds. In this connexion the observations of Weidman (1926) on a case of pedal ringworm sown with yeasts are of importance. Thus it would appear that many cases of tinea, when seen, will not show the causal agent, although such authorities as J. H. Mitchell (1922) and C. J. White (1919, 1927b) countenance a diagnosis on clinical grounds alone in the chronic keratotic pedal type. This may be admissible under certain circumstances, but the demonstration of the specific factor still remains a *sine qua non* in diagnosis, and the following statement by Ramsbottom (1931) may be taken as an expression of general opinion:

"There is, however, a tendency on the part of some investigators to assume a fungal origin of a disease on insufficient evidence. It is not enough to postulate the parasitic nature of a disease and then, bacteria, protozoa, and other parasites being eliminated, to describe it as mycotic; the evidence must be as clear as that for any other parasitic group."

In addition it must be remembered that all cases of interdigital scaling and maceration are not mycotic, although von Graffenried (1918) and Catanei (1932) have shown that such lesions form a suitable nidus. Dirt seems to bear no relation to infection, as the highest incidence is to be found among the wealthy and more leisured classes (Beintema, 1931). Therefore the practitioner will often be confronted with strong clinical evidence of fungus infection without microscopical or cultural confirmation, and a specific dermal test would be of great help in accurate diagnosis.

Previous Investigations with Mycotic Extracts

The use of mycotic extracts in the diagnosis of superficial tinea infers an altered state of reactivity in the body. That this state exists in certain bacterial diseases is a well-established fact, and for many years mycologists have endeavoured to find a corollary in ringworm infections.

The position with regard to deep ringworm has practically been cleared, but the occurrence of general changes in the superficial type has caused much controversy. The tendency towards spontaneous cure in deep tinea (Sabouraud; Greenbaum, 1924) has been explained by the development of a general immunity, and the rarity of scalp infection in adults has been cited as an example of acquired herd immunity. Supporting the development of immunity Greenbaum (1924) and Scholz (1934) showed that by inoculating guinea-pigs cutaneously with ringworm they rendered the animals immune to further inoculation, and Sabouraud found that he could not use the same animal twice in testing for the pathogenicity of moulds. Greenbaum also showed that by inoculating the skin of a human with *Trichophyton gypseum* spontaneous healing occurred in four weeks, and that the site of the lesion could not be reinfected as long as three months afterwards. Further, he claimed that the immunity thus produced was group-specific against the other ringworms, and said that "so sharply defined is this local immunity that reinoculation of a previously healed site only shows a take at the exact edge of the previous lesion." Disproving this cutaneous immunization as applied to superficial infection, Weidman (1927b) was able to infect the first interdigital cleft of his foot on two occasions with *Trichophyton cruris* after a lapse of two years, although the interval appears to have been unduly long. It is also not uncommon in tropical practice to see more than one attack of tinea cruris in the same patient, despite the fact that Gray (1934) thinks this is rather unusual.

The occasional development of distal phytid eruptions in superficial infections has been established by Jadassohn and Peck (1929), Sulzberger and Kerr (1930), Peck (1930), and Ayres and Anderson (1934), while a fungus has been recovered from the circulating blood in some cases (Sulzberger, 1928; Peck, 1930; Strickler *et al.*, 1932). It is of interest to note the inhibitory effect on ringworm cultures of serum from cases with phytid eruptions (Per and Braude, 1928; Ayres and Anderson, 1934). Ravaut and Rabeau (1932) even claim general changes in superficial yeast infections, and they have used an extract of yeast cultures, "levurin," as a diagnostic test.

From the foregoing evidence it is logical to conclude that superficial ringworm can sometimes cause local immunity, and there is reason to believe that the products of the causal fungus are not limited to the site of infection.

The diagnostic skin tests have given very variable results in the hands of different investigators. Such an eminent mycologist as Castellani (1934) places little reliance on them, although he publishes full details for the preparation of the extract; while Tarantelli (1926) and Garzella (1926) insist that to produce a positive result the lesion should be deep-seated. Amberg (1910), using linear scarification, had thirteen positives from 131 casual patients; four of the thirteen had no previous history of ringworm. Repeating this investigation, Rosen, Peck, and Sobel (1931) intradermally tested 102 dermatological patients—seventy-two of them with some pedal skin lesion—and found a positive reaction in sixty-seven. There were nine proved cases of tinea pedis in this series, and eight of them reacted strongly. Muskatblit and Director (1933) encouragingly report 72 per cent. positive results in 300 cases of superficial tinea, while Ruete and Scholz (1934) claim 100 per cent. success in proved cases. In addition, the last-named observers contend that the extract used will only react against the fungus from which it is prepared, although this is disproved by Sulzberger and Kerr

(1930). These group reactions are accepted generally, and are to be expected in view of the close biological relationship of the pathogenic moulds. Lomholt (1934) and Muende (1934) regard dermal tests as being of especial value in the differentiation of mycotic from dysidrotic eczema—a matter of importance in this country on account of the insignificance of the mycotic factor in the latter type of lesion (McLachlan and Brown, 1934).

The Present Investigation

The following investigation was carried out in ships of war during the last three summers. In all, 111 cases were examined, twenty-two of them with superficial pedal tinea; the remaining eighty-nine were casual patients used as controls. In every case a careful examination of the interdigital crural and axillary folds was made, and any skin surface showing maceration or desquamation was examined microscopically for fungus, while the previous dermatological history was assessed as to the possibility of a fungus infection. Polyvalent trichophytin (Hoescht) was used throughout in a dilution of 1 in 50, and the diluent was employed as a control. Dilutions of 1 in 250 and 1 in 800 were also used at the beginning of the investigation, but were soon dispensed with as unnecessary. The solutions were renewed at monthly intervals to ensure against deterioration. After a trial of the linear scar and patch methods the intradermal route was chosen as being the most reliable. The trichophytin was injected with a fine needle into the flexor surface of the right arm and the control was injected in a similar site in the other arm, care being taken that the syringes and needles used were dry and free from traces of chemicals.

Four types of reaction were met with: (1) immediate and transient urticarial reaction; (2) inflammatory reaction; (3) inflammatory reaction with vesicle formation; (4) late eczematoid reaction. Types 1, 2, and 4 correspond to those described by Sulzberger and Wise (1932), and Type 3 has been added on account of its frequent occurrence in the strongly positive group. Type 1 was occasionally seen in the negative group, and, according to Bray, is due to tissue damage at the site of injection. The results were read at intervals of one, two, four, and seven days, and were classified according to the following table:

Classification	Day of First Appearance	Maximum Diameter of Wheal	Vesicles	Desquamation
Strong	1st-2nd	20-40 mm.	+	+ +
Medium	1st-3rd	10-20 mm.		+
Weak	2nd-3rd	10 mm.		
Negative				

The trichophytin when fresh appears to be remarkably resistant to heat, as five positive and five negative cases retested after autoclaving the extract gave the same results. A summary of the investigation is set out in tabular form below:

Type of Case	No. of Cases	Strong Reaction	Medium Reaction	Weak Reaction	Negative Reaction
(A) Microscopically proved tinea	22	14	2	2	4
(B) Control with previous history of tinea ..	26	3	8	10	5
(C) Control with negative history	63	2	15	11	35

Owing to the difference in the numbers of cases in the three groups it is proposed to scrutinize the table as it stands and not in terms of percentages. The four negatives in Group A are disappointing, but all of those cases showed "mosaic" fungi microscopically. According to Weidman (1936) such "mosaics" are either a stage in the development of fungus or are due to some fungoid action on the superficial layers of the dermis, while Whitfield (1935) regards them as artefacts due to interaction between the epidermal scales and the caustic potash used in preparation of the microscopical specimen. Repeated attempts to culture scales showing such "mosaics" were made under ideal conditions in Ramsbottom's mycological laboratory without success. Their exact nature, therefore, must be a matter for doubt, but it is of interest to note their frequent association with the accepted type of fungus. The results in Group B are gratifying, but the unavoidable personal error in assessing the history makes it difficult to draw a definite conclusion; and this becomes especially noticeable in Group C. Many patients may have had a patch of superficial tinea not diagnosed as such, or an interdigital type may have escaped notice completely. The frequency of such previous infections has been regarded as negating the value of a positive reaction (Wise and Wolf, 1936), and this would seem to increase considerably the value of a negative reaction.

Conclusions

From this small series and from observation of the skin tests in other hands, it is permissible to state that a strong reaction in a clinically suggestive case with a negative previous history lends a bias towards a diagnosis of tinea, while a negative reaction in a similar case precludes a positive diagnosis completely.

An interesting outcome of the investigation was provided by two negative reactors in Group C. Some time after testing they developed microscopically proved tinea pedis—one with an associated tinea cruris—and when retested three weeks and twelve months after infection gave a strongly positive result.

Summary

Attention is called to the increasing frequency of pedal tinea in America and to the large number of statistical investigations published by American observers.

The lack of comparable figures in this country is noted, and appears to be due to non-appreciation of the condition among the great mass of the profession.

The difficulties and pitfalls of diagnosis are described and a conservative opinion by Ramsbottom is quoted.

The case for diagnostic dermal tests with fungus extracts is set forth and the results of some previous investigations are given.

An investigation is described and the results given in tabular form.

The limitations of a positive result are noted, and the value of a negative result in the differential diagnosis of pedal skin lesions is stressed.

Grateful acknowledgments are due to Surgeon Captain C. M. R. Thatcher and Surgeon Commander R. A. W. Ford for permission to use their cases; also to Miss F. L. Stephens for her cultural work and to Mr. T. J. Shields for his assistance with the references.

REFERENCES

- Amberg, S. (1910). *J. exp. Med.*, **12**, 435.
 Andrews, G. C., and Birkman, F. W. (1931). *N. Y. St. J. Med.*, **31**, 1029.
 Ayres, S., and Anderson, N. P. (1934). *Arch. Derm. Syph.*, Chicago, **29**, 543.
 Beintema, K. (1931). *Nederl. Tijdschr. Geneesk.*, **75**, 4302.
 Bruhns, C., Alexander, A., and Kadisch, E. (1929). *Med. Klinik*, **25**, 1769.
 Castellani, A. (1934). *J. trop. Med. Hyg.*, **37**, 365.
 Catanei, A. (1932). *Arch. Inst. Pasteur Alg.*, **10**, 195.
 Farley, D. L. (1921). *Arch. Derm. Syph.*, Chicago, **3**, 759.
 Garzella, N. (1926). *G. Ital. Derm. Sif.*, **67**, 280.
 Gilman, R. L. (1935). *J. Amer. med. Ass.*, **100**, 715.
 Graffenried, C. von (1918). *Derm. Wschr.*, **66**, 361.
 Gray, A. M. H. (1934). *British Medical Journal*, **2**, 541.
 Greenbaum, S. S. (1924). *Arch. Derm. Syph.*, Chicago, **10**, 279.
 Hulsey, S. H., and Jordan, F. M. (1925). *Amer. J. med. Sci.*, **169**, 267.
 Jadassohn, W., and Peck, S. M. (1929). *Arch. Derm. Syph.*, Berlin, **158**, 16.
 Legge, R. T., Bonar, L., and Templeton H. J. (1929). *J. Amer. med. Ass.*, **93**, 170.
 Lomholt, S. (1934). *Ugeskr. Laeg.*, **96**, 655.
 McLachlan, A. D., and Brown, W. H. (1934). *Brit. J. Derm. Syph.*, **46**, 457.
 Mitchell, J. H. (1922). *Arch. Derm. Syph.*, Chicago, **5**, 174.
 Muende, I. (1934). *Brit. J. Derm. Syph.*, **46**, 488.
 Muskatblit, E., and Director, W. (1933). *Arch. Derm. Syph.*, Chicago, **27**, 739.
 Peck, S. M. (1930). *Ibid.*, **22**, 40.
 Per, M. L., and Braude, R. (1928). *Acta derm. venerol.*, **9**, 1.
 Ramsbottom, J. (1931). Medical Research Council, *System of Bacteriology in Relation to Medicine*, **8**, 11, London.
 Ravaut, P., and Rabeau, H. (1932). *Presse méd.*, **40**, 1925.
 Rosen, I., Peck, S. M., and Sobel, N. (1931). *Arch. Derm. Syph.*, Chicago, **23**, 1041.
 Ruete, A. E., and Scholz, U. (1934). *Derm. Z.*, **68**, 241.
 Scholz, V. (1934). *Derm. Wschr.*, **99**, 1342.
 Strickler, A., et al. (1932). *Arch. Derm. Syph.*, Chicago, **25**, 1028.
 Sulzberger, M. B. (1928). *Ibid.*, **18**, 891.
 — and Kerr, P. S. (1930). *J. Allergy*, **2**, 11.
 — and Wise, F. (1932). *J. Amer. med. Ass.*, **99**, 1759.
 Tarantelli, E. (1926). *G. Ital. Derm. Sif.*, **67**, 108.
 Weidman, F. D. (1926). *Arch. Derm. Syph.*, Chicago, **13**, 374.
 — (1927a). *Ibid.*, **15**, 445.
 — (1927b). *Ibid.*, **15**, 427.
 — (1936). Personal communication.
 White, C., and Taub, S. J. (1932). *J. Amer. med. Ass.*, **98**, 524.
 White, C. J. (1927a). *Arch. Derm. Syph.*, Chicago, **15**, 889.
 — (1927b). *Ibid.*, **15**, 467.
 Wise, F., and Wolf, J. (1936). *Ibid.*, **34**, 1.
 Whitfield, A. (1908). *Lancet*, **2**, 237.
 — (1911). *Brit. J. Derm. Syph.*, **23**, 35.
 — (1935). Personal communication.

Clinical Memoranda

Pneumococcal Meningitis Treated with Prontosil Soluble

The case recorded below appears to afford further evidence of the efficacy of prontosil in the treatment of pneumococcal meningitis, and presents in addition several other points of interest.

CASE RECORD

A girl aged 5 years was admitted to hospital on February 22, 1938, with a provisional diagnosis of meningitis. For the past three weeks she had been suffering from whooping-cough.

On examination she was found to have a generalized purpuric rash, chiefly marked on the abdomen and limbs. There was rigidity of the neck muscles and head retraction, and a tendency to opisthotonos. Photophobia was present and Kernig's sign was strongly positive. The lungs, apart from a few catarrhal signs, appeared normal, and there was an apical systolic murmur. The temperature was 101° F., the pulse rate was 160, and the respirations 30. Haematuria was present. The child was very lethargic and wore an expression of frowning discomfort.

Lumbar puncture was performed, and the cerebrospinal fluid was found to be slightly turbid and under considerable pressure. The result of pathological examination was as follows: Cells, 11,200 per c.mm. (all polynuclear); chlorides 816 mg. per c.cm.; protein, 150 mg. per c.cm.; sugar absent; globulin reaction positive. Pneumococci in direct smears and on culture.