OCULAR HEADACHE

BY

**D. STENHOUSE STEWART, M.R.C.S.,
L.R.C.P., D.O.M.S.**

Honorary Assistant Ophthalmic Surgeon, Hull Royal Infirmary

Headache, together with constipation, and independently of it, may be claimed to be the most widespread effect of civilization. Headache is a symptom of which the pathology is speculative. It is believed that branches from the fifth nerve convey sensations of intracranial pain, and on this basis headache may arise by direct stimulation of these dural branches, sometimes in a manner not yet fully understood, or by the mechanism of referred pain from a focus of irritation elsewhere in the extensive area of trigeminal distribution.

These considerations are of little interest to the sufferer, and even less to the pharmacist from whom he obtains tablets, powders, and pills. A patient only consults his doctor if the headache is persistent, recurrent, or disabling, and seldom does a case of headache alone find its way to hospital, especially to a teaching hospital. Such a case is not likely to be used as clinical material, the diagnosis of the cause being commonly reached by exclusion, after which there is no case to demonstrate.

The student learns of headache as a casual concomitant of other conditions. He comes as a stranger to a case of disease, not as family doctor to patient, and he is unable to give fair weight to a subjective complaint, of which the reality and severity cannot be assessed with even relative accuracy. In general medical practice subsequently he will find that a patient desires relief of his symptom, easily becoming impatient, and the less reasonable from the very nature of his trouble.

Value of Systematic Investigation

"To waste no time in examining a patient" means to the lay public a mixture of clairvoyance and genius, by which treatment is given and relief obtained without any examination whatever. It should signify a rapid systematic inquiry and investigation rather than one which is inadequate and undirected. To the patient an ounce of treatment is worth a ton of diagnosis, but even on this basis if there be no diagnosis the treatment has no permanent value. This is particularly true in the case of headache.

Does the headache in any given case arise from dysfunction or disease of brain, kidneys, or rectum; eyes, nose, or teeth? However limited may be the time at his disposal, the practitioner has only to determine which of these is the most likely, and to find without difficulty someone eager to relieve him of any tedious further investigation in the particular sphere, but the specialist in any branch prefers to give his services or to accept his fee for positive results rather than for negative findings.

Headaches of ocular origin often fail to be recognized as such by the patient—as commonly perhaps as unsuccessful attempts are made to obtain relief from hyperpetic or renal headaches by buying spectacles. Faced by commercial publicity a patient is slow to learn that unnecessary suffering and expense may be avoided by consulting his doctor.

The Dental Act has ensured that the teeth of Britain shall receive none but competent attention. The Minister of Health believes that the eyes of insured persons are safeguarded by examination before the insurance practitioner issues the essential certificate for ophthalmic

benefit, and to secure that due heed is given to a request for full ophthalmic examination it is as well to cite the pathological condition found. Secretaries of approved societies are accustomed to regard certificates of incapacity as invalid if the cause is not stated, and it is possible (though not justifiable) that ophthalmic recommendations may be arbitrarily ignored if in the secretary's judgement they are incomplete.

It would be unreasonable to expect that with moderate equipment a general practitioner will make a comprehensive ophthalmic examination. He may be assisted to provide an adequate one by a summarized plan rather more comprehensive than that taught to the student in a crowded curriculum, a plan which will tend to exclude definite ocular causes of trouble.

In any case of chronic headache it is as necessary to examine the optic disks as the urine, blood pressure, or knee-jerks. With regular use of an electric ophthalmoscope the difficulties are rapidly overcome, even those presented by small pupils in the elderly where midriatics cannot safely be employed. The more normal disks one sees the more readily is an abnormality detected.

Measurement of Vision

It should be inquired whether each eye separately, with distance glasses if worn, can read 6/6 (that is, the seventh line of the usual Snellen test type from a distance of six metres, approximately twenty feet). Children under 10 years old may claim normal sight if at first trial they complete the sixth line, though letter knowledge varies with intellectual ability, and the mathematically or manually dexterous are often backward in spelling or reading. Sans-serif charts are usually more easily read, especially by children, who find the more complicated form of letters something outside their daily experience. Space is often limited in the consulting-room, but this is easily overcome by the use of types printed in the reverse, hung over the patient's head and read through a mirror on the opposite wall at a distance of ten feet. *It is not equivalent to employ smaller letters at a shorter distance.*

A visual acuity of 6/6 with either eye eliminates the possibility of uncorrected or under-corrected myopia. This is a matter of especial importance in juveniles and adolescents from the age of 10 to 25 years, but hardly concerns us here, as myopes as a class are not so liable to headache of ocular origin as are hypermetropes.

Hypermetropia and "Presbyopia"

A child or adolescent may be considerably hypermetropic, possibly unequally so in the two eyes, and yet read the distance test types accurately by exercise of his powers of focusing—powers which are considerable in infancy and which, it will be remembered, decline gradually throughout life until they become inadequate to focus close objects with comfort. The age at which this "presbyopia" is manifest depends upon how big a handicap of hypermetropia has to be made up by effort of accommodation, only the remaining surplus being really available. Any severe illness, accident, or operation, especially a gynaecological one, may leave accommodation permanently lessened if it occurs in the late thirties or after. Without actual measurement of any error of refraction or of limitation of accommodation it is fortunately possible to judge whether a discrepancy is present from the associated phenomenon of convergence. In focusing for near objects the normal individual instinctively converges his eyes to a proportionate degree, and if a handicap of hypermetropia already exists the individual must accommodate excessively and therefore converges excessively.

In a child, of course, this may lead to strabismus, transient and easily remediable at first, but if it be neglected even for a few weeks the child adapts himself to monocular vision and intractable amblyopia is liable to result.

Latent Strabismus

The established habit of binocular vision in adults prevents this excess of convergence from becoming apparent as a frank squint, but the effort of maintaining a satisfactory balance imposes a strain reflected in several ways. Common among them is the inability of students, otherwise capable, to concentrate upon uninteresting textbooks, a tendency of the aged to fall asleep over their newspapers, while those who persevere in close work in spite of their difficulty are very liable to experience headache.

If, in the course of an operation for squint under efficient local anaesthesia, traction by a squint hook is exerted upon an internal rectus muscle, one is struck by the fact that the discomfort experienced by the patient is described as headache.

The Use of the Convergometer

The headache due to latent squint at close range is commonly situated at the root of the nose or across the brow, and attacks of characteristic migraine or of

If, however, the lower arrow wavers even a little towards A on the right of the upper arrow—and certainly if it reaches this letter or beyond—the eye muscles are under a constant demand to exert a divergence (for which they have little capacity) up to and sometimes beyond the limit of comfort and almost of possibility, leading to headaches as the expression of such effort.

Correction by Eye-glasses

Great and perhaps undue stress has been laid upon small astigmatic errors as the cause of headache. The relief of headache by correction of such errors is often due to the restoration of normal muscle balance by adequate glasses. If a patient has bought for himself a pair of spectacles, estimated, fitted, and casually centred without correction of any muscle imbalance present, it is possible, should headaches persist, that he and his doctor will pursue their research for the cause elsewhere, and may part ultimately with expressions of mutual dissatisfaction. There is still some prejudice against the use of spectacles, and it has been suggested that numbers of people are wearing glasses unnecessarily as a result of commercial advertising campaigns. As long as all those not certifiably insane are assumed to be able, by exercise of their reason, to discriminate in such matters without expert guidance this possibility must exist, and the "eye

migrainoid type may ensue in a suitable subject. An ocular headache may be aggravated by close work and is seldom present on waking, but this history is not invariable. The binocular balance at reading distance is therefore a guide to the physiological comfort with which an individual performs close work, and there are several ways in which it can be tested. A practicable one is by the use of the convergometer. Its principles of construction were devised by the late Mr. Maddox many years ago, and they reflect his ingenuity in their simplicity. This very simplicity and the consequently inexpensive nature of the instrument may have prejudiced wider acceptance, for the wholesale manufacturers state that the demand has always been moderate. I am indebted to them (The Ellis Optical Company of Croydon) for the loan of the block.

A card bearing an arrow with a line of letters is seen through the eyepiece as two, one image being displaced upwards by a prism before the left eye and a second identical image being displaced downwards by a prism before the right eye. For the purpose of this test the patient wears reading glasses if he has them. A square of small newsprint pasted across the shaft of the arrow assists by concentrating the patient's accommodation on the card. If the lower arrow—that is, the one belonging to the right eye—points to the *left* of the upper arrow to some letter in the latter part of the alphabet all is reasonably well, for the internal recti have powers of convergence independently of those inseparable from accommodation and can make up for any lack in this respect.

massage" experts continue to reap the reward of their "successes" in enabling such patients to dispense with artificial aid. Patients often regard a headache as something too trivial to consult a doctor about, and prefer to buy acetylsalicylic acid tablets themselves, possibly imagining that their doctor can provide no better alternative.

Conclusion

The appreciation of pain and pleasure is very much a matter of temperament. Headaches are seldom incapacitating to those who may be described as "human vegetables," and when a phlegmatic, healthy individual complains diffidently of constant or frequent pains in the head investigation should be very thoroughly carried out. It must be a part of the treatment of the temperamental and neurotic patient to reduce to its true proportion each new symptom, whether of headache or any other kind; but to be effective this devaluation must follow such investigation as may be possible to exclude any real cause for such symptoms.

The use of test type and ophthalmoscope, and a rapid investigation of the lateral eye muscle balance at reading distance, will not reveal or exclude every ocular cause of headache, but they represent a simple, rapid, and sound method of ensuring as far as may be possible that appropriate ophthalmic attention is quickly given to those more obviously needing it, while others are referred for investigation to those quarters from which they are more likely to benefit.