produce a short period of anaesthesia. If too little is given the patient, when stimulated, may become quite uncontrollable; if more than sufficient is given the return to consciousness may be much delayed, and it may be some hours before the patient is fit to leave the surgery. With those addicted to alcohol, or in a highly nervous or apprehensive state, doses up to 5 or 6 c.cm. may be necessary.

The maintenance of an airway in a patient under evipan in the dental chair is particularly important. Obstruction such as is sometimes seen under nitrous oxide, when of necessity the dentist's fingers depress the tongue, cannot be tolerated. The amount of evipan given should never be enough to depress the respiration in any marked degree. Stimulation of the return of consciousness can be obtained by the inhalation of smelling salts or of carbon dioxide-oxygen mixtures.

Dental Operations in Nursing Homes

The induction of anaesthesia for dental operations in a nursing home, whether the patient is a child or adult, is similar to that for any major operation, and basal narcotics such as nembatal, avertin, and paraldehyde (which are described in another article in this series) may be used. To maintain anaesthesia ether vapour and oxygen or nitrous oxide and oxygen may be blown through a tube inserted through the nostril, leaving the mouth free for the dental surgeon. For the operation of removal of difficult wisdom teeth, in which the jaw is constantly depressed by the dentist, it is an advantage to employ an endotracheal tube, but for multiple extractions a short tube passed through one nostril into the oral pharynx will serve. A useful length for such a tube is the distance measured from the tip of the nose to the lobe of the patient's ear; the tube will not reach the vocal cords, and when employed it is necessary for the anaesthetist to support the mandible throughout the operation. If the patient's head is to be covered with towels care should be taken to see that the eyes are closed, as they may be injured by contact with such coverings in the movements of the head involved in a difficult extraction.

The progress of anaesthesia throughout the operation is estimated entirely by the respiration. The patient is maintained at a level of anaesthesia at which he does not cough when, from the manipulations of the dental surgeon's fingers on the throat pack, the posterior pharyngeal wall is irritated. At the conclusion of the operation the anaesthetic is discontinued and the throat pack removed. If the patient does not cough within a minute or so carbon dioxide and oxygen are blown through the nasal tube. On the return of the cough reflex the nasal or endotracheal tube is removed, but the mouth-prop is allowed to remain. If the prop is taken away too soon some patients tend to clench the teeth and allow the tongue to cause obstruction or grind the teeth, with the possible dislodgement of an inlay or filling. Once in bed the patient is turned well over to one or other side to ensure that any oozing from the sockets dribbles out of the mouth.

The following officers of the Women's Medical Service for India have received the Coronation Medal: Drs. H. M. Franklin, M. C. Murphy, E. Hamilton-Browne, H. M. Lazarus, Grace Stapleton, L. de Menezes, Ruth Young, G. Patel, N. Mucadam, Torrance-Allen, L. Ghosh, Roulston Mitton, K. McDermott, H. Keane, A. Dodhi, H. Acheson, D. Bolton, E. Wingate, U. Morton, I. Keess, N. Proctor-Sims, J. Orkney, C. L. Houlton.

"THE SURGEON'S LOG"

AN ANALYSIS OF THE HEALTH OF THE CREWS OF A LARGE FLEET OF MERCHANT SHIPS

BY

W. ARTHUR TATCHELL, M.R.C.S., L.R.C.P.

Medical Superintendent, Blue Funnel Line and Glen Line

There is no standardized "surgeon's log," each shipping company having its own form; nevertheless, though details may vary, the principal items recorded are the same. Our log is rather comprehensive, giving the name of the port where the ship is lying or to which it is bound, the numbers of the crew and passengers, the daily temperature, and a list of the sick, their treatment and progress. At the end of each voyage the log, with abstracts of cases and a list of accidents, however trivial, is either brought by the surgeon or forwarded to the medical superintendent at the head office of the company. There it is carefully examined by the medical superintendent and a summary of cases recorded. The log, when labelled and numbered, is placed in order with hundreds of others for future reference when necessary.

A Year's Statistics

These final statements, which I have made annually, are, I think, both enlightening and instructive, and it occurred to me that the publication of one year's statistics of the crews of a large merchant fleet would be of interest and possibly useful to those who are responsible for the health of seafaring men. To know what the principal diseases are from which seafaring men suffer and which cause them to be put off duty may help to guide those responsible for their health in an intelligent direction in the matter of prevention.

Health at sea, as on land, is largely ruled by conditions of living, food, and available treatment. The different standards for these set by owners will, I believe, be largely reflected in the entries in the surgeon's log. Climate must be taken into account. A North Atlantic company will probably have different results from one whose ships are trading for half their voyages within ten degrees of the equator. The fleet of the company concerned in the present analysis consists of eighty-five ships, varying in tonnage from 3,066 to 14,646 tons gross. The policy of the owners is to have a surgeon in each ship, although legally required to carry one in only fourteen of their ships.

The ships from which the statistics are taken sailed between the United Kingdom, and the Straits Settlements, China, Japan, the Dutch East Indies, and Australia. I have omitted for reasons of accuracy our Western Australia ships, which sail between Singapore and Fremantle, the transpacific service between Hong Kong and Vancouver, ships plying between New York and the Far East, and those flying the Dutch flag. The crews are composed of Europeans and Chinese; two ships were staffed entirely with Europeans. The Chinese are drawn principally from Hong Kong and Shanghai. All crews are carefully medically examined before signing on, the Chinese by the ship surgeon at the port of engagement. All are vaccinated at least once every three years, while the eyesight of each officer is tested once a year and a full medical examination made before each promotion. The food supplied is of good quality, well prepared, and not rationed.

The average time for the voyages was 119 days, and the average crew per ship was eighty-one, composed of forty-six Europeans and thirty-five Asiatics. In 1935 the total number of "men-days" at sea was 1,268,000, and the time lost through sickness and accident was only 4,786 days, giving the remarkably low percentage of 0.4. The average time lost per man was 1.26 days for Europeans and 0.69 day for Asiatics. This does not necessarily

include the days spent in hospital by the twenty-three Europeans and seventeen Asiatics who were landed abroad.

Chief Causes of Illness

The chief causes of illness were inflammatory and febrile conditions, especially influenza. There were only a few cases of malaria. The number of days on which Europeans were off duty from all febrile causes was 633, being 18.9 per cent. of the total days lost by Europeans from all causes—that is, 3,356. For Asiatics the figure was ninety-nine days, a percentage of the total of 1,430 days lost at sea of 6.9. Asiatics were less susceptible to influenza, only forty-six days being lost by them from this cause, or 3.2 per cent., whereas the corresponding figures for Europeans were 433, or 12.6 per cent. The same observation could be made with regard to "colds and coughs." It might be presumed that the Asiatics working in the hot engine room and stokehold and experiencing the great contrasts in temperature when they came on deck would suffer more than those whose work was on deck alone. But it appears that mere change of temperature, apart from other factors, is not prejudicial.

Injuries are classified under the headings of "fractures," "contusions," "wounds," and "burns and scalds," and the number of days off duty on board ship from all four injuries was 660 for Europeans, or 19.7 per cent., and 319, or 22.3 per cent., for Asiatics. Rheumatism and inflammation of joints seem to affect the Asiatics more than the Europeans. Neuralgia deserves mention with rheumatism, as *both* occur chiefly on encountering warm to hot weather. This is also the opinion of other observers.

Tropical diseases on board ship are very rare. Only six cases of malaria occurred among the 2,690 Europeans and three among the 2,065 Asiatics. There were seventeen cases of diarrhoea among Europeans and two among Asiatics; one European and one Asiatic had typhoid, and two Europeans had dysentery and enteritis (cholera). These were the only diseases of a tropical nature which were recorded, though it is possible that a few cases of coastal fever may have been classified with influenza as being clinically indistinguishable therefrom. The percentage loss of time from tropical diseases was 3.2 for Europeans and 2.5 for Asiatics.

The commonest types of skin diseases are those caused by various microsporons. Prickly heat rashes are confined almost entirely to the Tropics, and generally yield quickly to treatment with the use of fresh and the avoidance of salt water. Boils and pustular rashes are occasionally met with. Pediculosis is rare, there being adequate facilities on board the ships in the way of baths and washhouses for clothes to ensure personal cleanliness. Other types of skin troubles are so infrequent as to be represented by isolated cases. It may be noted that cuts, abrasions, ulcers, burns, and scalds, also most of the other skin affections, heal more slowly in the Tropics than in temperate climates. This may be attributable to the almost constant moist state of the skin rendering it impossible to keep lesions dry, and affording a more suitable medium for the growth of germs.

Effects of the Sun

There were eighteen cases of sun and heat strokes among Europeans and three among Asiatics. One must discriminate between the cases attributed to the direct rays of the sun and those due to heat. I think the vast majority of cases were, and are, due to heat. Real sun-stroke is reputed to occur at two periods of the day—namely, the rising and setting of the sun, whether the atmosphere be clear or dull, and the "stroke" is caused through either the eyes or the back of the neck. I have not come across a case at sea, but the two authentic cases I saw in China happened during the setting of the sun on dull days at an altitude of 3,500 feet. Heat stroke can occur at any period of the day, but seldom in the early

morning or evening. I may add that our officers have derived great benefit, including freedom from fatigue and headache, from wearing neutral-tinted glasses on the bridge and while working on deck during both bright and dull days. Racial susceptibility to disease is not well marked on board ship in our experience of the voyages considered. Pulmonary tuberculosis was rare, but a few cases were reported among the Chinese. Cases of beriberi, which is common in South China, were extremely rare. The latter are difficult to diagnose if the disease is in its early stages when the patient is examined, but any sufferer is soon restored to health with the good food supplies of meat, fish, eggs, and vegetables.

As to the morale of the crews, both the Asiatics and Europeans are a steady class of men. The amount of time lost through vicious propensities is negligible. Those members of the medical profession or general public who regard the seafarer as a hard-drinking morally lax person are entirely in error. At sea deck games are encouraged, and when in port football and cricket matches are frequently arranged; also other recreational facilities are provided in the company's clubs at various ports.

Conclusions

The different diseases, etc., recorded here are those causing loss of time only. There were many others mentioned in the logs, but their effect in this respect was insignificant. Considering the statistics as a whole and that the average time lost per man during the year was one day only, the conclusion must be that the seafaring life is a comparatively safe and healthy one on the routes and under the conditions considered where medical aid is constantly present. Is this a fair sample of the vast majority of trading routes and the experience of other shipping companies?

I wish to express my thanks for the valuable assistance of Dr. J. A. Tobin and my secretary, Mr. A. B. Tytler, in preparing the statistics from which this material has been compiled.

VITAL STATISTICS OF MALTA

The vital statistics of the Maltese islands (Malta and Gozo) showed a remarkable improvement in several respects in 1936, though the Chief Government Medical Officer (Dr. A. V. Bernard) in his annual report emphasizes the fact that in vital statistics based on small populations considerable fluctuations may occur in short periods due to ephemeral and fortuitous causes. The islands have a population of 262,165, which is about ten times as great as it was 1,000 years ago. The birth rate is steady at 33.8 per 1,000, and has not differed markedly during the last ten years; it stands where the birth rate for England and Wales stood fifty years ago, and where the birth rate for British India stands at present. The death rate for 1936 (17.61) shows a considerable drop, and compares with an average of 22.3 for the last nine years. The infant mortality, though still at the very high figure of 190 per 1,000 births, also registers a big fall from the average of the previous nine years of 280. The mortality of children under 5 has declined to 66 per 1,000 from rates which during the last ten years have approximated to 100. The high infant mortality accompanies a very high birth rate, and the population of Malta is steadily increasing at a more rapid rate than in countries where a low infant mortality is coupled with a diminishing birth rate. Of the notifiable diseases the commonest is undulant fever, of which there were 873 cases in 1936 (the lowest figure since 1927) and fifty-two deaths. The decrease in the incidence of this disease must be ascribed either to a diminution in the diffusibility and virulence of the infecting agent or to an increased resistance to the disease among a larger proportion of the population due to immunity derived from high prevalence. An undulant fever research station has been at work since the beginning of last year, and progress is also being made in the scheme for setting up a milk pasteurization plant.