ANAESTHESIA IN GENERAL PRACTICE

This concludes the series of articles on anaesthesia contributed by invitation

ANAESTHESIA AND THE LAW

BY

D. HARCOURT KITCHIN

Of Gray's Inn, Barrister-at-Law

The legal principles which govern the position of the anaesthetist are few and simple. He is bound to use reasonable care and skill, and he must not give an anaesthetic without first having obtained the patient's consent. Having said that, one has practically exhausted the law on the subject, and it only remains to consider some of the practical implications.

Reasonable Care and Skill

The degree of care and skill which is reasonable varies according to the circumstances in each case and according to the mind of the judge or jury. In practice, the standard required of a specialist is very high, and even a slight lapse by a man who makes anaesthesia his profession will render him liable to pay damages. If, however, as happens every day, the anaesthetist is a general practitioner, it is not reasonable to expect of him the same high degree of skill, and if the court is satisfied that he has taken every reasonable care and done his best it will not find that he has been negligent. At the same time, a reasonable degree of skill is always demanded, and a general practitioner who happens not to have given an anaesthetic for many years should not do so except in an emergency; if he has a mishap it will be no excuse to say that he did his best, because he should not undertake any task unless he has reason to believe that he can perform it safely. An error of judgement not due to lack of care or skill is not negligence; but when a doctor has actually made a mistake and is charged with negligence he will find it difficult to satisfy a jury of uninstructed laymen, and even a judge sitting alone, that his mistake was not due to negligence. The test question is always, Did the doctor do everything that could reasonably have been expected of him in the particular circumstances of the case?

Emergency may excuse a considerable amount of what might otherwise be negligence. Most medical men agree that only an emergency will justify a doctor in giving an anaesthetic without first examining the patient to see that he is fit to receive the type of anaesthetic it is proposed to use. Some authorities consider that, except in an emergency, a doctor should not act as surgeon and anaesthetist at the same time, but there are several exceptions to this rule in everyday practice. For example, it would be unreasonable to say that a dentist doing a simple extraction, or a doctor doing a minor operation, was negligent in not calling in a colleague to give the anaesthetic. In many parts of the country he would be unable to do so. Many doctors have to perform long and difficult obstetrical operations alone. The line will be drawn according to the evidence of what is considered reasonable by the consensus of medical opinion, and an emergency may justify a doctor in giving his own anaesthetic for an operation at which he ought not to give it in ordinary circumstances.

It is unwise to anaesthetize a woman unless some reliable third person—for example, a nurse—is either present or

within call, as, for instance, in an adjoining room with the door open or unlocked. The patient may charge the doctor with an indecent assault, and he may find it very difficult to defend himself.

Injuries associated with anaesthesia fall roughly into five classes: (1) death, which may not be due to the anaesthetic at all; (2) burns; (3) mechanical injury to the mouth, and other injuries due to swallowing or inhaling foreign bodies; (4) mishaps with local anaesthetics; and (5) injuries due to unconsciousness.

Anaesthetic Deaths

Death due to the anaesthetic itself is often promoted by the ill-health of the patient. Many operations are undertaken as a last resort; some are unsuccessful and no one is to blame, for the patient would certainly have died in any event. When, however, a patient dies from the effects of the anaesthesia alone, the two commonest causes are asphyxia from inattention to maintenance of the airway, with the action of the anaesthetic superadded; and overdose. Fatal injury results on rare occasions from rupture of the stomach caused by the insertion of an endotracheal tube into the oesophagus, if the mistake is not corrected. A number of deaths, though not many, are due to faulty assembly of the apparatus, so that a wrong mixture is delivered at the mouth-piece or unvaporized liquid is injected into the trachea. Anaesthetists have been known to use apparatus in which the gas cylinders were connected to the wrong inlets, or wrongly named, so that they gave carbon dioxide or nitrous oxide gas in mistake for oxygen. In spite of the ingenuity with which manufacturers attempt to make apparatus fool-proof, loopholes seem to persist. At least one accident has occurred because the bottles in a Shipway apparatus were not arranged in the way the anaesthetist expected. It is not safe to test Junker's apparatus only with the hand-bulb, because its feeble pressure cannot be relied on to drive the liquid anaesthetic up the inlet tube if the chamber has been connected up wrongly. The apparatus should be tested under working conditions, and the precautions taught in hospital carefully followed. One safeguard is for the anaesthetist himself to inhale a little of the mixture at the mouth-piece.

Patients have died under an anaesthetic from heart failure due to an overloaded stomach. It is the anaesthetist's duty to make sure that the patient is in a fit state to be anaesthetized, but he probably has the right to assume that the patient, or the hospital staff, has obeyed his clear instructions. In hospital practice, if an anaesthetist began to cross-examine the ward sisters about the preparation of every patient they brought to the operating theatre there would soon be a mutiny. One general principle of a doctor's liability for a nurse is that he may properly delegate certain tasks to her and assume she has done them. The preparation of a patient for operation is a task which is properly given to the nurse; the anaesthetist cannot be present to supervise her, and cannot, I submit, be made liable for the consequences of her negligence. It is probably his duty to make sure that routine general instructions have been issued to the ward staffs, but not to ask the sister every time whether they have been carried out. Nevertheless, the question whether in a particular case a particular task was rightly left to

a nurse, so that the doctor's responsibility for its proper performance passed entirely on to her shoulders, is nearly always a question of fact based largely on recognized practice, and not a question of law. A jury might conceivably, under the persuasion of a vigorous and skilful counsel, find that the anaesthetist ought to have asked the sister whether the patient had been properly prepared for the anaesthetic. The balance of probability, however, is rather against their taking this view.

The Coroner

The coroner is bound by law to hold an inquest on a person who seems to him to have died from an unnatural cause. Some patients die under an anaesthetic from perfectly natural causes. It is, however, difficult to decide which of many possible factors has caused the death of a given patient at an operation, and every registrar must, by order of the Registrar-General, report to the coroner any death which has occurred at an operation or before recovery from an anaesthetic. The coroner then has wide discretion to do what he thinks fit. He may, if he chooses, order a necropsy and an inquest, but he will nearly always wish first to consult the medical men in charge of the operation and hear their opinions about the causes of death. If he is satisfied that these were natural he will give permission for burial and do nothing more. If he thinks a post-mortem examination is necessary he will order one, and perhaps decide when he knows the result that an inquest is unnecessary. Strictly speaking, no citizen need notify any death to the coroner, for the duty is laid by statute upon the registrar. In practice, however, the surgeon or hospital authorities inform the coroner of every death of a patient under an anaesthetic, whether a death certificate is given or not. The coroner appreciates this courtesy. Even if death is certified as natural no necropsy should be performed without his permission, for he may still consider that an inquest is necessary.

Sydney Smith¹ says that the anaesthetist who is summoned by the coroner to give evidence at an inquest should prepare himself to answer the following questions:

Was the anaesthetic necessary?

Was it the most suitable one to give under the circumstances?

Was the patient properly examined before administration, and the condition of the heart, lungs, and urine ascertained?

Was there no contraindication?

Was the patient suitably prepared?

Was the anaesthetic administered with due care?

What amount of anaesthetic was given, and what was the duration of anaesthesia?

Were the necessary appliances for resuscitation at hand before commencing anaesthesia?

What treatment was actually adopted?

If the anaesthetist deals with all these points in his preliminary report to the coroner he will be giving that officer all the data necessary for his decision on what action to take, and will save him much trouble.

Burns and Anaesthetic Explosions

Burns due to the action of anaesthetic liquid on the skin, mucous membrane, or conjunctiva are fairly common. The face is sometimes burned when the mask becomes saturated and liquid overflows on to the skin. Some anaesthetists protect the patient's face with vaseline

or oil, and the eyes with castor oil; some cover the face with an elaborate dressing and apply strapping to the eyelids. Abrasion of the eye is an accident commonly met with during anaesthesia by the open method. It may occur in three distinct ways: (a) by the action of the liquid anaesthetic on the cornea or conjunctiva; (b) by the action of strong anaesthetic vapour, especially ether, when the patient is in a toxic or very debilitated state; (c) through over-handling of the cornea or conjunctiva by an inexperienced or nervous anaesthetist who feels that he must constantly reassure himself that the condition of the corneal reflex is satisfactory. Every precaution should be taken to prevent anaesthetic liquid from entering the conjunctival sac, and many anaesthetists drop castor oil into the eyes to prevent trouble if it does. When open ether is used it may be quite impossible to prevent a strong vapour reaching the eyeball, though every reasonable precaution should be taken. Patients often injure their eyes by rubbing them as they come round. To prevent this is part of the duty of the nurse, who should be with every patient who is under the influence of an anaesthetic.

ILLUSTRATIVE CASE

In a case of injury to the eye which attracted a certain amount of attention some years ago² the patient sued two medical partners for damages for loss of sight in his left eye. An operation was performed in hospital for acute obstruction; the patient was very ill and was on the table a long time. One partner gave the anaesthetic and the other assisted the surgeon. The anaesthetic consisted of two parts of ether to one of chloroform, and was given on an open mask, a Gamgee pad being placed on the face. This pad twice became so wet that the anaesthetist changed it. When the patient came round he complained of pain in his left eye, and some days later an ophthalmologist found that parts of the eye were very badly scarred. The expert evidence showed that the condition of the eye was consistent with damage done by the anaesthetic, also that liquid anaesthetic often entered a patient's eye but did not often harm it. The damage here was quite unusually serious. It was also fairly well established that liquid anaesthetic could get into an eye without any lack of care or skill by the anaesthetist. The surgeon said in evidence that there had been an epidemic of "anaesthetic eye" at one of his hospitals in the preceding year; the anaesthetics had been given by two highly experienced men who were absolutely certain that liquid anaesthetic had never entered any of the eyes. Most of the patients had been in a septic condition, and the eye trouble had cleared up within two or three days, except in one case where the eye had to be removed. It was not quite clear, from the mass of the evidence, whether any anaesthetic had entered the present patient's eye, but the concentration of vapour in the air had been very strong. The patient was in an extremely toxic condition. The jury failed to agree after two periods of retirement, and were discharged; there is no record of a retrial of the action.

Many anaesthetics are inflammable, and therefore apt to explode if they meet a red-hot cautery or spark. This risk was recently discussed in the *Journal*.³ Ether even when given by the open method is highly inflammable, but it is much more so when mixed with gas or oxygen. Explosions have been caused by badly fitting endoscopic lamps. In dry countries, such as the United States, explosion due to the discharge of static electricity is a recognized risk: tables and trolleys become charged by the friction of bedding, and powerful sparks are produced. The Joint Anaesthetics Committee of the British Medical Association and the Royal Society of Medicine recently requested the National Physical Laboratory to investigate this danger; a full report is promised for early

¹ *Forensic Medicine*. 1936. J. and A. Churchill Ltd.

² *British Medical Journal*, 1925, 1, 534.

³ 1937, 2, 383.

next year, and in the meantime the Laboratory recommend careful earthing of all equipment by a chain trailing on the floor or the use of electrically conducting rubber.

If a patient is injured in this way it seems very doubtful whether the anaesthetist can be made liable for the effects of his otherwise harmless mixture after it has been exploded by a diathermy needle or statically charged trolley, which was no concern of his. He could only be held liable if he had reason to suppose there was any peculiar danger in its use on that particular occasion.

The surgeon and the hospital governors are surely entitled in the first place to rely on the manufacturers of their electrical apparatus, as independent experts, to supply them with apparatus suitable and safe for the purpose for which it is to be used. None of them can be expected to be an expert electrician. It may, however, be the duty of the owner of the electrical apparatus used in the theatre, whether it be the surgeon or the governing body of the hospital, to see that it is periodically inspected and maintained by an expert technician to obviate the risk of a defect in it causing an explosion. It may perhaps also be the duty of the hospital governors, now that the risk is generally recognized and publicly known, to see that their theatre furniture is properly earthed to prevent it from accumulating dangerous static charges.

Gags and Foreign Bodies

In a dental operation under a general anaesthetic the patient is sometimes injured by the gag, or by a swab or other body entering the trachea, and the legal question may arise whether the dentist or the anaesthetist is responsible. If, for instance, a dentist does an extraction with an anaesthetist, a question which may arise is, Whose duty was it to pack the throat? In one case an assize court held that it was the duty of the dentist. No finding of responsibility in a particular case will ever amount to a rule of law binding another court. Each case is considered on its own merits. Nevertheless, in another case where the facts are similar the court may well come to a similar conclusion.

A dentist will often ask the anaesthetist to put in a gag. The anaesthetist will do this, and in doing so may accidentally dislodge some of the teeth. The question then arises whether he or the dentist is responsible for this injury. Both may well be, and the damage for interfering with healthy incisors may be serious, especially if the patient is a young woman and they have a high cosmetic value. Several anaesthetists have had to face claims of this kind. A dentist will often place a swab at the back of the throat to keep blood and foreign bodies out of the trachea. It is then his duty to take it away again, and if he has any difficulty in finding it he may ask the anaesthetist to help him. Moral, apparently, for every anaesthetist at a dental operation: Never, except in the direct emergency, step an inch outside your own province; let the dentist do his own work himself. In everyday practice an anaesthetist cannot be so strict. Everyone wants to help a colleague and, even more, to do his best for the patient. Nevertheless, this is a substantial legal risk, and anaesthetists have fallen into it. Injury has also been caused by the incautious use of too hot a gag.

Local Anaesthetics

Local anaesthetics also have their hazards. Mixtures may be made up carelessly, so that the patient receives an overdose of one of the constituents. Needles often

break, for the points are very fine and the shafts are flexible. The most usual point of fracture is at the butt, and the broken-off piece is very difficult to extract from the tissue. If an incision is made on to it, it is apt to move away. To guard against this known risk the doctor should not insert the needle to its full length; then, if it breaks at the butt, the broken piece will project and can be removed with forceps. A new needle, which enters easily, is less likely to break than an old, blunt, or tarnished one. Juries have been known to admit that a broken needle is not necessarily the result of negligence, but usually the presumption is the other way. The learned author of *Beven on Negligence*⁴ says:

"*Prima facie*, to sew up a sponge or an instrument in a patient after an operation is negligence. Very great care and method is to be observed in accounting for all the appliances used, and this in proportion to the easiness with which they may escape observation; but even here the fact that some needle or portion of an instrument has been left in a wound is not conclusive, but the conclusion from the fact must be determined by a jury on a view of the whole circumstances."

The technique of spinal anaesthesia is delicate and difficult, and one might expect to hear of a number of accidents and claims arising out of its use. None of the authorities I have consulted, however, seems to have heard of one, and this rarity seems to testify to the care and skill of those who use this method, and perhaps also to its essential safety in competent hands.

Accidents to Unconscious Patients

Hot water-bottles often burn unconscious patients, and sometimes the patient blames the anaesthetist. This kind of claim is fairly common, especially when the anaesthetist is also the patient's general practitioner. In hospital the duty of protecting the patient from this injury can properly be, and always is, delegated to a nurse, but in the home, even if a nurse is present, the general practitioner might reasonably be expected to supervise the patient's welfare before and after the operation.

Even before a patient goes under the influence of the anaesthetic he loses sensitivity and mobility, and a semi-conscious patient who is left with his feet near a radiator can easily be burned without noticing it or being able to attract attention. Patients are also insensitive after the operation, when they have gone back to bed. Many patients have been burned by bottles which have not been extremely hot, partly because their resistance had been lowered by the anaesthetic and the operation, and partly because they were unable to move out of reach. No patient who is under the influence of a general anaesthetic should ever be left alone for an instant.

Consent

To interfere, in however slight or innocent a way, with the person of another is technically an assault, and the assaulted party has a right to damages, which need not be in proportion to the gravity of the assault. Most forms of assault, however, are not legal wrongs if the person assaulted consents freely to them. It is undoubtedly an assault to deprive a person of his senses without his consent, and an anaesthetist who gives the anaesthetic without the consent of the patient takes the risk of being sued for damages. The same rule applies to a surgical operation, and it is the practice in most, if not all, hospitals for the surgeon first to obtain the patient's written consent on a form drawn up, for the

⁴ 1928, 2, 1359.

purpose. Some, but not all, hospitals use a form by which the patient consents to the anaesthetic as well as the operation. The question whether the anaesthesia and the operation can be separate assaults has probably never been raised at law. If a patient has given his consent in writing to a specified operation, no court is likely to give him damages on the ground that he did not also consent to the administration of the anaesthetic. It is an integral part of the operation, and his consent to it is necessarily implied. The question might possibly arise over a trivial operation for which a general anaesthetic was not absolutely necessary. If, however, the surgeon has told the patient he intends to give a general anaesthetic and the patient has had an opportunity of objecting and has not done so, he will in all probability be held to have consented. Both the surgeon and the anaesthetist run a definite risk if the patient is arbitrarily anaesthetized for a trivial operation without being given the opportunity of objecting. If an accident happens and he dies under the anaesthetic or suffers injury, both medical men are likely to have to pay damages, whether they have been negligent or not. If an operation is urgently necessary and the patient is unconscious, and no relative or friend is available to give consent in his place, the emergency will justify the assault.

It is possible that anaesthetists in general take the patient's formal consent to operation too much for granted. If an aggrieved patient or his widow can show afterwards that he did not give consent to the operation the anaesthetist as well as the surgeon will have been guilty of an assault and will be liable to pay damages; if the injury has been due to the anaesthetic these may be heavy. If, however, the substantial injury has been caused by the operative procedure, the assault committed by the anaesthetist is probably a merely technical one and damages will at the most be nominal. I have never heard of a case in which a patient suing a surgeon for operation without consent has also sued the anaesthetist—nor, indeed, of one in which an anaesthetist has been sued for anaesthetizing without consent; but the risk is not an imaginary one. When the patient is under 21 years of age he is a minor, and his parent or guardian must give consent for him.

Liability for Nurses' Mistakes

Nurses sometimes make mistakes which lead to serious injury or death. It is not often worth the while of a patient or his widow to sue the nurse for damages, and he will therefore usually try to show that some medical man was responsible for her negligence. The general rule is that the doctor is responsible for the negligence of his assistants. If, however, the nurse is doing skilled work which may properly be delegated to her and the doctor is not in a position to supervise her, she alone is responsible. Here again the circumstances of the particular case are all-important, and there are no rules of law governing particular circumstances. The principle is discussed in the leading cases of *Perionowsky v. Freeman* (1866 4 F. and F., 977), *Hillyer v. St. Bartholomew's Hospital* (1909, 2 K.B., 820), and *Anderson v. Glasgow Royal Infirmary* (1932, S.C., 245). The question often arises in cases of burning from hot water-bottles, and sometimes when a nurse has filled a bottle with the wrong anaesthetic fluid or given the anaesthetist a wrong bottle.

Liability of the Surgeon

A question of great importance to the defence societies is whether a surgeon is unconditionally liable for the negligence of every member of the theatre team at an

operation, including the anaesthetist. The only legal basis for asserting that he is so liable is contained in the famous dictum of Lord Justice Kennedy in the case of *Hillyer v. St. Bartholomew's Hospital* to the effect that the moment the door of the operating theatre closes behind the theatre team they cease for the time to be the servants of the hospital governors, inasmuch as they take their orders from the operating surgeon alone. The question at issue was whether the hospital governors were liable for an injury which a patient had suffered during an operation, and it was not necessary for the decision of that question that the court should express any opinion on the liability of a surgeon for the negligence of one of the theatre team. Whether the surgeon is liable for the anaesthetist's negligence or not is, however, immaterial to the question of the liability of the anaesthetist. Whatever the joint liability of the surgeon, the anaesthetist is responsible for his own errors. Whether he is also responsible for those of the nurses assisting him depends on whether he may reasonably be expected to supervise them in the particular task in question.

Surgeons sometimes prescribe a particular type of pre-medication which will affect the anaesthetist's choice of an anaesthetic. They occasionally go so far as to direct what type of anaesthetic should be used. Some day a mishap may well result directly from this restriction, and the legal question of whether the anaesthetist is a completely independent expert or is a subordinate of the surgeon will have to be argued, perhaps for the first time. When a question is still unanswered by the decision of a court every person must answer it in his own way. My tentative opinion is that as a matter of everyday fact the anaesthetist is independent and supreme in his own department, and the law ought to follow this fact. If this view is right, then the anaesthetist cannot shelter from legal responsibility behind these acts of the surgeon. If he considers that any requirement of the surgeon will make the giving of the anaesthetic unsafe he must say so and decline to give it, or else take the risk and shoulder the liability. In practice, any particular conflict of requirements can doubtless be solved by co-operation.

Medical Protection

Claims are not made as often against anaesthetists as against surgeons, but the anaesthetist's risk of being sued is substantial and constant. Every anaesthetist should therefore be a member of one of the large defence organizations, so that if a patient claims compensation from him he may fall back on its experience and resources. He will not only insure himself against damages and legal expenses, but will also avoid serious anxiety and loss of time. Moreover, the experts of these bodies are much better qualified than he is to deal with the demands of a claimant's solicitors. Finally, anaesthetists are recommended to read the sections in Taylor's *Medical Jurisprudence* which deal with their legal position.

I am much indebted to a number of medical and legal friends who have kindly helped me from their experience of these problems.

T. Hryntschak (*Med Klinik*, August 13, 1937) discusses the value of mandelic acid treatment in *B. coli* urinary infections. The usual preparation, ammonium mandelate, is often badly tolerated. The author has therefore tried another salt, calcium mandelate, and found it better tolerated and equally efficient. Another well tolerated and efficient preparation is sodium mandelate, which is given intravenously together with an equal amount of hexamine.