BRITISH MEDICAL JOURNAL

LONDON SATURDAY FEBRUARY 6 1937

CEREBRAL VASCULAR ACCIDENTS*

BY

C. M. HINDS HOWELL, M.D., F.R.C.P.

Physician to St. Bartholomew's Hospital and the National Hospital, Queen Square

I have chosen this subject for discussion because the vascular accidents to which I shall refer are matters of daily occurrence, are dramatic in their manifestations, and not infrequently give rise to much difficulty in diagnosis. Both the physician and the surgeon may be concerned in their treatment, and among these cases there will be many which will test our skill and experience to the utmost if they are to be brought to a satisfactory conclusion.

We can divide vascular accidents within the skull into those due to injury and those which result from disease processes. As the result of injury to the head there may follow concussion, haemorrhage, contusion, or laceration of the brain. With any of these there may or may not be fracture of the skull.

It is a matter of interest that in compensation cases, which are all too common nowadays, the legal mind appears to attach more importance to the fact of fracture of the skull than to what takes place inside it. This is a misleading point of view, for we know very well that the skull may be fractured without anything very serious happening within it, while the most damaging events may have occurred where no fracture has taken place.

It is seldom that haemorrhage occurs in an uncomplicated manner as the result of serious head injury. Concussion at least is of common occurrence, while contusion or laceration of the brain also frequently occurs. These events complicate the clinical picture, and make exact diagnosis difficult or impossible. I propose to consider first some uncomplicated cases which will illustrate the effects of haemorrhage in two common situations as the result of injury; and later the general principles which should guide us in more difficult cases.

Extradural Haemorrhage

In an uncomplicated case the patient receives some injury to the head, as the result of which he may suffer from a short period of concussion. He recovers from this, and after an interval of varying but short duration he develops symptoms of cerebral compression. There follow headache, vomiting, slowing of the pulse, which is full with a rising blood pressure, perhaps paralysis, then unconsciousness, and if the pressure be not relieved the next stage is death. The interval between the original injury and the later development of cerebral compression is the so-called lucid interval. All cases of extradural haemorrhage are fortunately not so serious as our hypothetical case, for the haemorrhage may cease before it causes serious compression.

Some years ago I was consulted by a young man who, while batting at cricket, had been struck on the left temple by the ball. He was knocked over, and for a few minutes was dazed

(period of mild concussion). He was able to continue playing for the rest of the afternoon, though his head ached (lucid interval). After the game his headache became much worse and he vomited (period of slight cerebral compression).

He continued to have headache, and three or four days later had a fit, which only involved the right side of his face. He was not unconscious in the fit, but could not speak during the attack. He had had several similar attacks, and to my great satisfaction had one in my consulting room. It was a typical Jacksonian fit, affecting the right side of the face only. The man was able to walk about during the attack, obey orders, and in fact do everything except talk.

In this case the diagnosis of extradural haemorrhage was not difficult to make. An operation was performed at the site of injury, and an extradural clot found and removed. The dura was opened, but no bruising or haemorrhage was present beneath it. The man made an uneventful recovery, but six months later had two generalized epileptic attacks. He was put on bromides for about eighteen months, but has had no further attacks.

This patient should have been put on some sedative—luminal 1 grain every night, for example—immediately after his operation. It is most important to remember that cases of this kind require this type of medical treatment even after a successful operation, such treatment being continued for a period of six months at least.

Subdural Haemorrhage

This may be diffuse or localized. In the former, which is usually associated with severe head injury, a venous sinus may be torn, when blood will diffuse rapidly through the subdural space, and symptoms of compression are likely to develop quickly.

The localized form of subdural haemorrhage is by no means uncommon, and it is important to recognize the clinical picture to which it gives rise. I will briefly describe two cases of this kind, one in an elderly subject, the second in a young man.

In the first case a man of 65 was hurrying to catch a bus when he was knocked down by a boy on a bicycle. He hit his head, was not unconscious but rather dazed. He was kept in bed for two or three days, but did not pick up. He complained of headache, and was irritable. This state of things continued for about a month, at the end of which he began to lose power in his left leg, and later in the left arm. He became drowsy and confused. When I saw him he had a low-grade papilloedema in both eyes and a slight left hemiplegia. I got Professor Ross to operate on him for chronic subdural haematoma. We found a large clot, wrapped in the peculiar membrane which forms in such cases. The underlying brain was guttered by the pressure of the clot. Next morning he was much brighter, but later became comatose and died. At the post-mortem examination a second similar but larger clot was found over the left cortex.

Lessons to be learnt from this case are that in elderly people a comparatively slight injury may cause a localized

* Part of a British Medical Association Lecture given to the Blackpool Division, November 18, 1936.

subdural haemorrhage, and that such haemorrhage is not infrequently *bilateral*. In this case there was an interval of four or five weeks before signs of paralysis appeared, but the man had never been well since his accident.

The second case I want to mention occurred in a young man who crashed on a motor bicycle. It was a very serious accident. The pillion rider was killed, and my patient lost a leg and was concussed. He recovered from the amputation, but continued to suffer from headaches. The accident occurred in August, and in November I was asked to see him as his headaches were still severe. I could find no signs of abnormality of the nervous system, but sent him into hospital for observation.

He improved very much in hospital under no particular treatment beyond rest in bed and one lumbar puncture for diagnostic purposes. The cerebro-spinal fluid was normal but under increased pressure (200 mm.). At the end of December the severe headaches returned and he developed papilloedema first in one eye, then in the other.

A diagnosis of chronic subdural haematoma was made and Professor Ross did a bilateral ventricular puncture, as there was nothing to show which side the haematoma was on. No fluid could be obtained from the right lateral ventricle; the left was found easily. We concluded the haematoma was on the right side; a bone flap was made, the haematoma found and evacuated, and the man made a good recovery.

In this case a period of four months had elapsed between the date of the accident and the development of definite signs of compression which the appearance of bilateral papilloedema indicated. In the circumstances the symptoms due to subdural haematoma might have been mistaken for those of so-called traumatic neurasthenia. The cases illustrate the importance of keeping patients who complain of recurring headache after a severe or even slight concussion under close observation. It is essential to examine the eye grounds regularly in order to detect the first signs of papilloedema in cases of this kind. And it is important to note that several weeks may elapse before anything of the sort appears.

The pathology of this condition is remarkable. A membrane develops round the extravasated blood in the subdural space. This membrane becomes vascularized, and a slow oozing of blood appears to occur, as a result of which the haematoma gradually increases in size and so comes to produce a slowly increasing compression of the brain.

As the result there is a latent period between the date of the injury and the development of obvious signs of compression. This latent period may vary in duration from a few weeks to a few months. During the latent period the patient is likely to exhibit symptoms which may be mistaken for those of traumatic neurasthenia. Of these symptoms headache, irritability, lack of concentration, and some memory defect are the commonest.

In cases where the clinical picture of haemorrhage is complicated by contusion or laceration of the brain the essential sign to watch for is that of cerebral compression.

In the earlier stages, after the initial period of shock, this is characterized by headache, vomiting, a full bounding pulse, and rising blood pressure. There may be signs of cerebral irritation either in the form of restlessness, delirium, or excitement, or perhaps by the occurrence of convulsions. As the compression increases the patient becomes unconscious (if he has previously been conscious), the pupils dilate, breathing becomes irregular or Cheyne-Stokes in character, the pulse, which was full and slow, becomes feeble and rapid, and the blood pressure, which rises in the earlier stages of compression, will fall, and if the condition is unrelieved death will follow.

It is important, therefore, to keep a close watch on the pulse rate and blood pressure. Lumbar puncture should be done, and the pressure of cerebro-spinal fluid measured. Any figure above 150 mm. indicates an increased intracranial pressure, and sometimes the pressure will be found to be as high as 300 mm. or even more. Rawling lays stress on the temperature chart, regarding a rising temperature as of serious significance.

Leaking Cerebral Aneurysm

Let us next consider haemorrhage outside the cerebral substance occurring as the result of disease processes. The commonest of these, though fortunately none are very common, is subarachnoid haemorrhage coming from a leaking aneurysm. Haemorrhages may occur under other circumstances, such as in the leukaemias, in purpura, and sometimes in septicaemia, but aneurysm is without doubt the commonest cause.

Cerebral aneurysms occur almost entirely in the vessels which constitute the circle of Willis or its branches, and therefore lie outside the brain substance. They may be the result of various morbid conditions, but the congenital aneurysm is the commonest. These aneurysms arise at the bifurcation of the arteries. Microscopical examination shows that the muscular coat of the artery is defective in such cases at the arterial bifurcation, which allows of the formation of the aneurysms. The aneurysms so formed are usually small. The name "berry aneurysm" has been given them. They may be multiple. Other causes of aneurysm of the cerebral arteries are arteriosclerosis and syphilis, though the latter are rare. Infective endocarditis produces mycotic aneurysm as the result of embolus. Trauma may also produce an arterio-venous aneurysm, usually from injury to the internal carotid artery in the cavernous sinus.

It is quite impossible as a rule to diagnose with certainty the presence of an aneurysm of a cerebral artery, previous to its rupture or leaking, from the clinical symptoms alone. X-ray photographs will sometimes show concentric circular shadows of calcified material in the wall of the aneurysm which are known as Albi's rings. Where an aneurysm is suspected the injection into the internal carotid of thorotrast, followed immediately by x-ray photography, will show this substance within the arteries, and in this way the presence of an aneurysm may be detected.

Subarachnoid haemorrhage from aneurysms seems to me to be commoner nowadays than when I was a student and house-physician thirty years or more ago, but I suppose the true explanation is that we are now more on the look out for such cases. The symptoms of a leaking cerebral aneurysm are usually dramatic. The patient calls out in pain, and may then fall unconscious; indeed, the diagnosis is often made on the history of a cry of "Oh! my head!" followed by unconsciousness. This diagnosis will be confirmed by finding uniformly bloodstained cerebro-spinal fluid if lumbar puncture is performed within a day or two of the accident, after which date the fluid becomes straw-coloured. If there is any doubt as to whether the blood is the result of the lumbar puncture this can be cleared up by collecting samples in two or three test tubes and comparing the tint in each. On standing the supernatant fluid will be yellow-tinted in subarachnoid haemorrhage, but colourless in cases of traumatic puncture. An interesting point to remember is that the urine often contains a heavy cloud of albumin for twenty-four hours or so after subarachnoid haemorrhage. If the patient is found unconscious this may suggest uraemia, but examination of the cerebro-spinal fluid will clear up

the diagnosis. Of course examination of the blood urea would do so too.

Occasionally glycosuria may result from subarachnoid haemorrhage. I remember such a case well at St. Bartholomew's Hospital. We were on duty, and my house-physician told me that he had admitted a patient in diabetic coma. The patient was brought into hospital unconscious, and the house-physician had found sugar and acetone in the urine in large quantity. He had given intravenous glucose saline and 50 units of insulin, but he reported that the patient was still semi-conscious, and apparently had some rigidity of the neck. I told him that if he would do lumbar puncture he would almost certainly find a uniformly blood-stained fluid, and such proved to be the case.

Rigidity of the neck, which is a very common feature of these cases, is due to irritation of the posterior roots in the cervical region by the extravasated blood, and often patients complain later of pains in the legs due to a similar cause. A cerebral aneurysm may produce palsies of some of the cranial nerves before rupture, those most likely to be affected being the sixth and third nerves.

A woman was admitted under my care at St. Bartholomew's Hospital with a subarachnoid haemorrhage. She made excellent progress and was due for discharge when she developed a third nerve paralysis in the right eye, and the next day had a second subarachnoid haemorrhage. She ultimately recovered from this, and for the last few years has been working as a cook-housekeeper, but the third nerve palsy persists.

Some patients have had as many as four or five haemorrhages recurring at intervals and have survived, but the condition is a very dangerous one, and the first attack may be fatal.

The youngest patient I have seen with subarachnoid haemorrhage was aged 9; he made a good recovery and is leading an ordinary life. Although a cerebral aneurysm may rupture at any age it is commonest between the ages of 20 and 40. Another point in connexion with aneurysm of the cerebral vessels is that some of them attain a large size and give rise to symptoms suggestive of tumour. You will find it stated, I think with decreasing frequency, that a bruit can be heard on auscultation where a cerebral aneurysm is present. I believe this statement to be quite untrue, except in the case of arterio-venous aneurysms, in which the patient is himself conscious of the bruit.

Accidents Due to Cardiovascular Disease

The common examples of vascular accidents in the brain which result from disease of the cardiovascular system are thrombosis, haemorrhage, or embolism of the cerebral vessels. But there is another group of cases of great interest, though by no means so common as those just mentioned, which I should like to deal with first. The condition has been called by McAlpine "the hypertensive attack." These occur in patients suffering from high blood pressure with or without obvious renal disease. I do not include among them cases of frank uraemia, in which the non-protein nitrogen in the blood is definitely raised, though it is probable that the cerebral manifestations that occur in uraemia—such as convulsions, temporary paralysis, amaurosis, and coma—are due in the main to a rise of intracranial pressure, associated with oedema and angiospasm, such as occurs in the cases to which I am referring.

Patients with high blood pressures in this group are liable to exacerbation of this pressure, with which obvious cerebral symptoms may be associated. In these attacks the patient complains of severe headache, and there may be in addition vomiting, drowsiness, or even coma and

sometimes convulsions. Transient paralysis may also occur. In some of these cases examination will show papilloedema, suggesting the presence of a cerebral tumour unless the picture presented by the fundus is considered very carefully and the associated blood pressure also taken into account. The appearance of the disk is not really like that seen with cerebral tumour, for haemorrhages form a striking part of the picture in addition to the swollen disk, and the vessels of the retina show marked nipping of the veins at the crossings, and the calibre of the arteries is seen to vary very much in different parts of the same vessel, due to angiospasm.

McAlpine has pointed out that young patients are more likely to suffer from cerebral oedema with papilloedema than older ones, and has demonstrated that the attack is associated with a very definite increase in blood pressure, the subsidence of which is followed by relief of symptoms.

Let us now consider some questions which always arise in connexion with embolism, thrombosis, or haemorrhage of cerebral vessels.

Cerebral Embolism

First, as to diagnosis. This is sometimes obvious, but often a matter of great uncertainty. As a rule embolus can be diagnosed with more certainty than the others. The embolus may come from a diseased valve, usually in this case the site of bacterial endocarditis. In cases of mitral disease it is more likely to be a clot detached from a dilated left auricle. But the presence of obvious valvular disease and the very sudden onset of symptoms usually leave no doubt as to the diagnosis. It is naturally likely to occur at an earlier age than either thrombosis or haemorrhage. We must not, however, overlook the possibility of embolism occurring from the surface of an atheromatous plaque in the aorta of older subjects. Apart from valvular disease it may arise as a sequel to dilatation of the heart, whatever the cause of such dilatation may be, for clots may form under such circumstances in what we might describe as the backwaters of the heart. This occurs sometimes in the puerperal state, so that embolism has a rather higher incidence in women than in men, for this reason and because of the higher incidence of mitral stenosis in women.

Following embolism of a cerebral artery there may be consecutive thrombosis, so that the symptoms at the onset become permanently established. On the other hand, if the clot be a small one it may retract and allow the blocked vessel to become patent again, when great improvement in the condition will follow, or if no secondary thrombosis results the symptoms are likely to improve as the oedema caused by the infarction subsides.

Cerebral Thrombosis and Haemorrhage

Thrombosis and haemorrhage both result from disease of the cerebral arteries. It may be easy to distinguish between them clinically, but, as you all know, it is often impossible to do so. There is a school of thought, to which I cannot subscribe, which maintains that all cases of cerebral haemorrhage are fatal. This is a comfortable doctrine, because if you believe it the treatment for all cases becomes the treatment for cerebral thrombosis. Let us consider to what extent it is possible to differentiate these cases with any degree of certainty.

In a young adult hemiplegia of sudden onset is more often due to syphilitic thrombosis than to anything else, provided you can exclude embolus, tumour of the brain, and disseminated sclerosis. I am assuming that hysteria is not in question. Subarachnoid haemorrhage and embolus we have already considered. Tumour of the

brain will usually have given some signs of its presence before causing a sudden exacerbation, but disseminated sclerosis is a more difficult thing to exclude. It is well known that the first onset of this disease may be by the appearance of hemiplegia. Serological tests will establish the diagnosis in syphilitic disease, and the proof should be obtained immediately, as energetic anti-specific treatment will do much to improve the condition in syphilitic thrombosis.

Thrombosis is also likely when arterial disease is associated with low blood pressure. It is strongly suggested when the attack occurs during sleep, or following exhaustion or in debilitating conditions generally. Often there are premonitory symptoms, such as numbness, tingling, and temporary loss of function, such as aphasia, monoplegia, and so on. But with regard to these I must point out that they may happen when the blood pressure is unduly high and as the result of temporary angiospasm in the arteries, and so these attacks may be followed either by thrombosis or by haemorrhage. They are always of serious import.

Of cerebral haemorrhage it may be said that it is uncommon with low blood pressure, and that usually the pressure is high, but it may not be excessively high; that it is more likely to take place during the day than the night, or in association with effort or strain; and that it is almost invariably accompanied by vomiting.

Of the three embolism is most sudden in its onset and thrombosis likely to be least. Consciousness may be lost as the result of any of the processes we are considering, provided the initial lesion is severe enough. It is probably true to say that it is more likely to be lost early in haemorrhage than in thrombosis, and early loss of consciousness which is prolonged almost always indicates a severe haemorrhage. The cerebro-spinal fluid in a serious haemorrhage is apt to contain blood, and it is good practice to do a lumbar puncture in any case of apoplexy with loss of consciousness, as it tends to lower the intracranial pressure.

Convulsions at the onset of apoplexy indicate irritation of the cortex, and are more commonly due to thrombosis than to embolism, though they may occur in the latter. They are very unusual at the onset of a cerebral haemorrhage.

Severity of the Lesion

The severity of the lesion can be estimated from consideration of three circumstances: the depth of the coma, the degree of depression of the nervous system which has followed the apoplexy, and the condition of the respiratory centre.

The depth of coma can be roughly estimated by seeing whether the patient responds to any cutaneous irritation. This may also serve to show the affected side if this is in doubt. The depression of the nervous system can be estimated by observing whether there is a complete flaccid paralysis of all the limbs, with loss of reflex action. If plantar reflexes can be obtained both are likely to be extensor at this stage. It is important to remember that even where the depression of the nervous system is not extreme retention of urine is common, and that it is important to prevent over-distension of the bladder.

Breathing is apt to be noisy and stertorous in proportion to the severity of the lesion, and grouped respiration of the Cheyne-Stokes type is a sign of failing activity in the respiratory centre.

The area of the brain supplied by the middle cerebral artery is the seat of election for all forms of apoplexy,

but while it is most unusual for embolism to occur elsewhere haemorrhage occasionally does so, and thrombosis more often still.

Prognosis in Apoplexy

In any case of apoplexy a prognosis will be anxiously sought by the relatives of the patient. The answer must deal with the immediate question of survival and the ultimate prospects of recovery of function. If the patient is deeply unconscious and clearly suffering from a cerebral haemorrhage a bad prognosis must be given. Such a patient will not survive. And in any patient in whom unconsciousness persists over twenty-four hours the prospects of survival are poor, and if survival occurs the chances of any useful recovery of function in the affected limbs are practically nil.

In elderly people the condition of the lungs, the heart, and the kidneys must all be considered carefully in coming to any conclusion as to their survival. So too must the condition of their arteries and their blood pressure. One must be very guarded in estimating the prospects of recovery of function during the early stages. If recovery of useful function is going to take place it will show signs of doing so at any rate by the end of a week. Quite obviously the sooner some recovery of power is noted the better the prospects. Cases of syphilitic thrombosis have probably the best chance of recovery if anti-specific treatment is efficiently carried out.

In thrombosis particularly, and in embolism to a less extent, one must remember the possibility that fits affecting the paralysed side will occur. These are as a rule infrequent, and may cease after a time. A patient whom I had under close observation for more than six years, suffering from a severe thrombotic hemiplegia, started to have unilateral fits affecting the paralysed limbs nine months after his apoplexy. These continued at intervals of six weeks to two months for about eighteen months, after which they ceased entirely.

P. Hauptstein (*Dtsch. med. Wschr.*, August 28, 1936, p. 1417) has conducted at the University Maternity Hospital in Freiburg i. Br. clinical investigations of the action on dysmenorrhoea of finely powdered potentilla anserina, whose popular name in Germany is "Gänsefingerkraut" or "Gänsekraut." Its medicinal properties were exploited in the Middle Ages, and pharmacological justification for its prescription as an antispasmodic was provided in 1933 (*Zbl. Gynäk.*, Nr. 37) by Schneider and Nevinny, who found it capable of relieving the barium cramp induced in the muscles of rabbits and rats. The author gave it in the form of capsules, each of which contained 0.5 gram of the finely powdered drug: From one to three capsules were given by the mouth three times a day while menstruation was painful and for two to three days before it was due. Even when large quantities were taken no gastric disturbances were observed, but it was found well to warn the patients against chewing the capsules on account of the disagreeable taste of their contents. The complete harmlessness of the drug and its cheapness, combined with the relief of pain it affords, make it, in the author's opinion, a valuable remedy. Though he does not state the actual number of cases thus treated, he says that about two-thirds of his patients were under the age of 30, and that only those patients were thus treated who came to hospital complaining of severe dysmenorrhoea, which, in many cases, had hitherto entailed rest in bed during menstruation. The results were most satisfactory in essential dysmenorrhoea associated with anatomically normal conditions or with hypoplasia of the uterus. But even when the dysmenorrhoea could be traced to chronic inflammation or displacements of the uterus, the results were so encouraging that the author recommends further investigations also in this field.