DEATH FOLLOWING BLOOD TRANSFUSION

NOTES ON TWO CASES

BY

F. PYGOTT, M.B., D.P.H.

*Late Assistant Resident Medical Officer, Mill Road
Infirmary, Liverpool*

Dr. N. S. Plummer (1936) has recently drawn the attention of the profession to the danger to life which may occasionally result from the transfusion of a patient with compatible blood. To the five cases he has reported I should like to add two more which have occurred within my own experience.

Case I

This was a case of aplastic anaemia in which death followed within five minutes of a third transfusion. The patient was a male, aged 42 years, admitted to hospital complaining of increasing pallor and dyspnoea on exertion for one month. There was a past history of syphilis eighteen years before, but the Wassermann had become negative after treatment over a period of two years.

His nutrition was good, though he was extremely pale and anaemic. There was no oedema of the ankles. He gave no history of loss of blood from any source. There was no enlargement of liver, spleen, or lymph glands. His pulse was regular and forcible. His heart was not enlarged clinically, and the sounds were regular and of good quality. His blood pressure was 175/95 mm. Hg. His urine contained no excess of urobilin; his icterus index was 5 and his blood Wassermann negative; his blood urea was 30 mg. per 100 c.cm. Examination of the blood on the first day of admission to hospital showed: red blood corpuscles, 1,400,000; haemoglobin, 20 per cent.; colour index, 0.7; white blood corpuscles 5,000 per c.mm. A differential count revealed: polymorphonuclear leucocytes, 18 per cent.; small lymphocytes, 80 per cent.; large lymphocytes, 2 per cent. The blood group was II (Moss), the bleeding time half an hour, and the coagulation time five minutes.

He was given 90 grains of ferri et ammon. cit. daily. As he showed no signs of improvement 550 c.cm. of Group II blood was administered by direct transfusion (Joubert method) on the seventh day after admission. No sequelae resulted. The day after the transfusion the blood count was: red blood corpuscles, 1,300,000; haemoglobin, 20 per cent.; colour index, 0.7; white blood corpuscles, 6,000 per c.mm.

On the tenth day there was a further direct transfusion of 550 c.cm. of Group II blood; again no sequelae resulted. On the thirteenth day a blood count showed: red blood corpuscles, 860,000; haemoglobin, 12 per cent.; colour index, 0.7; size of red cells, 8.2 μ ; white blood corpuscles, 4,000 per c.mm.; polymorphs, 27 per cent.; small lymphocytes, 69 per cent.; large lymphocytes, 4 per cent. Haemorrhages were now to be seen in both fundi and round the margins of the lips. His blood icterus index was 3.

The third and fatal transfusion was on the fourteenth day after admission. It is important to mention that in spite of his low haemoglobin he was quite able to sit up in bed without assistance immediately before the transfusion was begun. As on the two previous occasions the direct method was used. After about 150 c.cm. of blood had been transfused he became increasingly restless. This restlessness had been observed on the two previous occasions at about the same stage of the operation, but had passed off after a pause had been made. The operation was accordingly suspended for a minute or so, and then, encouraged by a slight improvement in the patient's condition, a further 50 c.cm. of blood was introduced, making a total of 200 c.cm.

His condition was now alarming; respirations were becoming rapidly intermittent, his pupils were starting to dilate, the pulse was absent at the wrist, and the heart sounds were faint. Intracardiac stimulants (adrenaline and strychnine) were of no

avail, and death occurred within five minutes of the end of the operation.

Each transfusion was from a different donor, who was carefully grouped and whose blood was directly tested with that of the patient before each transfusion was started. Post-mortem examination showed no abnormal changes in the lungs. The heart muscle was pale but otherwise normal, and there was no clot in either branch of the pulmonary artery. The spleen was normal in size. The kidneys were large (right 8 oz., left 11 oz.), and showed a mixture of chronic interstitial nephritis and toxic changes; microscopically there was extensive tubular damage and fibroid glomeruli.

Case II

This was a case of pernicious anaemia in which death followed five hours after the first transfusion. The patient was a male aged 50 years. He was a typical case of pernicious anaemia, with early subacute combined degeneration of the spinal cord. He gave a history of increasing breathlessness on effort, tingling and numbness in the hands and feet, and loss of energy, but no loss of weight.

He was fairly well nourished, but anaemic, and his skin and conjunctivae had a subicteric tinge. His peripheral arteries were moderately sclerotic and his heart sounds were regular, but of only fair quality. His blood pressure was 125/75 mm. Hg. Examination of the chest revealed a mild generalized bronchitis. The only abnormalities found in the central nervous system were slight nystagmus, absent abdominal jerks, bilateral extensor plantar responses, and absent vibration sense in both legs.

His urine contained a trace of albumin, but no excess of urobilin. No occult blood was found in the faeces. A fractional test meal showed a complete achlorhydria, with no delay in emptying of the stomach. His blood Wassermann was negative, Group IV (Moss), and his blood count was: red blood corpuscles, 1,250,000; haemoglobin, 25 per cent.; colour index, 1; average size of corpuscle, 8 μ ; white blood corpuscles, 2,500 per c.mm. A differential count showed: polymorphs, 54 per cent.; small lymphocytes, 42 per cent.; large lymphocytes, 4 per cent.; nucleated red cells present in fair numbers. A urea clearance test gave the following results:

Blood urea at the end of the first hour, 85 mg. per 100 c.cm.
Concentration, 2.1 per cent.
Clearance, 29 c.cm.

That is, there was evidence of very definite renal damage.

He was given 5 c.cm. of campolon daily intramuscularly. It was thought that perhaps his improvement might be hastened by a transfusion. Accordingly, on the fourth day after admission he was given 550 c.cm. of Group IV blood by the indirect (citrate) method. A transfusion service donor who had given blood on several previous occasions was used, and the patient's and donor's blood were perfectly compatible when mixed directly before the transfusion.

There was no immediate reaction of any kind, but one hour later the patient complained of feeling cold and had a rigor. He was given hot drinks, and he settled down again with no further complaints until four hours after the transfusion, when he suddenly collapsed and went into a coma, with cyanosis, dyspnoea, and a feeble pulse. His chest was then full of bubbling rales, and he died an hour later.

At necropsy the lungs were found to be oedematous. The heart was slightly dilated and its muscle was very flabby and showed a typical thrush breast appearance. There was a moderately severe degree of chronic interstitial nephritis. No other abnormalities were found, and no evidence of pulmonary embolism or clot formation in any of the larger vessels was demonstrated.

Classification

While these deaths following transfusion are not common, they are nevertheless disturbing. It would therefore be as well to consider in a general way the groups of conditions in which a transfusion may be thought a reasonable adjunct to other therapy, and at the same time to classify the fatalities in the same way.

The first big group of conditions for which blood transfusion is used universally is in the treatment of acute blood loss from any source. If reference is made to the fatalities it is clearly seen that not one of them falls into this group. Further, I think it is generally agreed that transfusion in these conditions is never followed by any reaction to which death could be directly attributed. Blood transfusion is employed so often in these cases, especially by the obstetrician, that had this not been so I feel sure that we should have found many more cases reported in the literature.

In another large group the object of the transfusion is not to replace actual blood loss, but to stimulate blood formation. The various anaemias, both primary and secondary, obviously fall into this group. Fatalities reported belonging to this group are (a) Plummer's Cases I to IV inclusive; (b) Carrington and Lee's case (1923); (c) both my cases.

A third group of conditions consists of those in which the object is to stimulate the body's reaction to infection or toxæmia, either acute or chronic. Case V in Dr. Plummer's series falls into this group.

Let us now consider the differences between the cases of the first large group and those of the second and third groups. In the first group we have a perfectly normal cardiovascular system deprived of blood. If we replace the blood lost the patient, other factors being equal, is in no worse state than he was before the onset of his hæmorrhage. In the second and third groups we have a cardiovascular system which has been injured, to a greater or less degree, by a deficient oxygen supply over a long period or by toxæmia, or by both in conjunction; here the amount of humoral fluid is at least normal. To this system we add a strain in the shape of an increase in blood volume, and thus place an extra load upon an organ which is already relatively inefficient. Fortunately the cardiac reserve is large enough to meet the demand in most cases, and nothing happens; but in a few it is not, and even death may result.

Possible Methods of Prevention

How, then, may this strain be avoided? One practical suggestion is that before transfusion in these cases venesection should be performed, and only the same volume of blood transfused as has been removed. The only possible alternative is to resort to frequent small transfusions, which are often a nuisance to both patient and doctor, and are naturally of very much less therapeutic value.

In considering certain factors which are common to all three groups, anaphylaxis at once comes to mind. This might suggest that it plays perhaps a smaller part in producing a fatal issue than that which Dr. Plummer would accord it. May I add in conclusion that it is my experience of direct transfusion that by increasing the rate of transfusion, especially after about 150 c.cm. of blood has been given, it is possible to produce in almost any patient a fit of severe coughing, which rapidly ceases if the transfusion is stopped for a moment or two, and does not as a rule return even when the rate of transfusion is increased again. I think this may be ascribed to a temporary over-distension of the right auricle, and until the distension of the auricles is equalized via the pulmonary circulation a certain amount of hold-up on the right side of the heart results. Once the cardiac output has been increased the constant addition of blood does not produce any significant effect upon the action of a normal heart.

Towards the end of a moderately large direct transfusion it is quite common for the patient to complain of backache and to become somewhat restless; whether this is in some way connected with the accommodation of the added blood by the splanchnic circulation would be a difficult point to settle; it is certainly quite different in character from the early severe pain in the back associated with the transfusion of incompatible blood.

Summary

1. Two cases of death resulting directly from transfusion with compatible blood are reported.
2. The importance of recognizing the causes at work in producing possible fatalities is stressed.
3. The mechanical rather than the anaphylactoid element is regarded as being the more important in their causation.
4. It is suggested that a venesection might be performed before transfusion in certain cases.

REFERENCES

- Carrington, G. L., and Lee, W. E. (1923). *Ann. Surg.*, **78**, 1.
Plummer, N. S. (1936). *British Medical Journal*, **2**, 1186.

Clinical Memoranda

Gastric Perforation in a Tabetic

In "acute abdomens" pain is usually the presenting symptom, and its absence may have disastrous results. I have been urged to publish the following case on account of its rarity and interest.

CASE RECORD

I was called a short time ago to see a man, aged 40, previously unknown to me, who complained of vomiting and diarrhoea. The onset had been acute the evening before, following a motor-cycle ride which had resulted in his becoming wet and thoroughly chilled. No article of diet could be incriminated, and there was no history of previous gastric trouble. When seen he was sitting up in bed, quite cheerful, not obviously very ill, and with no abdominal pain. His temperature was 99° F., pulse rate 84. The tongue was slightly furred, the abdomen tender, but not more so than one would have expected in a case of gastro-enteritis of this degree, nor was there more than slight rigidity. The vomitus and faeces had not been kept; the former had consisted at first of food, and later of "green bile"; the faeces were liquid and pale. He complained of pain, increasing in severity, across the back of his neck and shoulders.

I diagnosed gastro-enteritis, aggravated and possibly caused by the chilling, and fibrositis in the region of the trapezii from the same cause. I decided not to give a purgative, but prescribed a bismuth mixture and liniment. The next day his general condition and physical signs were about the same, except that the temperature was 102° F., the pulse rate 96. Vomiting had ceased, but the diarrhoea persisted. The pain in the neck and shoulders was worse. The day after that he appeared for the first time obviously ill. He was collapsed, blue, and sweating, with a temperature of 97° F. and pulse rate 120; there was still no abdominal pain, but the neck and shoulder pain persisted. I had him removed to the Buchanan Hospital, St. Leonards, where he was seen by Mr. J. A. Kerr, F.R.C.S., who confirmed the physical signs and advised continuation of medical treatment. The man's condition grew rapidly worse, however, and he died the following day.

Necropsy showed, to our astonishment, perforation of a large gastric ulcer on the lesser curvature of the stomach, with, of course, generalized peritonitis. In course of time the man's panel record card was forwarded to me. I then found out that (as I had tentatively suggested after the