about her condition. At first irritable, while mentally very alert, with rapid breathing and fits of screaming, she was admitted to hospital twenty-four hours after the ingestion of the aspirin. Her clinical picture, with deep rapid respirations 48 per minute, flushed face, and vomiting, was compatible with a diagnosis of acidosis; yet this was not confirmed by investigation. The alkali reserve was 50 volumes per cent., and the urine gave negative reactions to Gerhardt's and Rothera's tests. The treatment given was 8 ounces (227 c.cm.) of 2 per cent. glucose and saline solution intravenously, and 1 pint (568 c.cm.) of 10 per cent. glucose and 0.9 per cent. sodium bicarbonate solution orally during the first twelve hours. The child's breathing remained deep, and the rate was 40 per minute; she was still irritable. The alkali reserve had fallen to 29 volumes per cent. and the urine gave positive reactions to Gerhardt's and Rothera's tests, showing that a state of acidosis had developed.

Twenty-six ounces (738 c.cm.) of sodium lactate solution and 17 ounces (447 c.cm.) of 10 per cent. glucose and saline solution were given by the slow-drip method intravenously in twenty hours, the patient showing considerable improvement. She was able to take a normal light diet, and appeared quite well after a further three days.

Conclusion

In our cases the weight of evidence seems to be in favour of the intoxication being possibly a toxic action on the respiratory centre, but chiefly due to an acidosis. This is borne out by the rapid improvement in the clinical condition which accompanied the rise in the alkali reserve following the administration of sodium lactate solution. Odin (1932) warns us to be careful in using salicylate in nephritic patients. In two of his cases already quoted albumin was present in the urine before treatment was commenced, and both patients developed symptoms of poisoning with so small a dose as 45 grains a day. All of his cases had a moderate number of red blood cells in the urine, five had albuminuria, and two had some casts. This, considered with the necropsy findings of acute nephritis, indicates that it is difficult for nephritic patients to excrete salicylic acid.

Summary

1. A survey of the reported cases of aspirin poisoning is given, showing that large doses are necessary in adults to produce toxic effects.

2. Alkalis increase the absorbability of aspirin.

3. Two infants, who developed severe toxic symptoms after the ingestion of 30 grains (1.944 grammes) and 60 grains (3.89 grammes) of aspirin respectively, are described; both recovered after sodium lactate (Hartmann's solution) had been injected intravenously.

4. It is considered that the toxicity of aspirin is due to an acidosis, and possibly to a direct action of salicylic acid on the respiratory centre.

ACKNOWLEDGEMENTS

We wish to thank Dr. Stewart Ferguson for permission to publish the reports of these patients.

We are greatly indebted to Miss Splatt of the Walter and Eliza Hall Institute, Melbourne, whose estimations of the alkali reserve served as a valuable guide to treatment; to Dr. Bolton, medical superintendent of the Royal Melbourne Hospital, who supplied us with the sodium lactate solution; and to Dr. Ian Wood for his advice and assistance.

REFERENCES

- Balázs, J. (1930). *Med. Klinik*, **26**, 1664.
 Dyke, S. C. (1935). *Lancet*, **2**, 613.
 Hartmann, A. F. (1935). *Arch. int. Med.*, **56**, 413.
 Martindale and Westcott (1928). *Extra Pharmacopoeia*, 19th ed., 1, 75.
 Ibid., 19th ed., 1929, **2**, 15f.
 Neale, A. V. (1936). *British Medical Journal*, **1**, 109.
 Odin, Martin (1932). *Acta med. Scand.*, Supp. **50**, 177.
 Paiseau, G., Friedmann, E., and Vaille, C. (1934). *Bull. Mém. Soc. méd. Hôp. Paris*, **50**, 1201.
 Quincke, H. (1882). *Berl. klin. Wschr.*, **19**, 709.
 Wyllie, A. M. (1935). *Lancet*, **2**, 768.

CONTINUOUS INTRAVENOUS SALINE INFUSION

BY

HAMILTON BAILEY, F.R.C.S.

Surgeon, Royal Northern Hospital

WILFRED I. B. STRINGER, M.D., M.C.P.S.

Resident Medical Officer, Royal Northern Hospital

AND

KENNETH D. KEELE, M.D., M.R.C.P.

Casualty Physician, St. Bartholomew's Hospital;
 Late Medical Registrar, Royal Northern Hospital

Continuous intravenous saline infusion is now being used more extensively in this country; unquestionably it has helped to save many lives. We feel that a time has been reached when it would be well to review its advantages, to try to enumerate its indications and contraindications, and particularly to dwell upon the dangers connected with its use.

Indications

1. *Dehydration* from any cause, such as starvation, vomiting, diarrhoea, or excessive sweating.—In acute intestinal obstruction the combination of continuous gastric aspiration and continuous intravenous saline infusion both before and after operation is a marvellous therapeutic advance.

2. *Prevention of Post-operative Shock*.—A typical indication is at the close of the operation of partial gastrectomy, especially when it has been performed on a poor subject.

3. *Shock*.—Here we appear to have a helpful method of combating shock. In these cases we have added one ampoule of coramine to each pint of saline solution with advantage. French observers have reported favourably on the results of adding adrenaline to the solution in doses of about 5 minims of 1 in 1,000 solution to each pint.

4. *Diabetic Coma*.—This is probably the best method of introducing large quantities of glucose and insulin, a measure which is so necessary in this condition.

5. *Post-operative Thyrotoxicosis*.—It has long been recognized that a copious fluid intake is beneficial in these cases. If administered early enough continuous intravenous saline infusion will often tide the patient over a crisis. If hyperpyrexia and other signs indicate that the patient is being flooded with the toxin, the injection of 50 minims or more of Lugol's solution into the delivery tube, associated with the application of an ice-pack to the limbs and precordial region, and, above all, the employment of an oxygen tent, is the best possible course to adopt in these anxious, but now fortunately rare, catastrophes.

6. *In profound jaundice, after the obstruction has been removed*, it is often remarkable how quickly the jaundice clears when continuous intravenous saline treatment is administered (see also Contraindications).

7. *Septicaemia*.—That benefit accrues from continuous intravenous medication in septicaemia is not generally agreed. In our experience in some cases continuous intravenous saline therapy, by diluting the circulating toxin and stimulating the excretory activity of the kidneys, appears to have done good. The addition of a capsule of intravenous mercurochrome to each filling of the reservoir definitely has done no harm, and we believe that in several cases of septicaemia it has helped the patient to combat the infection.

8. *As a Convenient Means of Inducing Evipan Anaesthesia.*—In cases of perforated peptic ulcer we give the patient continuous intravenous saline infusion while the theatre is being made ready. With the saline solution still dripping into the vein the patient is brought to the theatre. The abdomen is opened under local anaesthesia. As soon as the peritoneum has been reached a small dose of evipan is introduced into the rubber tubing of the saline apparatus, and evipan flows into the vein together with the saline solution, which has been turned to a steady flow. The anaesthesia produced by the evipan gives ample time for the performance of the operation. The infusion before and during the operation mitigates shock, and the absence of a general anaesthetic helps, in a large measure, to prevent a major complication of treated perforated peptic ulcer—pulmonary involvement.

9. *Continuous Intravenous Sodium Sulphate in Oliguria and Anuria.*—Providing that in obstructive cases the cause of the anuria has been removed, continuous intravenous saline, or, what would be better still, continuous intravenous sodium sulphate, are diuretics without a peer. I. L. Dick found that in every case the diuresis produced by intravenous sodium sulphate was greater than that produced by sodium chloride, and in no case was there any evidence of storage of sodium sulphate in the body. An isotonic solution is made by dissolving 42.85 grammes of Glauber's salt in one litre of water. We have used this solution in a variety of cases of anuria, oliguria, and "latent" uraemia. Needless to say, in cases where there is a mechanical obstruction to the outflow of urine this must be first removed. Some of the patients were actually moribund. In only one instance did it fail to produce diuresis, and that was in a case of polycystic disease of the kidneys which at the post-mortem examination showed no evidence of any renal tissue remaining. We generally give only one or one and a half pints of this solution daily; during the rest of the twenty-four hours normal saline solution is employed. There is no reason for limiting the dosage to this amount other than timidity on our part.

Contraindications

1. *The Failing Heart.*—A history of dyspnoea on exertion, uncompensated valvular disease, or any possibility of cardiac weakness should call for hesitation in increasing the bulk of circulating fluid. When the integrity of the heart muscle is suspected a physician should be consulted before employing continuous intravenous saline infusion.

2. *Pulmonary Congestion.*—In all conditions where there are signs of oedema or consolidation of the bases of the lungs the method is inadvisable.

3. *Hypertension.*—If the blood pressure is high it is obviously courting danger to burden the circulation still further. Essential hypertension and the sthenic form of uraemia with convulsions are especially mentioned as contraindications, for in these cases some degree of cerebral oedema may be present.

4. *Bright's disease* in all its stages is a grave contraindication. In advanced uraemia, where Bright's disease is a possible underlying cause, venesection, one pint, followed by continuous intravenous sodium sulphate, one to one and a half pints, may be tried. The same precaution of preliminary venesection and cautious replacement of circulating fluid should be taken in all cases of chronic bilateral pyelonephritis. Polycystic disease is a definite contraindication.

5. *In Obstructive Jaundice where the Obstruction has not been Removed Completely.*—Death is hastened by this measure, as was observed in one of our patients where

the obstruction was due to carcinoma of the common hepatic duct. At the post-mortem examination the kidneys were deeply stained and choked with bile.

Preparation of the Solution

Obviously a solution which is introduced directly into the circulation must be absolutely sterile, but this is not enough. It must also be free from dead bacteria and other extraneous matter such as fur from the sterilizer. Boiled tap-water does not fulfil these requirements, and if it is used reactions are bound to occur. Triply distilled water must be employed in all cases. Although the Pharmaceutical Society is less stringent in its regulations we have made it a rule that distilled water more than a week old is discarded. With a view to this flasks are stamped with the date of preparation.

It is, of course, equally important to have the sodium chloride sterile and pure. Tablets sold in tubes for the preparation of normal saline are not prepared with a view to intravenous therapy. Sterile ampoules for making a 5 per cent. solution of glucose are readily obtainable; it would be an advantage to have similar ampoules of sodium chloride.

In fully equipped institutions the whole of the work of preparing the saline solution can be undertaken by the dispensary staff. At the Royal Northern Hospital flasks of sterile saline solution prepared by the dispenser with newly triply distilled water are supplied on demand to the wards. Periodical bacterial examinations are made in the pathological laboratory.

Saline or Saline and Glucose?—Normal saline solution alone meets most requirements. It has two outstanding advantages over glucose or combined saline and glucose. First, it causes less chemical phlebitis and thrombosis, and secondly, it is manifestly physiological. When glucose is employed we are introducing a substance into the circulation which requires katabolism, and in our opinion if glucose is to be used there should be adequate reasons for its introduction; the patient must be in need of glucose, as one who has been starved, and only sufficient glucose for his immediate needs should be introduced by the intravenous route. A 5 per cent. solution of glucose in six pints of saline contains about 180 grammes of glucose, so at the usual rate of flow the patient is receiving 720 calories in the twenty-four hours. This may be increased if no other source of nourishment can be utilized by the patient. As regards insulin, if there is no reason to think that there is a deficiency in this hormone there is no reason to give more.

Technique

Inserting a Needle or Cannula.—It is not always necessary to cut down upon a vein; a small nick in the skin and the insertion of a special hollow needle (Fig. 1)

FIG. 1.—A, Needle for inserting into a vein by a puncture. Once it is in place satisfactorily it is kept at rest by a strip of adhesive plaster across each flange. B, Cannula for use when vein is too inconspicuous for a needle to be inserted. This is made in two sizes, for adults and children.

is equally effective in many instances. A vein in the forearm is usually chosen. It is a great mistake to choose the most prominent vein. One which is easily palpable but well obscured and supported by subcutaneous fat will

be found to be less mobile, and thus more readily entered. The large prominent veins of the aged are notoriously difficult to enter, and in these cases it is as well to transfix the vein transversely with a fine needle before carrying out the puncture. When the veins are small and difficult to locate, if time permits it is useful to soak the arm in a hot bath for fifteen minutes before attempting the puncture. When it is necessary to cut down upon a vein a small transverse incision is used. If no veins can be seen an incision should be made over the commencement of the internal saphenous vein. Even in a new-born infant we have never failed to find this just anterior to the internal malleolus. There is no need to dissect out a vein with the point of a scalpel; blunt longitudinal dissection by opening the jaws of a small haemostat (Fig. 2) is more efficient and saves time.

FIG. 2.—A rapid and efficient method of displacing a subcutaneous vein through a small transverse cutaneous incision.

Splinting the Arm.—A posterior plaster strip well padded with gamgee with a turn around the wrist forms an ideal splint readily adapted to each individual patient.

Splinting the Leg.—There is no better method than Thomas's knee splint, as described in a previous article.

The Temperature of the Fluid

In spite of the many ingenious devices which have been invented for keeping the fluid warm we are still of the opinion that the simplest is the best—namely, the patient's leg or arm into which the saline solution is flowing should be kept warm with a hot-water bottle or a Q-pad. In addition to simplicity this leaves the reservoir and the delivery tube in full view.

The Administration

The actual administration is largely in the hands of the nursing staff. It is of paramount importance to be sure that the nurse appreciates the simple yet vital responsibilities connected with the care of the apparatus and the maintenance of strict asepsis. The nursing instructions are really very simple, and can be carried out by any conscientious State-registered nurse, providing that if she has not been trained in the method she receives special instructions. With the help of a competent nurse there is no reason why continuous intravenous saline solution should not be administered to a patient in a private house, as contrasted with a fully equipped hospital.

Nursing Instructions

1. Every portion of the apparatus must be freshly boiled and washed through with sterile water or saline solution before the reservoir is filled. New rubber tubing

should be boiled, stretched, and washed through to remove chemicals before the apparatus is assembled.

2. Every utensil used in the transfer of the saline to the reservoir must be absolutely sterile (Fig. 3).

3. To prevent the entry of bacteria from the air the top of the reservoir must be covered with sterile gauze.

4. The reservoir must always be kept more than half full of saline solution.

5. The patient's limb into which the solution is flowing should be kept warm with a hot-water bottle or Q-pad.

FIG. 3.—Replenishing the reservoir.

6. The average rate of flow is fifty drops per minute. This should be timed at frequent intervals. Fifty drops per minute is about a quarter of a pint per hour—six pints in twenty-four hours.

7. If the flow stops: (a) do not pinch the tubing; (b) see if the tubing is kinked; (c) if the flow is not restarted by some simple adjustment of the limb or the tubing report the matter at once.

8. Watch for and report immediately: (a) rigors; (b) redness along the vein; (c) oedema of the feet, face, or arms; and (d) any sign of respiratory distress.

9. Measure the amount of urine the patient has passed. If the output is less than the intake report the matter. Also report if the specific gravity becomes low.

Special Instructions for the Medical Officer

1. If the patient does not appear to be responding take the blood pressure. Intravenous saline infusion should not be continued in the presence of a falling blood pressure.

2. Especially after forty-eight hours watch the bases of the lung for signs of cedema.

3. Watch the urinary output, paying particular attention to the specific gravity.

4. Unless there are definite indications to the contrary there is seldom need to continue the flow after the third day. A good working rule is to remove the needle or cannula at the end of seventy-two hours and if necessary to reinsert it into another vein after six or twelve hours.

5. If the flow stops do not dissemble any part of the apparatus without due thought of the possibility of air embolism. Rather remove the needle or cannula, and if necessary reinsert it into another vein.

6. In some cases, especially when glucose is employed, redness and tenderness appears along the course of the vein. This is hardly ever anything more than a chemical phlebitis, which soon settles down after the flow into that vein has been discontinued.

Conclusion

A patient who is ill enough to warrant this form of therapy needs watchful care and attention. The continuous administration of saline solution intravenously does not therefore add materially to the nursing or medical attention which such a patient requires.

The advantages of this over other methods of administering saline solution are great; the rate of flow and the amount of fluid given can be regulated with mathematical precision. Like all good things the principal evil lies in its abuse, and the most obvious danger is oedema. It should be realized that it is not until about eight pints of fluid have collected in the tissues, a quantity equal to the volume of the circulating blood itself, that the sign of pitting on pressure becomes manifest. It would be a very great help if there existed some simple test which would indicate when the patient had received sufficient fluid and that more would be harmful. At the present time the only choice lies between an estimation of the plasma protein, which is far too complicated and time-consuming for general application, and the rough-and-ready method of charting the output of urine and its specific gravity in one column and the intake of all fluid in the other (see table). Providing that reasonable care

*Continuous Intravenous Saline Balance Sheet**

24 hours ending	Intake	Output
	Pints	Pints
	At 50 drops per minute	Urine { S.G. a.m. } { S.G. p.m. }
	„ 30 drops per minute	Vomit
	„ drops per minute	Normally { Faeces }
	By mouth	1½ pints { Sweating and lungs, say
	Total	Total

* Pads of these charts can be obtained from the G.U. Manufacturing Co., 28A, Devonshire Street.

is exercised the latter is enough for practical purposes. A balance sheet of fluid loss and intake should be drawn up. For loss of fluid by sweat, lungs, and normal defaecation a total of one and three-quarter pints has to be calculated. To this is added the urine output plus vomitus if necessary. From this total the maintenance dose for the next twenty-four hours can be calculated.

REFERENCES

- Bailey, H., and Carnow, J. M. (1934). *British Medical Journal*, 1, 11.
 Baudouin, A., Lewin, J., Bernard, H., and Sallet, J. (1935). *Paris méd.*, 1, Suppl., 569.
 Dick, I. L. (1934). *Edinb. med. J.*, 41 (Med.-Chir. Soc. Trans.), 203.
 Smith, G. G. (1935). *Canad. med. Ass. J.*, 33, 262.
 Walther, W. W. (1937). *Lancet*, 1, 6.

CLIMATO-PHYSIOLOGICAL INVESTIGATIONS AT THE SEASHORE*

BY

OTTO KESTNER, M.D.

For nearly two hundred years convalescent children have been sent to the seaside, and it is generally known that this treatment is successful. When, as a physiologist, I began to study the curative power of the seashore I remembered that pharmacology had greatly increased the efficiency of drugs by modern methods of experimental study. In recent years many investigations have been made upon children at the small seaside town of Wyk on the island of Föehr, on the German side of the North Sea, where there are two large children's homes belonging to Berlin and Hamburg, and a great number of smaller homes for convalescent children under national insurance. The town built a small laboratory for us and the insurance organizations made a grant of some money. Dr. Häberlin, who has practised in Wyk for more than thirty years, and other local medical men, were associated with me in this investigation.

Physiological Effects of Climate

Medical climatology makes a true distinction between climates that are stimulating or "bracing" and those that are sheltered or "sedative." But a stimulating climate, so called, includes both the stimulus of the solar heat, which we do not feel so bracing, and the ultra-violet rays of the sun, which we do not feel at all. Obviously climatic stimuli of all kinds—of the sunshine, wind, cold air, or rolling breaking waves—operate upon the body through the skin, and it is true also that it is by means of the skin that the human body controls its temperature-regulation by changing the size of the capillaries at different external temperatures.

The observations which formed the basis of our work may be summarized as follows:

1. There is a very rapid reaction of the capillaries (Häberlin, Kestner, *et al.*, 1923), both dilating and contracting, in healthy children living at the seaside.

2. In pale, weakly, and undernourished children, damaged by town life and sent by health officers or doctors to the seashore, the same reactions occur, but they are slower and weaker. We found that these children, not protected by immediate reactions, lose many calories, and that the temperature of the body falls considerably after the shortest sea-baths, even in August, and even whilst merely playing on the seashore. Such children cannot at first tolerate sea-bathing, but during the cure the capillaries gradually recover their normal "gymnastic" function. The circulatory reactions improve week by week, and the same children who felt "terribly cold" and cried at their first bath could play later on in the open air without clothing, even in stormy weather. The time and completeness, or otherwise, of the reaction can be determined and measured, and these measurements are very helpful in checking the degree of exposure and the recovery of health. This behaviour of the cutaneous circulation explains an observation of Curschmann. He found that healthy young men, mostly students, had the same systolic blood pressure after sea-bathing as before, but that in patients with contracted blood vessels (arterial hypertonus) the blood pressure was lowered after a sea-bath. For this reason he called the seashore "the paradise of the hypertonic."

* Abstract of a lecture given to the Kensington Branch of the British Medical Association, October 16, 1936.

† The reference to Dr. Carl Häberlin's address, reported in the *British Medical Journal*, August 22, 1931 (p. 331).