R. G. Gordon (1936) in a recent paper drew attention to the metabolic factor in chronic rheumatism, with special reference to fibrositis, and observed that the pendulum had swung too far, and that we must consider some cases of chronic rheumatism to have a gouty basis, quite apart from the question of the presence of a focus of infection. This same viewpoint I raised at the discussion on chronic rheumatic diseases at the Oxford meeting of the British Medical Association last year. Further, I pointed out that there were cases where there might be no evidence of an infective focus. For instance, patients tell you that they get pains and aches in their muscles (a sprained feeling) after drinking beer or Burgundy; after eating strawberries, asparagus, etc.; or after taking a purge; and patients suffering from glycosuria often complain of such symptoms. Another group of patients will tell you that when they get out of bed in the morning their feet feel tender, their hands are stiff, and their grip is weak. By the time they have completed dressing this discomfort has passed off. On examination there may be no physical signs beyond perhaps some tenderness on squeezing the metacarpophalangeal joints as in a vigorous handshake. Such cases improve much by the use of sulphated or sulphide waters, and by trying to correct the errors of metabolism, quite apart from general physical treatment.

Post-malarial Cachexia

The waters of these spas are also distinctly useful in the cachexia following malaria. Enlargement of spleen and liver derives much benefit from a course of these waters, combined with baths, exercise, change of air, etc. In the winter and early spring Cheltenham and Leamington are more suitable, while in the summer months the more bracing climate of Harrogate and Strathpeffer works wonders. In fact, I know of no type of case which shows greater and more rapid improvement with a course of these waters. Many residents in the Tropics, from India, Malay, West Coast of Africa, etc., make it a practice to have a "cure" during their periodic leaves home, mainly as a prophylactic measure.

REFERENCES

- Bain, William (1935). *The Pharmacological Action of the Harrogate Drinking Waters*. Churchill, London.
Brown, D., and Woodmansey, A. (1929). *Proc. roy. Soc. Med.*, 1928-9, 22, 487.
Gordon, R. G. (1936). *British Medical Journal*, 2, 1243.
Lockie, L. M., and Hubbard, R. S. (1935). *J. Amer. med. Ass.*, 104, 2072.
Rehfuß, M. E., and Nelson, G. M. (1935). *Medical Treatment of Gall-bladder Disease*. Saunders, Philadelphia.
Tidy, H. Letheby (1936). *British Medical Journal*, 2, 418.
Wilkie, D. P. D. (1934). *Ibid.*, 1, 909.

A comprehensive survey of obesity and emaciation due to pathological causes, by Professor Francesco Galdi of Pisa and three of his colleagues, is published in the proceedings of the forty-second congress of La Società Italiana di Medicina Interna, held in Rome in October, 1936. The authors comment on the extensive literature, especially from German and American sources, dealing with obesity, and the comparative neglect of pathological loss of flesh. The relation of metabolism, of the nervous system, and of the endocrine glands to these conditions is fully discussed, and some sixty pages are devoted to more detailed separate analyses of the metabolism of proteins, carbohydrates, and fats. Following a description of the pathological anatomy of both conditions, the several forms of obesity and emaciation are considered in their clinical aspects.

POST-SCARLATINAL NEPHRITIS A STUDY IN PREVENTION

BY

B. A. PETERS, M.D.Cantab., D.P.H.

*Medical Superintendent, Ham Green Hospital, Bristol
Lecturer in Infectious Diseases, University of Bristol*

AND

IRIS M. CULLUM, M.D.Lond.

Late Assistant Resident Medical Officer, Ham Green Hospital, Bristol

By giving large amounts of alkalis in the course of an attack of scarlet fever Carter and Osman (1927) showed that a considerable reduction in the incidence of nephritis occurred in cases so treated compared with a control series. Peters (1932) brought forward evidence showing that the administration of thyroid and iodine for the first fortnight of an attack of scarlet fever was followed by a considerable fall in the incidence of nephritis.

Below is given a table showing the incidence of post-scarlatinal nephritis in five-year periods during the past twenty-five years in Ham Green Hospital.

TABLE I

Period	Total Cases of Scarlet Fever	Cases of Nephritis	Percentage Incidence of Nephritis
1910-14	2,771	53	1.92
1915-19	1,329	25	1.88
1920-24	2,496	63	2.5
1925-29	3,148	28	0.88
1930-34	2,427	9	0.37
1935-36	1,350	43	3.2

Thyroid and iodine were given during the period of 1930-4. Comparing the different incidence in the two periods of lowest prevalence, it will be found that the difference is 2.3 times the standard deviation. The difference is very much higher if the whole twenty-five-year period is taken into consideration. On these grounds we were disposed to think that our method was proving effective, especially as 1930-4 figures include two cases which were admitted with nephritis already present in whom preventive methods had not been applied.

In the winter of 1934-5 a more severe type of scarlet fever appeared. It was characterized by a great increase of secondary attacks whilst in hospital; by several second attacks in patients within a few weeks of discharge; by the increase of our return case rate to three times its normal number; by an increase of all the more severe complications; and by the occurrence of attacks of scarlet fever in six Dick-negative reactors among the staff, an event which had not occurred for many years.

In spite of administering thyroid and iodine, cases of nephritis cropped up with distressing frequency. We therefore investigated our method afresh, using alternate cases as test and control. Table II shows the result of this investigation. By the term "albuminuria" is meant symptomless albuminuria for three consecutive days or longer. "Nephritis" includes cases with signs or symptoms of renal failure. The observation of the sedimentation rate of the red blood corpuscles by Cookson (1936), working in this hospital, suggests that there is a sharp line of demarcation between the two conditions.

TABLE II.—*Series of Scarlet Fever Cases Treated with Thyroid and Iodine and Controls*

		Albuminuria	Nephritis
Test cases	 165	34 (21%)	11 (6.6%)
Controls	 162	26 (16%)	9 (5.5%)

It was evident that our method was having no beneficial effect in this type of scarlet fever. We therefore submitted another series of alternate cases and controls to Osman's method, giving the alkaline mixture to the test cases in the doses he recommended continuously throughout the period of their stay in hospital, while the controls received alkalis for the first two days of their stay in hospital only. The results are shown in Table III.

TABLE III.—*Series of Scarlet Fever Cases Treated with Alkalis (Osman's Method) and Controls*

		Albuminuria	Nephritis
Test cases	 124	15 (12.1%)	5 (4.3%)
Controls	 134	25 (18.6%)	6 (4.1%)

In this series it was also evident that no statistically significant improvement was being effected by Osman's method. On referring to Table I it will be seen that the incidence of nephritis was higher during 1935-6 than during any other period in the last twenty-seven years in Bristol.

Conclusion

It would appear that the various strains of streptococci causing the symptoms of scarlet fever vary greatly in their power of producing renal complications, and that methods which seem to be effective in a controlled series of cases in preventing the complication of nephritis caused by certain strains may fail completely against other strains. A reliable means of preventing this serious complication which is effective against all strains of streptococci is still to be found.

REFERENCES

- Carter, H., and Osman, A. A. (1927). *Proc. roy. Soc. Med.*, **20**, 1405.
 Cookson, J. S. (1936). *Brit. J. Child. Dis.*, **33**, 251.
 Peters, B. A. (1932). *Practitioner*, **129**, 614.

R. B. Engelstad (*J. Radiol. Electrol.*, February, 1937, p. 53) reports the results of radium therapy in fifty-six cases of cancer of the buccal cavity. These included cancer of tongue, of the mouth proper, and cancer of the gums. The treatment consisted of teleradium therapy alone, or teleradium therapy plus radium implantation plus electro-coagulation, or of teleradium therapy plus surgical operation. The cervical metastases were treated either by teleradium therapy or by teleradium therapy plus operation. The source of radiation for the teleradium therapy consisted of a radium unit containing 2,600 mg. of radium element. The distance between the source of radium and the skin varied between 5.7 and 12.5 cm. Several fields were irradiated depending on the size and situation of the tumour. One field was treated each time and the dose varied between 3 and 6 Dominici units ($1\frac{1}{2}$ to 2 hours), and a total of 50 Dominici units was applied. Skin and other reactions all cleared up in two to three weeks. The results of the treatment of the fifty-six cases were as follows: $1\frac{1}{2}$ to 4 years' cure, twenty-nine patients (53 per cent.); cure after recurrence, three patients; cured for the time being, thirty-two cases; alive but not cured, four cases; dead, twenty cases (36 per cent.).

CARCINOMA OF BREAST WITH WIDE-SPREAD METASTASES TWO CASES OF RECOVERY

BY

W. BARRINGTON PROWSE, M.R.C.S., L.R.C.P.

Honorary Consulting Radiologist, Royal Sussex
County Hospital

I venture to report these two cases in view of the extremely satisfactory results which followed treatment. Both were cases of mammary carcinoma in which wide-spread metastases had occurred, and both patients are now, between one and two years after cessation of treatment, in good health and free from any sign of carcinoma. One of them had had no operation, as she did not present herself to a doctor until the disease was too far advanced for surgery; the other had had a radical amputation of the breast eighteen months before the presence of secondaries was discovered and treatment begun.

Case I

An unmarried lady of middle age was sent to me in August, 1934. There was in her right breast, adherent to the skin, a carcinomatous lump, which was broken and discharging slightly. There were secondary glands in the axilla and above the clavicle, and also secondary deposits in several bones of the thorax and shoulder-girdle. Her general condition was very poor. I thought she was not fit for intensive x-ray therapy, and suggested that she should submit to a course of treatment by Todd's method. She agreed to this, and after some delay treatment was begun early in November. At that time she was entirely confined to bed, and had to be brought by ambulance and stretcher for all x-ray treatment. She was wasted and had much pain, for which morphine had been necessary. She took food badly and was often sick.

On beginning treatment all morphine was stopped. She was given a weekly intravenous injection of compound colloid of selenium and sulphur, followed two days later by a mild dose of high-voltage x rays; a dressing of Paré's ointment (lead amalgam) was applied to the breast. It was found impossible to give her the calcium mixture recommended by Todd; she could not tolerate it, and she only took radiostoleum and thyroid extract intermittently. Her veins were very small and there was considerable difficulty with the injections, which led to some delays and irregularity in the course of the treatment. Eventually the injections had to be stopped altogether—sooner than would have been the case under more favourable circumstances. The breast began slowly to improve not very long after starting treatment, but otherwise there was no change, except some relief from pain.

In the early part of 1935 there was complaint of fresh pain in the lower part of the back and pelvic region, and radiographs revealed extensive secondary deposits in the spine and pelvic bones. The outlook appeared very gloomy, but nevertheless treatment was continued. The one bright spot was the condition of the breast, which by this time had vastly improved. I must pay tribute to the patient's perseverance and great tenacity of purpose. General improvement became evident at last in April, 1935. Pain was much less and the sickness stopped; the appetite improved, and before long the patient was eating more than she had done for years. In the middle of May she could sit up out of bed, and on June 1 was just able to take a walk in the garden. No treatment was given between the end of July and the middle of October.