to many different environmental situations and bodily states, whether emotional, physical, or a mixture of the two; and to assume that in no two cases is it due to an identical combination of factors. At all events it is quite possible to persuade most of the children to change the habit by simple encouragement or by other suggestive means; nor is the task very laborious.

Summary

1. A series of seventy cases of enuresis in mentally normal children is described, with the procedure by which they were treated.

2. The results of treatment are analysed and show at least 60 per cent. of cures maintained over a length of time by the one method, and additional cures by other methods to suit the individual case.

3. Theories of aetiology and different methods of treatment are briefly discussed.

4. Most of the cases may be cured by simple procedures based on encouragement or suggestion.

My thanks are due to Dr. H. C. Cameron, in charge of the children's department, Guy's Hospital, and to Dr. J. E. Roberts, school medical officer for the City of Portsmouth, for permission to include in this series cases collected from their departments.

BIBLIOGRAPHY

- Addis, R. S. (1936). *Proc. roy. Soc. Med.*, **29**, 1515.
 Batty, R. J. (1933). *Enuresis*, London.
 Friedell, A. (1927). *Amer. J. Dis. Child.*, **33**, 717.
 Goldman, L. M., and Malavazos, A. (1936). *Urol. cutan. Rev.*, **30**, 729.
 Hamill, R. C. (1929). *J. Amer. med. Ass.*, **93**, 254.
 Stocks, A. V. (1936). *Med. Officer*, **56**, 87.
 Usher, S. J. (1931). *Canad. med. Ass. J.*, **24**, 665.

ACUTE LYMPHOCYTIC MENINGITIS

BY

WILLIAM HUGHES, M.D., M.R.C.P.

Acting Physician, General Hospital, Singapore

Acute lymphocytic meningitis is a specific disease which has been systematically described only in recent years. It is characterized by symptoms of meningitis, often preceded by upper respiratory tract infection, certain distinctive findings in the cerebro-spinal fluid, and a benign course usually ending in complete recovery.

The modern study of the aetiology of the disease began when Armstrong and Lillie (1934), during the St. Louis epidemic of encephalitis in 1933, isolated from a fatal case in a negro a virus which was pathogenic for monkeys and mice. They found that this virus differed in many respects from the virus of encephalitis on which they were working at the time. Later Armstrong and Dickens (1935) showed this virus to be identical with one recovered from a series of cases of acute lymphocytic meningitis. Bengtson and Wooley (1936) reported successful cultivation of the virus on the chorio-allantoic membrane and brain of the developing chick. They make the interesting observation that inoculated chicks hatched out were definitely subnormal in many features. Findlay, Alcock, and Stern (1936) isolated a virus from a mouse which had developed symptoms of nervous disease. This virus they showed to be identical with Armstrong's virus, and they further established the identity of the American lymphocytic meningitis with the European type. They were able to demonstrate the presence of antibodies in the blood of a Dublin patient previously reported by Collis (1936), and they give a full account of their studies on two cases in England which they investigated themselves.

They also found antibodies in the blood of a certain proportion of wild mice, which suggests that these rodents may form the natural reservoir of the infection. The present position, then, is that lymphocytic meningitis is known to be widely distributed in Europe and North America. Recently Allen and Spencer (1935) have reported six cases from New Zealand. I cannot find any record of cases reported from the Tropics, but I propose to describe later on a series of patients who showed all the symptoms of lymphocytic meningitis. This makes it probable that the distribution of the disease is universal.

Pathology

Armstrong and Lillie found that intracerebral inoculation of the virus was usually fatal for mice. The mice die in convulsions after six to eight days. In monkeys the condition is characterized by drowsiness and apathy, and recovery is the rule. Findlay, Alcock, and Stern found an intense basal meningitis in their experimental animals. The infiltration extended on to the posterior surface of the upper part of the cord. There was extensive round-celled infiltration in the lining membrane of the whole ventricular system. Armstrong and Lillie found oedema and infiltration of the choroid plexus in two-thirds of their specimens. The name "chorio-meningitis," which is often used to describe the disease, is based on this finding. The lesions are confined for the most part to the meninges and choroid plexus, and there is very little perivascular infiltration in the brain tissue.

Symptoms

The clinical symptoms of the disease are hardly sufficient to distinguish it from other forms of meningitis and meningismus. It begins often as a "cold," sometimes with coryza. Even in the early stages there may be generalized aching. Headache can be intense. One of our cases had a peculiar combination of drowsiness and insomnia. The temperature is raised, usually running between 100° and 103° F. Kernig's sign is present in practically all cases.

Armstrong and Dickens say there is usually no paralysis and "no definite neurological signs." Allen and Spencer, however, found an occasional extensor plantar reflex in their cases. In one of our cases there were extensor plantar reflexes, weakness of the legs, and retention of urine. Findlay, Alcock, and Stern record paraplegia, paraesthesia, and cranial nerve palsy in their cases, both of which had retention of urine. The cerebro-spinal fluid shows characteristic changes. The cells are increased, and findings of 50 to 2,000 per c.mm. are recorded (Armstrong and Dickens). Lymphocytes may constitute up to 100 per cent. of the cells, and are rarely below 90 per cent. Sugar, chlorides, and urea are normal. No bacteria are found.

Diagnosis

In the Tropics there are special difficulties about the diagnosis of such cases. Apart from diseases like dengue, influenza, and spirochaetosis icterohaemorrhagica, which may simulate the condition clinically, there are three comparatively common diseases which give similar cerebro-spinal fluid findings. I refer to tuberculous meningitis, syphilitic meningitis, and lead encephalopathy. The important though not the only factors in excluding tubercle are the absence of bacilli, the normal chloride content of the fluid, and the benign outcome of the disease. In the case of syphilis we sometimes have to rely on the Wassermann or other serological reaction. How far these tests are reliable in an individual case it is often

difficult to say. One would expect, however, that the Wassermann or Kahn test would be positive in the spinal fluid in 100 per cent. of cases of syphilitic meningitis. Even so, one reads of fairly well-authenticated cases of recovery in tuberculous meningitis on the one hand and undoubted cases with normal chlorides and absence of tubercle bacilli on the other hand. In syphilis there can be a "clinical silence," as Craps (1936) puts it, during bismuth treatment. It is a legitimate speculation that meningitis could occur before the Wassermann reaction became positive again.

Lead encephalopathy can give a picture not unlike that described for acute lymphocytic meningitis, and the cerebro-spinal fluid findings are similar, except that the cells are rarely increased to the same extent. In children lead line and paralysis are not to be expected and gross anaemia is rare. Encephalopathy in children and adults due to the use of toilet and face powders containing lead is common, especially among Chinese in the Far East.

Keeping the above considerations in mind, I think it is still possible to report with confidence the occurrence of two cases of lymphocytic meningitis in Malaya. Both were treated in the wards of the General Hospital, Singapore.

Case I

A male European, aged 23, admitted on January 11, 1937, was discharged on January 26. His illness began eleven days before admission with fever, cough, and generalized aching. He recovered sufficiently to get up from his bed on the fifth day. On the seventh day the fever returned. He now complained of headache, weakness of the legs, and subjective disturbances of sensation—"pins and needles." He was put to bed and had to be catheterized on the tenth day. On this day he was said to have coughed up some blood or sputum streaked with blood. On admission to hospital (eleventh day) he looked ill. Prominent symptoms were headache and photophobia, and drowsiness with insomnia. The bladder was distended and he was unable to pass urine. Kernig's sign and head retraction were present. Knee-jerks were exaggerated, and there was bilateral ankle clonus. The right plantar reflex was "extensor," the left normal. Abdominal reflexes were absent. There was no ocular paralysis, and the fundi were normal. The lower limbs were weak, but there was no gross paralysis and no objective sensory loss. No signs of disease were observed in the chest. Routine examination of blood (for malaria) and stools (for ova), and repeated examination of the sputum (for acid-fast bacilli) revealed no abnormality.

Lumbar puncture on January 12 revealed a clear fluid under increased pressure. The count gave 110 cells per c.mm.; globulin was present. The total protein amounted to 120 mg. and the chlorides to 710 mg. per c.cm. Sugar was found. A differential count showed that 85 per cent. of the cells were lymphocytes. There were no organisms in stained or unstained smears. The Wassermann and Kahn tests were negative. The lumbar puncture relieved the headache. On the fifth day after admission the patient was able to pass urine, and on the sixth day his reflexes were all normal. His temperature, which was 101° F. on admission, fell by lysis to normal on the fifth day, and remained so. On January 21 he developed a small swelling at the angle of the mouth. A boil formed, which discharged itself on the inner surface of the lower lip after two days. Otherwise his convalescence was rapid and without event.

Case II

A male European, aged 26, admitted on January 28, 1937, was discharged on February 16. On January 27, in the afternoon, he suddenly developed severe headache with a temperature of 102° F. His bowels were opened twice that afternoon. The next day he had diarrhoea—six motions. The headache and fever persisted and he was admitted to hospital in the evening. On admission his temperature was 102.6° F. and the pulse 100.

He looked ill and was restless. He had nausea and retching, and complained of severe headache. The tongue was moist and coated; the fauces were injected. No signs of note in the chest or abdomen. Routine examination and inspection revealed no abnormalities in stool, urine, or blood smears. His fever remained at 102° F. for two days, after which he ran an irregular temperature, varying between 99° and 101° F. for the next five days. White blood counts on January 30 and February 1 showed 9,500 and 9,100 cells, with 79 per cent. and 86 per cent. of polymorphs respectively. Stool culture was negative for dysentery and enteric groups, as was the blood culture for the enteric group. The Widal reaction was twice negative for O agglutinins in enteric, *abortus*, and *proteus* (X19 and XK) groups. H agglutinins for enteric were present, but were without significance owing to previous T.A.B. inoculation. On January 30 he complained of severe abdominal pain. There was some general tenderness but no rigidity. The pain continued for the next three days, during

Cerebro-spinal Fluid Findings in Case II

Date	3/2/37	5/2/37	10/2/37
Cells per c.mm.	268	116	68
Lymphocytes	93.5%	100%	100%
Polymorphs	6.5%	nil	nil
Total protein	100 mg. %	90 mg. %	80 mg. %
Globulin	Present	Present	Present
Chlorides	703 mg. %	733 mg. %	—

which time his symptoms became aggravated. His restlessness increased, and there was that marked alteration in behaviour which is often associated with tuberculous meningitis. Further examination on February 3 revealed head retraction, a positive Kernig's sign on both sides, flexor plantar reflexes, sluggish knee-jerks, and brisk abdominal reflexes. Lumbar puncture on this date gave a slightly opalescent fluid under increased pressure. There were 268 cells per c.mm., of which 93.5 per cent. were lymphocytes and 6.5 per cent. polymorphs. The sugar content was normal, and the protein 100 mg. and chlorides 703 mg. per 100 c.cm. Wassermann and Kahn reactions of the fluid were negative. Gram and Ziehl-Neelsen stained smears showed no organisms.

Headache was relieved by lumbar puncture and improvement set in rapidly. Further lumbar punctures were performed on February 5 and 10, the results of which are set out below. On the 9th all his symptoms had disappeared and he was allowed up. His convalescence was rapid and uneventful.

Comment

Birch (1936), who has recently described some cases, gives a good account of the differential diagnosis in temperate climates. In view of what has already been written in regard to diagnosis in the Tropics, elaborate comment on the above cases is unnecessary. In future it is hoped to arrange for biological tests. It should then be possible to decide whether the lymphocytic meningitis which occurs here is identical with that in Europe and in North America.

The fact that in both instances quoted above the patients happened to be Europeans might give a wrong impression of the racial incidence of the disease. Actually they are representative of half a dozen similar cases—Asiatic and European—seen by me in the past two years. My notes in regard to the other cases are not complete and do not justify publication at this stage. I believe the incidence of the disease in Malaya is about the same as it is in Europe, and that it affects all races in the Peninsula.

Summary

Two cases of lymphocytic meningitis in Malaya are described. The symptoms were comparatively severe, but

recovery was complete in both cases. Clinically the disease was identical with the acute lymphocytic meningitis of Wallgren.

My thanks are due to Dr. Benjamin Chew for the routine investigations and for the blood counts. I am indebted to the Government pathologist, Singapore, and his staff for the Wassermann and Widal tests; and to the Director of Medical Services, Straits Settlements, for permission to publish this article.

REFERENCES

- Allen, I. A., and Spencer, F. M. (1935). *Med. J. Austral.*, **2**, 275.
 Armstrong, C., and Dickens, P. F. (1935). *Publ. Hlth. Rep., U.S.A.*, **50**, 831.
 — and Lillie, R. D. (1934). *Ibid.*, **49**, 1019.
 Bengtson, I. A., and Wooley, J. G. (1936). *Ibid.*, **51**, 29.
 Birch, C. A. (1936). *Practitioner*, **137**, 219.
 Collis, W. R. F. (1935). *British Medical Journal*, **2**, 1148.
 Craps, M. (1936). *Bull. Soc. Clin. Hôp. Brux.*, p. 78.
 Findlay, G. M., Alcock, N. S., and Stern, R. O. (1936). *Lancet*, **1**, 650.

BILATERAL TUBAL PREGNANCY

REPORT OF A CASE

BY

MURIEL B. McILRATH, M.B., B.S. Sydney,
 F.R.C.S. Eng., M.C.O.G.

Resident Obstetrical Officer, Withington Hospital, Manchester

Multiple extra-uterine pregnancy is a sufficiently unusual condition to merit a report of any case that may be encountered. Such cases may be grouped under the following three headings:

1. Intra-uterine pregnancy combined with extra-uterine gestation.
2. Multiple pregnancy in a single Fallopian tube.
3. Coincident pregnancy in each tube.

This last group may be subdivided into two groups: (a) successive or repeated gestation as seen in the case of Ferguson (1899); (b) simultaneous binovular gestation as seen in the cases reported by Launay and Seguinot (1911) and Lockyer (1917). The second subdivision of the last group is undoubtedly the rarest form of multiple extra-uterine pregnancy, and it is often extremely difficult to prove that a given case falls into this category.

Case History

The present case is that of a woman aged 28, married nine years, who was admitted to hospital on February 14, 1937. The following are the particulars:

Menstrual History.—Menstruation started when the patient was 15 years old; regular twenty-eight-day interval lasting three to six days; no vaginal discharge. The last menstrual period occurred on December 24, 1936; there had been no loss per vaginam since then.

Previous History.—There had been one normal confinement eight years ago, and one miscarriage at six weeks four years later. Three years ago the patient had pains in the lower abdomen, with vomiting and constipation. Two months later she had a similar attack, and six months ago another attack in which the pains were less severe but lasted longer.

Present Illness.—At 6 p.m. on February 13, 1937, the patient suddenly felt faint and had slight pain in the lower abdomen shooting upwards. This pain became worse and caused the patient to vomit. She fainted, and does not remember the events very clearly. The vomitus was brownish in colour. During the night the pains came on again, starting in the pelvis and shooting up into the abdomen. They were colicky in nature. She felt she was choking and vomited constantly, and on attempting to pass urine she fainted. On the morning of February 14 the pains were still present, and she also

complained of pain in the right shoulder. She was admitted to hospital on February 14 at 2 p.m. On examination the patient appeared shocked and the skin was cold; she was very pale, and the tongue brown and dry. Temperature 97.2° F.; respirations 22; pulse 120, thin and running. Heart and chest were clear. The abdomen moved evenly on respiration, and was not distended; there was no rigidity, but marked voluntary guarding. Very tender in right iliac fossa.

On vaginal examination a soft boggy mass was found in the pouch of Douglas, and an ill-defined soft swelling in the left fornix. The uterus was very tender, and voluntary guarding made adequate examination of the right fornix very difficult.

A diagnosis of ruptured tubal gestation was made.

Operation

Under general anaesthesia the abdomen was incised by a mid-line infra-umbilical incision, and free blood escaped on opening the peritoneum. On exploring the pelvis a mass was found on the left side formed by the left tube surrounded by omentum. Through a gap about one centimetre long on its anterior aspect chorionic tissue and old blood clot were seen protruding. Clamps were placed on the uterine end of the tube, the mesosalpinx, and the proximal end of the omentum, and the tube and adjacent omentum removed. Clots were removed from the pouch of Douglas and haemostasis secured. Then fresh blood was seen to be escaping from the fimbrial end of the right tube, in which there was now observed to be a swelling the size of a hazel nut impinging on to the uterus. The right tube was removed. Some difficulty was experienced in checking the bleeding from the uterine cornu. The right ovary was the seat of a corpus luteum of pregnancy, but no similar change was seen in the left ovary. Clots were taken from the peritoneal cavity and the abdomen closed. The patient was given a continuous intravenous saline transfusion for three hours, and then continuous rectal saline for forty-eight hours. Apart from some difficulty in getting the bowels to act the patient made an uninterrupted recovery, and left hospital in three weeks.

Macroscopically both tubes were found to be thin and tortuous. In the left tube on the anterior aspect about three centimetres from the uterus there was a rent one centimetre long through which chorionic villi and old blood clot protruded. The omentum had wrapped itself round the rent, but the adhesions were plastic and therefore of quite recent origin. The foetus unfortunately was not found. The attachment of the omentum to the site of rupture recalls the findings of Paul Bloch in the case reported by him (1931), though in his case the attachment was of longer duration. He too found no trace of either foetus. The right tube was unruptured; there was a swelling the size of a hazel nut at the site of foetal implantation. Haemorrhage had occurred around this area and in the lumen of the tube; the amniotic sac was ruptured, but no sign of a foetus was discovered.

Unfortunately the specimens were thrown away before a microscopical examination was made. Microscopical study was therefore impossible, but the macroscopic examination definitely revealed two areas of foetal implantation separated from each other by the fundus of the uterus. On the left side there was definite rupture and, on the right, evidence of tubal abortion. The foetus was not recovered on either side. There was only one corpus luteum of pregnancy, but according to Lockyer (1917) this is the usual finding, though in the case reported by Schockaert (1928) there were two, one in each ovary. Apart from this Schockaert's macroscopic findings closely resemble those seen here. That this case is not one of haematosalpinx on one side and tubal pregnancy on the other is proved by the presence of chorionic tissue on both sides and absence of distension of the tube in its whole length, as was found in Doran's case (1891).

The attempt to correlate the operative findings with the history is interesting. The first onset of pain was slight, but rapidly became worse and was apparently extremely severe. It was accompanied by internal haemorrhage and shock, as the patient fainted and was only semi-conscious for several hours. This apparently was caused by the rupture of the