ADVANCED EXTRA-UTERINE PREGNANCY

BY

A. PATRICK, M.B., F.R.C.S.Ed.

Honorary Assistant Surgeon, Hull Royal Infirmary

(WITH SPECIAL PLATE)

Extra-uterine pregnancy is not a very common condition, although within the last two decades it appears to have become more frequent, this probably being due to greater accuracy in diagnosis. Advanced extra-uterine pregnancy is, however, still a rarity, and the following case is worthy of mention from that point of view alone.

Case Record

Mrs. X., aged 33 years, was admitted to hospital on September 12, 1935, suffering from severe abdominal pain, sickness and vomiting, and collapse. The previous history showed that she was well advanced in pregnancy. Her last menstrual period commenced on April 16, 1935. During the first few months of pregnancy she complained of severe sickness and vomiting, which was always accentuated at the suppressed menstrual periods. After the third month the nausea and vomiting became much less. During the second and third months of pregnancy she had several what she termed bad attacks of lower abdominal pain, associated with an increase in the sickness and vomiting. The abdomen just above the pelvis was always very tender to the touch. At no time during the pregnancy was there any uterine haemorrhage.

Past History.—The patient had been married for six years, during which time she was never pregnant. She had consulted her family doctor about her sterility, and, in fact, it had been suggested that she should have a dilatation and curettage. Her menstrual history was that the menses were always regular—every twenty-eight days and lasting six to seven days. There was no history of leucorrhoeal discharge or of any frequency of micturition. She had always enjoyed perfect health, and there have been no previous serious illnesses.

Examination.—The patient looked extremely ill. She was absolutely blanched, and her respirations were of a sighing character. Every time she attempted to sit up in bed she collapsed. Her pulse rate was 120, rapid and feeble. Her tongue was dry and she complained of great thirst. Her conjunctivae were practically colourless. She had severe abdominal pain associated with sickness and vomiting. The abdomen did not move on respirations. There was a fullness of the lower abdomen in the middle line. The abdomen was held very rigid, and she complained of the greatest tenderness in the left iliac fossa. On vaginal examination, which was extremely difficult owing to the rigidity of the abdominal muscles, one could palpate a large mass filling the pelvis; this I considered was the normal pregnant uterus. Other signs of pregnancy were well advanced.

Diagnosis.—It was obvious from an examination of the patient that the condition was a very acute one, and that she was suffering from internal haemorrhage, which I considered was due to a ruptured uterus. The fact that she had had no uterine haemorrhage during the pregnancy, and that the pregnancy was so advanced, made me disregard the probability of this being a ruptured ectopic gestation. However, the seriousness of the condition was realized, and I advised immediate operation.

Operation.—On September 12, under general anaesthesia, the abdomen was opened through a lower mid-line incision. On reaching the peritoneum I noted the bluish-black coloration behind it indicative of intraperitoneal haemorrhage. On opening the peritoneum I found the abdominal cavity filled with partly clotted and partly fluid blood. In the lower part of the wound was a large mass, and through a tear in the free surface of this mass could be seen a foetus covered by the amniotic sac. On further examination it was found that this was a large ectopic gestation. The uterus was slightly enlarged and congested. Both ovaries were normal. I resected the

grossly distended Fallopian tube, and then, hastily removing all the large pieces of blood clot, closed the abdomen. No intravenous saline was given, and the patient rapidly improved; within two weeks she had made a complete recovery.

Specimen

The large mass to the left as shown in the photograph is the grossly distended Fallopian tube. The placenta with the vessels running over its surface can be plainly seen, firmly attached to the Fallopian tube. The amniotic sac was opened and the foetus placed in this position in order to photograph it (see Special Plate).

Commentary

According to Schumann ectopic gestation occurs in only 1 in 267 pregnancies, or 0.38 per cent. The age incidence, according to Wynne, is between 24 and 33 years.

The position of the gestation varies. It can therefore be (1) cornual, (2) isthmal, (3) ampullary, (4) tubo-ovarian, or (5) ovarian. The commonest type is the ampullary, into which category this case belongs. This part of the tube is capable of great distension, and allows an ectopic gestation to develop much further than any other part.

Numerous conditions, such as inflammation in the lower pelvis, salpingitis, diverticulitis, and appendicitis, have been found to be predisposing factors to ectopic gestation, but in this case no history could be obtained of any such cause, nor was there evidence of anything pathological in the other tube. Congenital anomalies are often cited, and this might well have been the case here, but I have no proof.

It is common in cases of ectopic gestation to find that there is some history of sterility. Farrar quotes 16.6 per cent. of cases in which ectopic gestation followed a long period of sterility, and 30 per cent. in which there had been one previous child.

One interesting feature in the present case was the absence of the atypical menstruation as described by Bandler. In the majority of cases some slight uterine haemorrhage is complained of, but here the patient strongly denied that there had ever been any.

Schumann states that the greater proportion of ectopic pregnancies rupture at the eighth week, and that very few continue after the twelfth week of pregnancy. The most interesting point about the case here recorded was the fact that the pregnancy developed up to nearly five months.

I wish to express my indebtedness to Dr. D. Stenhouse Stewart, who took the photograph.

L. T. Davidson, K. K. Merritt, and T. T. Chapman (*Amer. J. Dis. Child.*, January, 1937, Part I, p. 1) gave eighteen premature and fifteen full-term infants irradiated evaporated milk during the first six months of life to determine the degree of protection from rickets afforded by this means alone. Judged by x rays, none of the eighteen premature infants were entirely free from rickets; six showed slight, nine moderate, and three marked rickets. Of the fifteen full-term infants, four were free from x-ray signs of rickets throughout the period of the study, one had moderate rickets, and in the other eleven there was only slight rickets. In the group of premature infants in this study, who were given irradiated evaporated milk as their sole source of vitamin D, the rickets tended to develop earlier than in the full-term infants. Irradiated evaporated milk has been shown to be considerably less efficacious for the protection of the premature infant against rickets than is metabolized vitamin D milk. When the two are given under identical conditions the inferiority of the irradiated milk is believed to be due entirely to the smaller concentration of vitamin D units.

JAMES L. BROWN: INTRAPLEURAL PRONTOSIL IN STREPTOCOCCAL EMPYEMA

FIG. 1.—March 1, 1937. Very large left basal empyema.

FIG. 2.—April 1, 1937. No sign of any fluid.

A. PATRICK: ADVANCED EXTRA-UTERINE PREGNANCY

Photograph showing the foetus and, on the left, the grossly enlarged Fallopian tube, with the placenta attached.

EDGAR P. WATERS: PARTIAL EXTRA-UTERINE PREGNANCY

Anterior view of the uterus showing the wide gap of which the rounded edges had become adherent to the parietal peritoneum.