BRITISH MEDICAL JOURNAL

LONDON SATURDAY JUNE 5 1937

EARLY DIAGNOSIS AND TREATMENT OF HEART FAILURE*

BY

WILLIAM EVANS, M.D., F.R.C.P.Lond.

Assistant Physician, London Hospital

(WITH SPECIAL PLATE)

Few illnesses can more readily compel a medical attendant to apply immediate therapeutic measures than heart failure. So alarming may be the symptoms and so great the anxiety surrounding the patient that a demand for hurried action is everywhere evident. Never is the necessity greater, however, for the doctor to make haste slowly, to act promptly but not precipitately, and yet again to treat cautiously but expeditiously. The need to do something is always present, but it should be no greater than the need to know everything before doing anything; so that an immediate perception of the expected course of the illness, and the processes involved, is an essential preliminary to the competent treatment of heart failure.

The Mechanism of Heart Failure

Although the true mechanism of heart failure continues to remain an unsolved problem, our knowledge of it on the one hand has helped us to conceive certain principles in treatment, and on the other hand our lack of knowledge is unlikely to have withheld anything revolutionary from present-day methods of dealing with the condition. The changes during failure are familiar to us; it is the evolution of these changes which eludes precise exposition.

The back-pressure failure theory envisages stasis of blood in the vascular areas which drain in the direction of the affected heart chamber, so that obstruction in the mitral valve is expected to produce the following sequence of changes: distension of the left auricle, engorgement of the lungs, distension of the right ventricle and of the right auricle, engorgement of the veins of the neck and cyanosis, distension of the liver, and oedema of the lower extremities. Recently it has been shown (Thompson and White, 1936) that left ventricular strain from hypertension, aortic valve disease, or infarction of the left ventricle is the commonest cause of hypertrophy of the right ventricle, which occurs regardless of the presence of clinical evidence of left ventricular failure, although this increases the degree of hypertrophy.

The forward-pressure failure theory, supported by Sir James Mackenzie, postulates systemic failure following inadequate output of the left ventricle, while pulmonary symptoms in the same way are the outcome of failure of the right ventricle. This deficiency in the outflow of blood results in a depletion of the vascular areas and in ischaemia of the tissues situated in front of the affected heart chamber.

It is likely that both back-pressure phenomena and a diminished cardiac output are features that exist together in many if not in most instances of congestive heart failure, but the knowledge of the part which each plays

in the development of failure is unlikely to contribute in a greater measure to the treatment of the condition than to indicate the need of relieving congestion and of increasing the efficiency of cardiac contraction.

Size and Position of the Heart in Failure

Cardiac enlargement is usual in failure, but it is rarely the direct result of failure, and its distribution and extent depend chiefly on the lesion which is providing the cause of the failure. Either with or without failure the progression of cardiac enlargement is very gradual, so that sudden dilatation of the heart seldom occurs. Whenever such an event has been implied from a clinical examination, and displacement of the apex beat to the left has been noticed, three influences contriving to produce this change must be kept in mind before attributing it to cardiac enlargement. They are tachycardia, distension of the liver and ascites producing elevation of the diaphragm, and a right-sided hydrothorax. Even on radioscopy caution must be exercised when assessing the development of cardiac enlargement, so that the "squatting" posture of the heart, assumed as a result of a raised diaphragm from any cause, must not be interpreted as enlargement. Again, the possibility of pericardial effusion enlarging the cardiac silhouette as seen on radioscopy should receive attention in any case where an exacerbation of failure signs has occurred with the formation of effusion within serous cavities.

It is held by some that hydrothorax (usually right-sided) from heart failure does not cause cardiac displacement,

FIG. A.—Displacement of heart in failure. Diagram showing superimposed outline of heart in teleradiographs 1 and 2 of the Special Plate.

but this is fallacious (see Figs. 1 and 2 on Plate and Fig. A in text). Certainly it does not displace the heart to the same extent as does an inflammatory pleural effusion, because the quantity of fluid is not so great, and, further, whenever the effusion in heart failure has become considerable pulmonary congestion with pleural effusion

* The substance of a lecture delivered before the North Glamorgan and Brecknock Division of the British Medical Association, March 16, 1937.

occurs on the opposite side, and this opposes the initial displacement.

Evidence of Early Failure

The need for recognizing early heart failure is only equalled by the need for early treatment, and early treatment depends on early diagnosis. In order to gain this end the earliest evidence of failure must be sought in those patients known to suffer from heart disease which is expected to give rise to heart failure sooner or later. In them symptoms develop gradually and almost imperceptibly, so that our awareness of this happening will usually lead us to seek for premonitory signs and not to wait for the fully developed clinical picture of heart failure characterized by cyanosis, considerable breathlessness, conspicuous engorgement of the veins of the neck, pulmonary congestion with right-sided hydrothorax, enlargement of the liver with ascites, oedema of the lower extremities, and scanty urine with albuminuria. The recognition of early congestive heart failure depends on attention to the following essentials: undue breathlessness to customary exertion, venous engorgement, pulmonary congestion, hepatic distension, and a familiarity with the causes of heart failure.

Undue Breathlessness

Breathlessness on exertion occurs in health. With advancing age or with the onset of obesity this natural reaction to increased physical performance becomes accentuated. The breathlessness which heralds the commencement of heart failure is characterized by its occurrence during exercise which hitherto produced no breathlessness, or which is in excess of that measure which was customarily induced by a standard form of exercise. Thus the hall-mark of pathological breathlessness is present when exercise which usually produced no breathlessness induces it, when exercise which previously initiated breathlessness causes undue breathlessness, and when the breathlessness has shown progression within a relatively short period of time.

Venous Engorgement

An increase in the venous blood pressure may be regarded technically as the first evidence of heart failure, but unfortunately its estimation cannot be carried out with the same ease as the registration of the arterial blood pressure. The direct method of recording the venous pressure by connecting a venepuncture needle to a manometer filled with citrate solution cannot for obvious reasons become common practice in clinical medicine, so that some other means denoting a change in the venous pressure is desirable. Sir Thomas Lewis (1930) directed attention to a method which permits an approximate clinical estimate of the venous pressure. He pointed out that in health all veins which lie at a higher level than the manubrium sterni remain collapsed and those below are distended. Thus in a healthy patient in the reclining posture with the head resting on a pillow the external jugular veins are distended for about one-third their course from the clavicle to the angle of the lower jaw. In a patient presenting a slightly raised venous pressure the column of venous distension reaches the centre of the sternomastoid muscle or higher. The veins in the neck are not evident in a healthy subject in the erect position, but in established heart failure they are visible as cords reaching from the clavicle to within a variable distance of the mandible. This clinical indicator of the venous pressure should gain significance not only on account of its relative accuracy, but also on account of the simplicity of the observation.

Pulmonary Congestion

When persistent crepitations are elicited over the bases of the lungs, mostly on the right side, from heart failure, the presence of a considerable degree of pulmonary congestion can be presumed. Earlier evidence of pulmonary engorgement is marked by vascular congestion in the hila of the lungs. Careful auscultation along each side of the spine over the middle zones will often disclose, especially after coughing, isolated coarse crepitations at the end of a deep inspiration, but commonly the presence of this degree of congestion can only be made out on radioscopy; certainly the extent of hilar congestion can only be told by this method. The picture obtained is a characteristic one presenting wing-like shadows spreading from each hilum, and caused by an increase in the density of the pulmonary vessels, which are depicted here and there in cross-section and appear like ink blots (Plate, Figs. 3 and 4).

The help provided by radioscopy in the recognition of early heart failure has not yet gained the reputation it deserves. Nor should the method be applied solely in diagnosis, for it also provides an index in the evaluation of progress during treatment, so that the extent of hilar congestion as determined at intervals by radioscopy determines the advance or retreat of the process of failure (Plate, Figs. 3 and 4).

Hepatic Distension

An increase in the volume of the liver is a sensitive indicator of an increase in the venous blood pressure. Although this natural manometer does not supply accurate quantitative values, when enlargement of the liver has been made out it serves as early evidence of heart failure, so that whenever failure is suspected or expected distension of the liver should be sought. Palpation immediately below the right costal margin may elicit tenderness in a patient presenting hepatic distension, but a quest for the descended liver edge as the hand travels upwards from a low position is a more reliable method of examination.

Specific Causes of Heart Failure

Familiarity with the conditions capable of precipitating heart failure ensures timely and efficient attention to a patient in whom failure threatens. To anticipate heart failure is often to postpone it; to remain unaware of its causes and its early manifestations is to allow its grosser signs to develop. The causes of heart failure may be conveniently considered in two groups, as suggested by Parkinson and Clark-Kennedy (1926); but here a reference to these will only include in the first place any special features which facilitate early diagnosis, and secondly those details concerning the treatment of each group which cannot be regarded as common to all cases of failure.

Treatment of Established Heart Failure

Certain general principles apply to the treatment of most of the common examples of heart failure, so that in the first place it is opportune to consider these and later to detail any particular remedy or procedure which may apply when certain specific causes are present. An outline of those therapeutic measures to be adopted in the general case of heart failure is facilitated by referring to three distinct clinical types—established heart failure, early heart failure, and abrupt heart failure.

The designation "established heart failure" is applicable when a patient presents characteristic physical signs of failure which have developed gradually and have pro-

gressed during the absence of treatment or with inadequate treatment. No rigid rules can be formulated to govern the management of all cases of established heart failure, but although every patient must be studied individually the treatment should generally provide for adequate rest, digitalization, and diuresis.

REST

In ambulatory cases a limitation of activity is of first importance in treatment, and in order to gain the fullest benefit from this curtailment of exercise the patient's usual daily habits should be ascertained and a programme modifying the day's activities should be complete in detail. A later rising time in the morning and an earlier retiring time at night, periods of rest in the reclining posture after meals, walking slowly on level ground and avoiding hurrying and climbing, are among the instructions imparted to the patient in whom established heart failure is controlled by digitalis medication. At night the provision of extra pillows promotes rest and sleep.

In non-ambulatory cases absolute rest is sought, and a patient is discouraged from indulging in any exertion, not even turning in bed. Again, the head and shoulders should be supported in the upright posture, and although devices to meet this necessity have been improvised in the form of chairs, extension of these amenities to the construction of special beds has received but scant attention. Such a bed was designed by Sir Thomas Lewis, but the adoption of this helpful instrument in the treatment of heart failure in the wards of a hospital has proceeded very slowly.

Inseparable from any method adopted to ensure rest in a case of heart failure is the alleviation of the patient's natural apprehension and the provision of sleep. Alcohol, in that it is conducive to sleep and causes a feeling of well-being, may be dispensed with benefit, but in no sense must it be regarded as a "heart stimulant." A mixture containing 20 grains of potassium bromide and chloral hydrate often suffices as a hypnotic, or resort may be made to the barbiturates (sodium luminal 1 to 2 grains) or morphine (1/2 grain dissolved in the mouth, or 1/4 grain hypodermically). Certainly these drugs should be used with determination, as they are important agents in the treatment of heart failure.

DIGITALIS

Good wine needs no bush, and certainly the utility of digitalis in the treatment of heart failure does not need to be advertised; it has proved its worth. What does require emphasis is that it should be used in adequate doses, so that our duty in treating heart failure is not so much the giving of digitalis as establishing a state of digitalization. Digitalization is only present when the patient's symptoms are lessened, when diuresis has resulted, when the pulse rate has been slowed to 70 per minute or under, when the ventricular rate has been similarly diminished and the pulse deficit annulled, and when depression of the R-T segment is a feature of the electrocardiogram.

A choice of the preparation of digitalis must necessarily be made before determining the dosage. It is most desirable to adhere to the administration of a single preparation, and acquire experience of its effects in different patients. Nothing is gained, in fact a good deal is lost, by substituting various preparations of digitalis, either in the same patient or in different patients. It is far better to acquire exact knowledge of the value of a single preparation than to nourish a doubt concerning the comparative value of a number of preparations. The following may be regarded as representative of the digitalis group

and dependable when used in the indicated daily doses: powdered leaf (4 grains), tincture (45 minims), Nativelle's granules (1/125 grain), and digoxin tablets (0.45 mg.). Although these doses are comparable and basic they are not to be regarded as absolute standards in all patients with established heart failure. The dosage should be adjusted according to the severity of the failure, weight of the patient, and particularly the length of time required to establish a state of digitalization. In most instances the suggested standard dose is the approximate one, but often it will need increasing and sometimes decreasing.

DIURESIS

To produce efficient diuresis should be a constant aim in every case of established heart failure, and there is more than one method available to attain this object.

That rest in bed combined with a diminished fluid intake can produce prominent diuresis has not gained sufficient recognition. Fig. B illustrates this effect in a

FIG. B.—Chart showing diuresis following rest in bed and a diminished fluid intake in a patient with mitral stenosis and heart failure with normal rhythm.

female, aged 36 years, with mitral stenosis, in whom the features of early heart failure presented. Cognizance of this happening is necessary in the evaluation of any remedy credited with diuretic properties: in the patient instanced here unmerited repute might have been accorded any drug professing a diuretic action if it had been introduced when the patient was voiding scanty urine on admission to hospital. The lesson in this connexion is twofold: first, it impresses the need of prescribing rest combined with a diminished fluid intake (1½ pints daily) in patients presenting heart failure; and, secondly, it emphasizes the importance of adequately controlling all clinical tests conducted to assess the diuretic properties of any drug.

Foremost among the drugs which can effect diuresis in heart failure, when administered by mouth, is digitalis, and in most instances it should be preferred to other remedies in this group, such as ammonium chloride or the purine diuretics—for example, theobromine, diuretin, theophylline, or euphylline.

Intravenous and intramuscular salyrgan or novurit in 2 c.cm. doses are pre-eminently the most powerful diuretics we possess, and they form important implements now available to combat the effects of heart failure and to help restore cardiac efficiency. Parkinson and Thomson (1936) have shown that novurit gives satisfactory results when used as a suppository, although of course its diuresis is not so profuse as when it is introduced intravenously

or intramuscularly; but the advantages afforded by the simplicity of this method of administration are obvious. The prescribing of ammonium chloride along with salyrgan increases diuresis, but this is not always considerable. The most recent advancement in the treatment of heart failure has been the addition of mercurial diuretics to the other effective therapeutic agents—rest, digitalis, and hypnotics.

The effect of total ablation of the thyroid gland in heart disease presenting paroxysmal nocturnal dyspnoea and heart failure has to be discussed. Impressions concerning the benefit gained from this procedure are best got from a study of the results obtained in seventy-eight cases [thirty-six by Blumgart and others (1935), and forty-two by Levine and Eppinger (1935)]. The therapeutic benefit enjoyed by some of these patients corresponded with the lowering of the basal metabolism, and when this became reduced to minus 30 per cent. myxoedema frequently intervened, and this in turn responded to small doses of thyroid without materially detracting from the improvement resulting from the lowering of the basal metabolism. If heart failure was present with persistent oedema the operative risk increased considerably, so that it is evident that total thyroidectomy is not often desirable in instances showing failure, and certainly this procedure should be undertaken only when the recognized methods of treatment have failed and when the operation can hold out a reasonable prospect of considerable improvement.

Treatment of Early Heart Failure

A patient with heart disease is commonly examined during the pre-failure period, when he will derive greater benefit from observation and instruction than from the initiation of any form of zealous treatment. Medicinal treatment, apart from a placebo, is unlikely to produce material benefit during this period. Certainly any attempt at digitalization at this stage is ill advised and without promise of improvement, so that it should be discouraged even though it might be given without harm.

In particular the patient is to be instructed on the subject of activities and diet. While it is permissible or even desirable to indulge in limited or graduated exercise, exertion is prohibited. In connexion with diet it is desirable that it should not be heavily salted, and stress is directed to two main essentials—namely, the avoiding of large or indigestible meals and, if obesity is a feature, the resorting to a fat-reducing diet.

To gain the co-operation of the patient during this pre-failure phase should be a first consideration. No amount of enthusiasm or even heroism in treatment will produce improvement if the patient is not the doctor's ally. Nor must the psychological side be neglected, so that encouragement and just optimism should be dispensed along with firm dissuasion where needed. The over-cautious should receive encouragement, and the reckless should be restrained and be made to realize the consequences of any disregard to instructions. Improved health should be promised on conditions and as a reward of attention to advice.

If the signs of early heart failure are beginning to collect and there is increased breathlessness, slight venous engorgement in the neck, and hilar congestion detected by radioscapy, apart from the precautions already enumerated concerning exercise and diet, periods of controlled rest should be outlined; these will be in the mornings and evenings, after the midday meal, and at the week-ends. Digitalis should be added if auricular fibrillation has set in. The administration of mercurial diuretics combined

with a much-reduced fluid intake should be conjoined with the week-end rest.

The treatment adopted for this group of patients almost merits the designation "prophylactic treatment"; at any rate, it aims at postponing the signs of established heart failure. The measure of success gained from the care devoted to these cases justifies the zeal involved in collecting cases within the scope of this group.

Treatment of Abrupt Heart Failure

In some patients in this class the precipitation of heart failure may be an unforeseen contingency, and this particularly applies to cardiac infarction. It may also occur unexpectedly in hypertension, although paroxysmal nocturnal dyspnoea will often serve as a premonitory sign. Sudden exacerbation of failure symptoms may also occur in cases with established heart failure hitherto controlled by treatment, or even in some instances of early failure; in both of these groups either a period of increased activity and exertion or some intercurrent illness has caused a damaged heart to fail suddenly.

It is in this class of cases that the urge to do something is so strong and good judgement so often yields to mismanagement created by over-enthusiasm. The term "cardiac stimulant" designates the drug whose purpose it is to help the heart in this sudden crisis. Many drugs have gained a reputation in this field, but usually this has not been merited and has been gained through uncontrolled clinical impressions. Strychnine, camphor, and pituitary are in this category, and have far survived their usefulness in this connexion. A warning is overdue concerning the danger of the wholesale and indiscriminate use of proprietary preparations imputed to be of this class of "cardiac stimulants" in virtue of their action, variably claimed to be on the medulla, on the peripheral circulation, on the coronary circulation, or as "cardiac hormones." A plea for resort to controlled clinical assay is made in order to regain a proper perspective of the value of these potions.

Strophanthin has seldom shown superiority over digitalis in cases of precipitate heart failure. Digitalis, particularly digoxin, has great value in abrupt heart failure associated with auricular fibrillation. Oral administration of 1 mg. of digoxin can lower the ventricular rate in auricular fibrillation within a few hours. Its success when the heart rhythm is regular is not so assured. Little or no advantage is gained by intravenous administration of digitalis.

Whenever cyanosis is a prominent feature and there is conspicuous venous engorgement in the neck, venesection is good practice. If restlessness and sleeplessness present, hypnotics must be administered in adequate doses after providing support for the head and shoulders and making the patient comfortable.

If oedema and ascites are features salyrgan is indicated. In the case of a female patient with much oedema it is necessary to anticipate the considerable diuresis by tying in a urinary catheter in position so that the patient's energy is in this way conserved. Salyrgan medication has dispensed with the necessity of purging an oedematous patient for the purpose of ridding him of superfluous tissue fluids.

Heart Failure in Specific Conditions

Certain special features in the mechanism, early diagnosis, and treatment of specific conditions which may initiate failure will now be described.

HYPERPIESIA

Whenever congestive heart failure appears in a patient and the heart rhythm is regular, hyperpiesia may be regarded as the likely cause. To bear this in mind is to recognize the underlying condition when the failure has produced a fall in the blood-pressure value. Sometimes when the onset of failure has caused the systolic figure to fall to a normal level the diastolic value has continued to remain high, permitting the diagnosis of hypertension. Often, however, a fall in the diastolic blood pressure has accompanied a similar change in the systolic blood-pressure value, and the diagnosis then rests on a record of a raised blood pressure during the pre-failure period and the finding of cardiac hypertrophy chiefly involving the left ventricle. An ominous sign has appeared when heart failure has been added in a case of hyperpiesia. Improvement may occur for a time, but longevity can hardly be mentioned in connexion with prognosis.

Digitalis is without any great reputation in the treatment of hypertensive heart failure, but it always merits a trial in individual cases, as it will usually produce some degree of improvement. Certainly it is the treatment of choice in the less common instances where auricular fibrillation is present.

The mercurial diuretics salyrgan and novurit hold first place in the treatment of hypertensive heart failure, and the administration of these should be augmented by providing organized periods of rest and a diminished intake of fluid (Figs. 5 and 6).

Further, when failure has set in as a complication of hypertension, paroxysmal nocturnal dyspnoea (cardiac asthma) is a frequent accompaniment, and demands specific treatment in addition to the general regime adopted for the failure. The patient must be propped up in bed, and potent hypnotics such as sodium luminal, $1\frac{1}{2}$ grains, or morphine $\frac{1}{4}$ grain, administered each night until the general condition is improved from the use of diuretics. Digitalis medication should also be given a trial in these cases.

AORTIC INCOMPETENCE AND AORTIC STENOSIS

Aortic incompetence from syphilitic aortitis and endocarditis invariably precipitates heart failure with a regular cardiac rhythm, and so does rheumatic aortic incompetence if the mitral valve has remained healthy or is only slightly damaged. Similarly the cardiac action is regular in aortic stenosis with heart failure if the aortic valve is the only valve affected. As in the case of hyperpiesia the onset of failure is a serious complication, and signals a grave outlook.

The failure associated with aortic valve disease also simulates hypertensive failure in that digitalis assumes second place to salyrgan or novurit, except in the less frequent instances where failure has recurred with auricular fibrillation.

CARDIAC INFARCTION

Although arrhythmia, especially auricular fibrillation, is sometimes present in cardiac infarction, in the majority of cases normal rhythm is preserved. The signs of failure may appear abruptly in this condition, and if the area of infarction is a large one the signs are correspondingly increased; thus pulmonary congestion becomes prominent, and not infrequently crepitations are found scattered over both lungs. The liver, too, may become distended early in the illness, and slight oedema of the ankles may appear.

Treatment of the failure in this instance resolves itself into dealing with cardiac infarction, so that rest ensured by morphine is the procedure to adopt in patients in this group.

CONGENITAL CARDIOVASCULAR DISEASE

Heart failure is not a common feature of congenital cardiovascular disease, and pulmonary infection and subacute bacteriaemic endocarditis are the more likely complications to ensue. Cyanosis in these cases usually only serves to indicate the presence of a venous-arterial shunt, and is seldom an expression of the presence of heart failure. In those instances of congenital cardiovascular defects where cyanosis appears later in life the onset of heart failure may have caused cyanosis by producing a reversal of an existing arterio-venous shunt into a venous-arterial shunt. The symptoms of dyspnoea and cyanosis, therefore, in congenital heart disease are not commonly manifestations of congestive heart failure; they do not call for special care or treatment except in those few instances which show frank signs of failure.

The cardiac rhythm continues to be regular after failure has set in unless congenital heart-block is present as a rare feature of interventricular septum defect. When failure has set in it is usually progressive, although adequate rest and treatment with mercurial diuretics and digitalis may produce temporary improvement.

EMPHYSEMA AND BRONCHITIS

Too often in the past a patient giving a history of a productive cough occurring mostly during winter months and presenting great breathlessness, cyanosis, and abundant adventitious sounds in the chest, has been regarded as a case of heart failure. More notice should be taken of emphysema as a common cause of extreme breathlessness, and less of its influence on the heart. The severe breathlessness associated with emphysema is the direct result of a diminished pulmonary ventilation, and is only rarely the sequence of heart failure. The effects of emphysema on the heart have recently received attention. Parkinson and Hoyle (1937) determined the size and shape of the heart by radiology in eighty patients suffering from a high grade of emphysema. Enlargement of the pulmonary artery stem and its branches and of the conus of the right ventricle was present in about one-half of the cases. Changes due to hypertension were also commonly met with. About one-third showed no cardiac enlargement. Heart failure was infrequent (thirteen out of eighty), and in seven of the thirteen cases hypertension was a feature. They stated that congestive failure purely from emphysema and chronic bronchitis was rare and had a grave prognosis, so that recurrent failure was almost unknown.

Of even greater importance in the management of a case of heart failure is the recognition of the effects of emphysema and chronic bronchitis in a patient presenting heart disease. The occurrence in these cases of attacks of acute bronchitis produces an exacerbation of symptoms and may cause alarm, for the breathlessness may be extreme and accompanied by a distressing cough with the voiding of thin sputum and often haemoptysis. In the presence of mitral stenosis and auricular fibrillation such symptoms have been attributed too readily to heart failure, and the added pulmonary signs have been erroneously assigned to pulmonary congestion from failure instead of to diffuse bronchitis and emphysema. The treatment of each condition is so dissimilar as to urge the necessity

of recognizing in many cases of heart disease this prevalent "bronchitic element," and of deciding its influence in producing the symptoms before treatment is commenced. Adrenaline or ephedrine and a mixture containing potassium iodide, stramonium, and ammonium carbonate prove more beneficial than an increase in the dose of digitalis. It is likely, from experience with this type of case, that certain proprietary adrenaline-like preparations have gained an over-measure of trust in the treatment of all cases of heart failure.

NEPHRITIS

Heart failure is a common event in nephritis when hypertension is present. In chronic nephritis cardiac asthma is often an added feature. This frequent occurrence of heart failure in nephritic hypertension is not universally realized, particularly in hypertension of acute nephritis. Ellis (personal communication) found evidence of heart failure in twenty instances among 100 patients with acute nephritis, and in nine of these it was severe. The importance of a more general acceptance of the common incidence of heart failure in nephritic hypertension is found in a wider application of specific remedies in these cases to cardiac rather than to renal failure.

ADHERENT PERICARDIUM

Adhesions may form between the parietal and visceral layers of the pericardium, partially or completely obliterating the sac, and yet produce no cardiac embarrassment nor result in cardiac enlargement unless valvular disease is present. In another group calcification of the pericardium may be added, but without symptoms. Where extrapericardial adhesions have bound the heart to the surrounding structures (mediastino-pericarditis) cardiac enlargement is often present, but even here coincident valvular disease may account for the enlargement. A special feature of a fourth group is the constriction of the superior and inferior venae cavae, and to a less extent the heart, by adhesive pericarditis, giving rise to a characteristic clinical syndrome (Pick's syndrome). The effects of these changes is to produce a condition comparable to that of heart failure but without pulmonary congestion, so that the symptom of breathlessness is never prominent in the initial stages. The means of recognizing cases in this group has been clearly detailed by Paul White (1935), who refers to the leading clues in the diagnosis in the following order: The insidious onset of dropsy and oedema of the lower extremities in a young person. Preponderant enlargement of the liver with ascites. Undue prominence of the jugular veins with increase of the venous pressure. Absence of any heart disease or nephritis in a case presenting dropsy. Small and regular pulse with a low systolic blood pressure and pulse pressure. Evidence of chronic pleurisy. Calcification of the pericardium seen on radiology. Low-voltage curve in the electrocardiogram with flattening or inversion of the T-wave in one or more leads. Previous history of pericarditis and polyserositis.

Pericardial resection by an experienced thoracic surgeon is the only method of treatment which holds promise of recovery from the effects of constrictive pericarditis, and the number of patients who have gained benefit from this surgical procedure is fast increasing.

ARTERIO-VEIN ANEURYSM

When a small communication has been established between an artery and a vein there is no demonstrable change in either the anatomy or function of the heart, but

if the fistulous opening is a large one then symptoms and signs of heart failure develop. It is the arterio-venous aneurysm resulting from trauma, therefore, and situated in the extremities, particularly the thigh and leg, which specially produces changes in the heart. Here breathlessness and palpitation are common symptoms, and considerable enlargement of the heart, especially the left ventricle, is demonstrable on radiology. The pulse is regular, a little rapid, and is collapsing in character with an increase of the pulse pressure; obliteration of the arterio-venous shunt by digital compression raises the systolic and particularly the diastolic blood pressure, and the collapsing nature of the pulse disappears; the pulse rate also falls abruptly. With the fistula open the venous blood pressure is increased and pulmonary congestion is visible on radiology. Extirpation of the aneurysm causes the cardiac enlargement and the pulmonary congestion to disappear within a short period of time (see Figs. 7 and 8), thus providing a rare example of reversible cardiac enlargement.

MITRAL STENOSIS

Most instances of mitral stenosis showing evidence of heart failure present arrhythmia, usually in the form of auricular fibrillation, but occasionally as auricular flutter; in some the rhythm remains regular. It is likely that early indications of heart failure are invariably present before the onset of fibrillation, and that the adoption of efficient treatment during this pre-fibrillation period postpones the onset of fibrillation. Once auricular fibrillation has become established in a case of mitral stenosis digitalis is pre-eminently the medicine to prescribe. In the uncommon instances of mitral stenosis presenting prominent and aneurysmal dilatation of the left auricle the absence of gross heart failure symptoms may be a notable feature.

THYROTOXICOSIS

Slight cardiac enlargement occurs commonly in patients with thyrotoxicosis in the absence of auricular fibrillation and heart failure, but failure is seldom if ever present with a normal cardiac rhythm. Again, not all cases of thyrotoxic fibrillation show evidence of failure. When it has set in digitalis will produce some improvement, but subtotal thyroidectomy after adequate treatment with Lugol's iodine is unquestionably the proper course to adopt. An attempt to re-establish normal rhythm by quinidine administration during the pre-operative period is of doubtful value.

COMPLETE HEART-BLOCK

When complete heart-block is complicated by heart failure and when Adams-Stokes attacks are absent digitalis is worthy of trial because it often produces clinical improvement and the failure symptoms disappear without causing undue slowing of the pulse. In cases subject to Adams-Stokes attacks digitalis is contraindicated as it usually increases the incidence of the attacks.

Conclusion

In conclusion it may be said that in the diagnosis and prevention of heart failure it is important to seek early evidence of this in patients presenting heart disease which can ultimately precipitate failure, and that in treatment it is necessary to individualize, to treat with determination but not over-zealously, to prescribe remedies whose worth has been established by repeated and controlled clinical trial, to gain the full co-operation of the patient, and to give constant encouragement.

REFERENCES

- Blumgart, H. L., Riseman, J. E. F., Davis, D., and Weinstein, A. A. (1935). *Amer. Heart J.*, **10**, 596.
 Ellis, A. W. M.: Personal communication.
 Levine, S. A., and Eppinger, E. C. (1935). *Amer. Heart J.*, **10**, 736.
 Lewis, Sir T. (1930). *British Medical Journal*, **1**, 849.
 Parkinson, J., and Clark-Kennedy, A. E. (1926). *Quart. J. Med.*, **19**, 113.
 — and Thomson, W. A. R. (1936). *Lancet*, **1**, 16.
 — and Hoyle, C. (1937). *Quart. J. Med.*, **30**, 59.
 Thompson, W. P., and White, P. D. (1936). *Amer. Heart J.*, **12**, 641.
 White, P. D. (1935). *Lancet*, **2**, 539, 597.

LUPUS VULGARIS

TREATMENT BY INTRADERMAL INJECTION OF
HYDNOCARPATES

BY

J. EDGAR WALLACE, M.D.,

Assistant Tuberculosis Officer, West Sussex County Council,
late of Lancashire County Council

(WITH SPECIAL PLATE)

The numerous methods which have been devised for the treatment of lupus vulgaris bear adequate testimony to the obstinate nature of this disease. For this reason the encouraging results that have recently been obtained by the intradermal injection of esters of hydnocarpus oil deserve more widespread attention than they have as yet received. Leprosy workers have for some years made increasing use of this method in dealing with cutaneous lesions, and the analogy between this disease and tuberculosis led to the account by Sir Leonard Rogers (1933) of a case of extensive lupus vulgaris successfully treated in this way.

Soon after this publication treatment was commenced in the present series of fifteen cases, and preliminary reports were issued in the succeeding two years (Wallace, 1934 and 1935). More recently Burgess (1935) has described a series of eleven cases, in which clinical cure was obtained in seven and improvement in the remaining four.

Choice of Preparation

The substance first used in the present investigation was that known as moogrol. This is prepared by esterizing the total acids of hydnocarpus oil, and consists of ethyl hydnocarpate and ethyl chaulmoograte, together with the esters of the other acids present in this oil. As reactions were sometimes unduly severe, iodized moogrol (containing 0.5 per cent. of iodine) was also tried. This, however, seemed less efficacious, and in addition produced in the scar a bluish-brown discoloration of considerable persistence.

Later, at the suggestion of Dr. T. A. Henry of the Wellcome Chemical Research Laboratories, both these forms were abandoned in favour of a third preparation, phenyl-ethyl hydnocarpate. The change proved to be a satisfactory one, as reactions became much less marked, and treatment could thus be advanced more rapidly. Phenyl-ethyl-hydnocarpate, which is therefore the preparation of choice, is now on the market under the trade name of eulykol (Burroughs Wellcome and Co.). Its less irritant effect has been made possible by eliminating the small uncrystallizable portion of the total fatty acids during the preparation of the ester.

Technique

Injections are made intradermally with an ordinary 1 c.cm. hypodermic syringe fitted with an agla 240 needle

(B.W. and Co.). This is a 3 mm. needle fitted with a guarded point. Patients attend weekly, and at each sitting a series of wheals 1/2 cm. to 1 cm. in diameter are raised in the lupus patches and nodules. The total amount of injection at any one time has not exceeded 1 c.cm., but Burgess, using phenyl-ethyl hydnocarpate exclusively, has injected as much as 5 c.cm. Reactions can be diminished by spacing the wheals as far apart as possible, and no area should be injected a second time until all inflammatory changes have subsided.

Lastly, it is important to avoid subcutaneous injection. This not only causes more severe diffuse reactions, but also fails to produce the desired effect, for, as Beatty (1932) points out, the lupoid foci are mostly in the corium, and there is a layer of corial connective tissue between them and the subcutaneous tissue. These precautions are particularly to be observed in lesions around the eyes, where swelling of the loose tissues may be greater than elsewhere, and may cause some concern to the patient.

Effect of Injections

No general disturbance of any sort has been noted. The injection itself is practically painless, but during the next forty-eight hours a certain amount of redness and swelling appears, accompanied by a slight degree of pain and tenderness. The pain lasts for one or two days only, but the swelling may not subside for a week or longer.

As treatment continues a slowly progressive flattening of nodules and plaques occurs, together with a disappearance of scales and crusts. The granulation tissue between the lupoid areas diminishes. Tiny ulcers may appear in and around the site of injection. A varying degree of redness prevails, but gradually the active lesions are replaced by pale smooth scar tissue. Several months may elapse after the cessation of treatment before the final effect is seen. During this time the tendency to relapse is no less apparent than in other forms of lupus treatment. Careful observation is necessary to ensure that any recurrent nodules can be reinjected at the earliest possible moment.

The esters of hydnocarpus oil are absorbed very slowly from the skin, and their effect on lupoid tissue is continued long after the purely irritant reaction has worn off. For this reason one hesitates to regard the action entirely as a non-specific one, unless it be due, as Muir (1932) suggests, to the liberation of antigens.

Summary of Cases

The fifteen cases summarized in Tables I and II all attended a tuberculosis dispensary of the Lancashire County Council. Previous treatment had consisted of local and general artificial light therapy and occasional "spiking" with acid nitrate of mercury, but the lesions had mostly reached that curiously resistant stage where the disease appears to be neither advancing nor healing. All other forms of treatment were stopped at the commencement of the new method so that a fair trial could be made.

The length of treatment has varied in individual cases from six months to three years, although, in the longer cases, as the condition improved injections have frequently been spaced out at fortnightly or even monthly intervals. Treatment has been unduly prolonged in some cases, since considerable caution was exercised at the onset owing to the experimental nature of the method.

Progress has undoubtedly been more rapid than that made by artificial light treatment alone, but there is no reason why eulykol injections should not be supplemented by general carbon-arc baths, as in some of the cases

WILLIAM EVANS: EARLY DIAGNOSIS AND TREATMENT OF HEART FAILURE

FIGS. 1 AND 2.—Teleradiographs showing displacement of heart in failure. Fig. 1, before treatment, shows greater displacement of heart from right-sided hydrothorax than Fig. 2, after treatment.

FIGS. 3 AND 4.—Teleradiographs showing pulmonary congestion in heart failure. Characteristic hilar shadows appear in both skiagrams, but are less conspicuous in Fig. 4, taken after three weeks' rest, than in Fig. 3, taken before resting.

WILLIAM EVANS: EARLY DIAGNOSIS AND TREATMENT OF HEART FAILURE

FIGS. 5 and 6.—Teleradiographs showing relief of pulmonary congestion from heart failure by rest in bed along with a diminished fluid intake and salyrgan administration. Fig. 5 was taken before treatment, and Fig. 6 one month after commencement of treatment, in a case presenting hypertensive heart failure.

FIGS. 7 and 8.—Teleradiographs showing arterio-venous aneurysm. Heart diminishes in size and pulmonary congestion is relieved following resection of aneurysm. Fig. 7 taken before operation; Fig. 8 taken one month after removal of arterio-venous aneurysm of lower thigh.