first pregnancy. Spontaneous labour occurred at thirty-six weeks, with a difficult breech delivery. The baby was somewhat shocked, and weighed 4 lb. 15 oz. at birth. On discharge at 2 weeks his weight was 5 lb. 4 oz.

Case 11.—Baby boy. Normal healthy parents. Spontaneous delivery at thirty-five weeks. His weight was 5 lb. 3½ oz. at birth, 5 lb. 5 oz. at 2 weeks, and 8 lb. 12 oz. at 2 months.

Table of Results (No Deaths)

Case No.	Sex	Weight		Age on Discharge
		At Birth	On Discharge	
1	Girl	4 lb. 7½ oz.	4 lb. 7½ oz.	2 weeks
2	"	3 lb. 10 oz.	3 lb. 10½ oz.	2 "
3	"	3 lb. 12 oz.	3 lb. 14 oz. + operation	4 "
4	"	3 lb. 12 oz.	3 lb. 10 oz.	2 "
5	"	4 lb. 8 oz.	5 lb. 0 oz.	2 "
6	Boy	3 lb. 13 oz.	4 lb. 1 oz.	2½ "
7	Girl	3 lb. 8 oz.	4 lb. 2½ oz.	2½ "
8	"	4 lb. 7½ oz.	4 lb. 12 oz.	2 "
9	"	4 lb. 1½ oz.	4 lb. 7½ oz.	2 "
10	Boy	4 lb. 15 oz.	5 lb. 4 oz.	2 "
11	"	5 lb. 3½ oz.	5 lb. 5 oz.	2 "

Summary

Details are given of eleven premature and one sick baby treated with progynon. This preparation of oestrin, given orally, appears definitely helpful to premature infants. The babies cause less anxiety. The feeds are taken better; the initial loss of weight appears less and is usually made up sooner. No baby to whom it was given died.

REFERENCES

- Aschheim (1927). *Arch. Gynäk.*, **132**, 179.
 Martin, E. (1929). *M Schr. Geburtsh. Gynäk.*, **82**, 66.
 Moncrieff (1936). *Arch. Dis. Child.*, **11**, 2.

R. Opsahl (*Norsk Mag. Laegevidensk.*, April, 1937, p. 9) has scrutinized the records of necropsies at the Ullevaal Hospital in Oslo for the twenty-year period 1916–35, with special reference to primary malignant disease of the lungs. During this period 2,005 cases of carcinoma and sarcoma came to necropsy, the malignant disease being in some of these cases an incidental finding. In addition to the fifty-six established cases of primary malignant disease of the lungs (twenty-eight men and twenty-eight women) there were fourteen cases in which this diagnosis was uncertain and eight in which it was possible but not probable. The age incidence was greatest between 50 and 60 (seventeen cases)—a little earlier in life than in other forms of malignant disease, which reached their maximum frequency between 60 and 70. Only in eleven of the fifty-six cases were the new growths diagnosed clinically. The duration of the disease was usually between four and twelve months. The fifty-six cases were classified in four five-year groups and in relation to the total number of cases of malignant disease in each five-year group. In the first five years primary malignant disease of the lungs constituted 1.8 per cent. of all the cases of malignant disease in this period. The corresponding figures for the next three five-year periods were 2.7, 3.1, and 3.1 respectively. Putting these figures to the test of modern statistical criteria, the author comes to the conclusion that his material gives no support for the assumption that there is a real increase in the incidence of primary malignant disease of the lungs.

VACCINE TREATMENT OF MEASLES

BY

**A. A. CUNNINGHAM, M.D.Dub., M.R.C.P.Lond.,
D.P.H.**

*Senior Medical Officer, North-Western Hospital (L.C.C.)
late Senior Medical Officer, Park Hospital (L.C.C.)*

Measles is typically a disease of childhood, the majority of the patients being attacked in the first five years of life, and although it is still regarded by the general public as a trivial illness, its complications, both immediate and remote, are severe enough to constitute a major medical problem. Bronchopneumonia remains the most serious and dreaded complication, and is largely responsible for the present mortality rate.

Wynn of Birmingham has for some years advocated the use of a special vaccine in the treatment of pneumonia and bronchopneumonia. His results have apparently been good in those cases in which the diagnosis is made early and the appropriate vaccine treatment instituted in the first twenty-four to forty-eight hours of the illness. The vaccine for bronchopneumonia contains equal numbers of pneumococci, haemolytic streptococci, and Pfeiffer's bacilli. Fresh cultures are used and the organisms are killed by heat. Wynn (1936) recommends 600 million organisms—that is, 200 millions of each organism—as a suitable injection dose for an adult. Children require proportionately smaller doses, but at 1 year old 20 millions of each organism can be given with impunity. Three such injections subcutaneously at intervals of twenty-four hours are recommended.

Production of Reactions

The vaccine treatment of acute infections has not found favour owing to the fear of producing reactions in a patient who is dangerously ill. According to Wynn this is a fallacy, as reactions occur in chronic infections only when patients are sensitized and specific antibodies are present in the circulation. In the early stage of an acute infection antibodies are absent, and therefore these reactions do not occur. Immunotherapy by vaccine treatment aims at the production of antibodies at an earlier stage than would usually be the case. Specific antibodies do not appear for several days, but the vaccine used is stated to have an immediate effect in that it stimulates the production of non-specific antibodies—various bactericidal substances which appear in the circulation. These are later supplemented by the specific antibodies, but the greatest stress has been laid on the value of the non-specific immediate response to the vaccine injections, which if given sufficiently early are claimed to abort the infection.

In measles bronchitis is present in the early stages in all but the mildest and modified cases. It usually clears up in a few days; but occasionally, and especially in poorly nourished and debilitated children, it progresses to bronchopneumonia. The detection of bronchopneumonia is very difficult at times, and it is only by taking into consideration the general condition of the patient, in conjunction with the various physical signs and symptoms, that one can hope to arrive at a reasonably accurate diagnosis. Areas of lung collapse are met with in young children with acute bronchitis and may easily be mistaken for areas of pneumonic consolidation.

In this investigation we decided to treat only those patients suffering from measles in the various stages of development up to the time of the appearance of the full-blown morbilliform eruption. All late cases with

fading rashes were therefore excluded. The question of using alternate cases on admission as controls was carefully considered, but this was regarded as impracticable, so it was decided to treat by vaccine those cases admitted to five specified wards and to treat as controls those admitted to five other wards. Admissions to the various measles wards were in strict rotation as vacancies occurred.

Treatment

The stock vaccine used was prepared in the Southern Group Laboratory under the direction of Dr. J. E. McCartney. It was killed by heating at 60° C. for thirty minutes, and was made from young active cultures. The vaccine was so constituted that there were 600 million organisms in each cubic centimetre, 0.1 c.cm. being regarded as a suitable dose for a child 1 year old, 0.2 c.cm. for a child 2 years old, and so on. The first injection was given for the most part on the day of admission, but in a few instances, when the child was admitted late at night, the injection was deferred until the next morning. The routine treatment consisted of three subcutaneous injections of equal amount at intervals of twenty-four hours. All cases of measles had a small prophylactic injection of diphtheria antitoxin on admission.

The following tables give briefly the incidence of the various complications in the specially treated group and in the corresponding set of controls.

TABLE I.—*Chest Complications*

Series	Cases	Lobar Pneumonia	Bronchopneumonia (measles)				Bronchopneumonia (measles and pertussis)		Empyema (strep.)	Deaths (due to measles)	
			No.	%	Deaths	Mortality	No.	Deaths		No.	%
Vaccine	236	1	24	10.1	5	20.8	2	1	1	5	2.1
Control	278	—	25	9.0	7	28.0	1	1	—	7	2.5

TABLE II.—*Ear Complications*

Series	Cases	Measles					Measles (with intercurrent disease)		Deaths (directly due to ear complications)
		S.O.M.* (right)	S.O.M. (left)	S.O.M. (bilateral)	S.O.M. Total	Mastoidectomy	S.O.M. (measles and pertussis)	S.O.M. (scarlet fever)	
Vaccine	236	10	9	16	35 = 14.8	1	1	1	—
Control	278	10	15	14	39 = 14	1	1	1	—

* S.O.M. = Suppurative otitis media.

TABLE III.—*Complications in Age Groups*

Age Group	Vaccine Series (236 Cases)						Control Series (278 Cases)							
	No. in Group	Broncho-pneu-monia		Suppur-ative Otitis Media		Deaths		No. in Group	Broncho-pneu-monia		Suppur-ative Otitis Media		Deaths	
		No.	%	No.	%	No.	%		No.	%	No.	%	No.	%
0-2	95	15	15.8	20	21.0	4	4.2	106	16	15.1	7	16.0	4	3.8
2-3	53	3	5.6	9	16.9	1	1.9	67	5	7.4	7	10.4	3	4.5
3-4	32	1	3.1	5	15.6	—	—	40	1	2.5	4	10.0	—	—
4-5	23	3	13.0	1	4.3	—	—	31	2	6.4	3	9.7	—	—
5+	33	2	6.0	—	—	—	—	34	1	2.9	8	23.5	—	—

Discussion

In this investigation 236 patients with measles were treated with a vaccine prepared according to Wynn's formula, and a corresponding control series of 278 cases was treated on similar general routine lines without vaccine. The majority of the patients were under 5 years of age, the youngest being about 9 months old. The injections were without exception innocuous, and we did not observe any general febrile reaction which could reasonably be attributed to the vaccine. Slight local reactions were seen from time to time, but were rare.

An analysis of results shows that there has been no demonstrable diminution in the incidence of the two most common complications of measles—namely, bronchopneumonia and otitis media—nor has the mortality rate been affected to any appreciable extent. Mr. B. E. Spear of the statistical department of the London County Council, to whom I am indebted for the following report, states "that from the statistical point of view the results justify the negative conclusion reached as to the treatment." The claims made by Wynn as to the value of the vaccine in aborting pneumonic infections and in reducing the mortality rate have not been confirmed by this investigation in young children with measles as the primary infection. It is true that the total number of patients treated was relatively small, but one feels that no useful purpose would be served by the continuance of vaccine treatment in measles with the vaccine as at present constituted.

I wish to express my gratitude to Dr. H. Stanley Banks, medical superintendent of the Park Hospital (L.C.C.), for his advice and criticism, and for permission to publish this paper.

REFERENCE

Wynn, W. H. (1936). *British Medical Journal*, 1, 45.

Clinical Memoranda

Slimming Drugs and Cataract With Notes of a Case

During the past few years compounds of dinitrophenol have been used as metabolic stimulants in the treatment of obesity, especially in America. In 1935 Tainter, Stockton, and Cutting published results of treatment with 2:4-dinitrophenol in doses of 100 mg. daily for one week, then 200 mg. for several weeks, with gradual increase to 300 mg., followed by reduced doses if toxic symptoms occurred, such as dyspnoea, oedema, urticarial rashes, and prostration. They reported that no pathological changes were produced in the blood cholesterol, liver, or blood count in four years of treatment, and weight was reduced. Koch, Lee, and Tainter (1935) reported no changes in the liver or any other tissue in dogs killed by lethal doses.

In 1935 Horner, Jones, Boardman, and other authors published cases of cataract in human beings following this treatment. They found that after the first signs appeared the development was rapid. The cornea was normal; the aqueous showed a slight slit-lamp flare; the anterior capsule was spotty, dry, and lustreless; and fine grey subcapsular cloudy opacities were present. Later irregular pearl-grey opacities appeared deeper in the cortex, with polychromatic specular reflection from the posterior cortex. Marked swelling of the lens followed. The cause was thought to be increased permeability of the lens capsule. Secondary glaucoma sometimes occurred.