ABNORMALITIES OF GROWTH AND DEVELOPMENT

THE CLINICAL AND PATHOLOGICAL ASPECTS*

BY

H. GARDINER-HILL, M.D., F.R.C.P.*Physician to Out-patients, St. Thomas's Hospital*

ARREST OR RETARDATION—DWARFISM AND INFANTILISM

The conditions of dwarfism and infantilism can be considered together, for although dwarfism is not infrequently found without infantilism the two are more frequently combined. Both represent disturbances in the growth sphere, but whilst in dwarfism the defect is skeletal and limited to the skeleton, in infantilism the retardation of development is general.

The term dwarfism implies a defect in stature below the normal standards but does not necessarily signify a pathological state, for dwarfism under certain conditions may occur without any evidence of disease. In simple dwarfism the skeleton is of small size, but the bodily characters and functions—especially in the sex sphere—are appropriate to the age of the individual. The skeleton may be proportioned or disproportioned. If disproportioned the defect may be either in the trunk or the limbs—an abnormality of the spine or the long bones. In either case the underlying cause may be developmental or acquired.

The term infantilism is more difficult to define, and is often used ambiguously. Strictly speaking, it should be reserved for more or less permanent conditions in which the physical and psychological attributes of childhood persist into adult life—smallness of stature, underdevelopment of the musculature, a head large in proportion to the body, comparatively short limbs, lack of function of the genitalia, and certain qualities in the mental sphere. This strict interpretation of the term, however, is not generally adhered to, and there is a tendency to include conditions of temporary delay in development in which the stage reached is less than that appropriate to the age of the individual. This is perhaps justifiable if the wide sense in which the term is used is clearly understood. The term infantilism, however, should not include pathological states such as eunuchoidism, in which retardation or arrest of development in one sphere—in this instance the sex sphere—without any general disturbance of growth and development is found.

It is obvious from our consideration of the many factors which influence the growth process that there must be a number of different causes of dwarfism and infantilism. Physiological and pathological varieties both occur. Dwarfism may be a separate entity in simple hereditary dwarfism and in achondroplasia and allied conditions. The various forms of rickets are also accompanied by growth defects, and in the coeliac and renal forms, where there is a pronounced interference with general metabolism, infantilism results too.

The different forms of infantilism are difficult to classify, but there are two main groups. In the first, due to endocrine deficiency, the underlying cause may be hypopituitarism, hypothyroidism, or hypogonadism. If the underlying condition is untreated the physical and psychological attributes of childhood persist into adult life. To the second group the term cachectic has been

applied, and it includes those in which retardation or arrest of development has resulted from malnutrition, metabolic disturbance, or chronic infective disease. A convenient classification of the different varieties of dwarfism is as follows:

Simple Dwarfism:

1. Simple hereditary dwarfism.
2. Simple dwarfism due to developmental skeletal disease: (a) achondroplasia, chondro-osteo-dystrophy, and dyschondroplasia; (b) osteogenesis imperfecta.
3. Simple dwarfism due to acquired skeletal disease: (a) simple rickets and late rickets; (b) spinal caries and spinal deformities due to other causes—for example, poliomyelitis.
4. Hypergonadal dwarfism.

Dwarfism and Infantilism:

1. Cachectic.
2. Cachectic infantilism associated with more specific changes at the growth cartilages: (a) congenital syphilis; (b) scurvy; (c) coeliac rickets; (d) renal dwarfism.
3. Endocrine: (a) hypopituitary; (b) hypothyroidic; (c) hypogonadal.

1. True Hereditary Dwarfism

True hereditary dwarfism should not be regarded as a pathological condition, for apart from the defect in stature neither the skeleton nor other organs of the body show any evidence of abnormality or disease. These individuals are entirely normal apart from their small size. Puberty occurs at the normal age and sex function is normal. Hereditary dwarfism is essentially a hereditary characteristic and dependent on chromosomal constitution. The racial form is typified by the Norwegian Lapps, Central African pygmies, and inhabitants of the Andaman Islands. Many striking examples of the familial variety have been reported, and a complete description of them has been given by Hastings Gilford (1911). Growth is slow and far below average, and this applies to the differential growth of the skeleton too.

2. Simple Dwarfism Due to Developmental Skeletal Disease

(a) ACHONDROPLASIA, CHONDRO-OSTEO-DYSTROPHY, AND DYSCHONDROPLASIA

Amongst the developmental conditions producing simple dwarfism achondroplasia and the closely related entities, chondro-osteo-dystrophy and dyschondroplasia, must be considered. Some authorities take the view that achondroplasia, dyschondroplasia (Ollier, 1899), chondro-osteo-dystrophy (Morquio, 1929), and hereditary deforming dyschondroplasia (metaphyseal aclasis) are only different manifestations of one pathological state—dyschondroplasia, by which term is implied a nutritional defect of the growth cartilage. This view is held by Parsons (1936), who suggests that achondroplasia could be regarded as the extreme form—namely, an aplasia—with the others as intermediate varieties.

From the pathological and clinical aspects there are certain points of distinction.

In achondroplasia the most striking feature is a mucoid degeneration and absence of cell proliferation in the growth cartilages. The normal palisade arrangement of the cells—that is, the normal scaffolding—is absent. Ossification is consequently hindered by the absence of this scaffolding, and longitudinal bone growth interfered with.

The term chondro-osteo-dystrophy has been applied to several ill-understood conditions. The earlier cases were reported by Brailsford (1929) and Morquio, who described

* The Oliver-Sharpey Lectures delivered before the Royal College of Physicians of London on March 16 and March 18, 1937. Lecture I was published last week.

irregular development of the metaphyses and epiphyseal ossification centres. The latter appeared as multiple foci which either fused gradually or became distorted and fragmented in joints subjected to pressure. In the vertebral column affected vertebral bodies became irregular in shape and size, the result of compression from the superimposed weight of the vertebral column. The chondro-osteo-dystrophy of Brailsford and Morquio bears a close resemblance to cases reported by Dickson Wright (1930-1) as generalized osteochondritis, in which every epiphysis in the body became fragmented and distorted, and by Thursfield and Lightwood (1930-1) as "stippled epiphyses"—a case with clinical features suggestive of achondroplasia but, radiologically, fragmented epiphyses. The fundamental disturbance in the growth cartilage of this case, described later by Harris (1933e), was a mucoid degeneration, more pronounced in the centre of the epiphysis than at the metaphyseal line, but otherwise identical with that in achondroplasia. There was an attempt in certain areas to form cartilage cell columns, but no continuity in the orientation of the cells for any distance.

The term dyschondroplasia has been coined by Ollier to describe the condition at the growth cartilage in which cartilage cell proliferation—that is, the scaffolding—is present in excess, but the subsequent senescent processes of degeneration, calcification, and ossification do not occur. As the bone develops masses of unossified cartilage persist at the metaphysis. Hereditary deforming dyschondroplasia is a variety of this condition with additional features—an unmodelled metaphysis and exostoses.

From the clinical aspect all these conditions have much in common, and dwarfism is generally pronounced. The achondroplasiac certainly appears to represent the extreme form in which the whole skeleton is affected and the changes are present at birth. The characteristic appearance is produced by a combination of normally sized trunk and short limbs, particularly the proximal segments, a relatively large head and small face with depressed nasal bridge, and trident hand. The disproportion between limbs and trunk is most pronounced. A lumbar lordosis usually produces an undue prominence of the abdomen, and the gait is of an odd waddling character. Dentition, sex, and mental development are normal. Radiology of the skeleton shows characteristic changes. In the child and adolescent the space between the epiphysis and metaphysis is smaller than normal and the metaphyses tend to be unduly splayed and broadened near the joints, producing the characteristic "petunia"-like appearance, due to failure in the modelling process.

The dwarfism of chondro-osteo-dystrophy is very similar, but the child is normal at birth and the head and facial changes are not found as in achondroplasia. The proportions of the skeleton depend on whether the vertebral column or limb bones are more affected. In some cases both are the site of characteristic changes; in others the defect is localized to one or other. The diagnosis can be confirmed by radiology, when fragmentation of the epiphyses will be found. In some instances this condition improves as the child grows older, but although a satisfactory growth rate is resumed a degree of dwarfism results.

The clinical pictures of dyschondroplasia and hereditary deforming dyschondroplasia are similar to the above, but the limb bones are chiefly affected, and deformities, bowing or genu valgum, may be present. In both a short-limbed skeleton is usually found. In the hereditary type the characteristic radiological features are an irregularly expanded metaphysis with exostoses. The progress of

the condition can be judged from the story of a family, and particularly two sisters, who came under my personal observation. In the younger, a dwarf aged 8, the features of dyschondroplasia were first noticed at the age of 18 months. At 8 she had a strikingly disproportioned skeleton: symphysis-vertex 54.5 cm., symphysis-soles 35.5 cm. The limbs were short and there was enlargement and typical radiological changes at the metaphyses. She happened to be the ninth child of a family of ten, of which the third, fourth, and tenth members were also subjects of this condition. Radiological examination of the metaphyses of her elder sister (No. 4, aged 14) by the same radiologist, at the age of 11, had shown an identical condition, but a second examination carried out three years later—at the age of 14—demonstrated that the metaphyseal abnormalities had disappeared. At the age of 14 this girl was still strikingly dwarfed and her height was only 4 ft. 4½ in. Her skeleton was disproportioned and of the short-limbed type: symphysis-vertex 28½ in.; symphysis-soles 24½ in. The changes of puberty were normal and well advanced.

(b) OSTEOGENESIS IMPERFECTA

Osteogenesis imperfecta must be included among the developmental skeletal diseases leading to simple dwarfism owing to the scoliosis, stunting of long bones, and deformities following the fractures which invariably occur. Apart, however, from the deformities from healed fractures, the bones are actually shorter, thinner, and smaller than normal. The epiphyseal cartilage is normal, but in the metaphysis the trabeculae are slender and delicate and are widely separated by interstices filled with cellular connective tissue. The underlying abnormality in osteogenesis imperfecta is a generalized disturbance of osteogenesis; according to Knaggs (1926), a failure of differentiation of osteoblasts. The resulting growth defect depends much upon the deformities and the age at which the condition manifests itself. In the infantile form there may be stunting and evidence of fractures at birth, but these children seldom survive to adulthood. In the adolescent type the child may appear normal at birth, but the tendency soon manifests itself, and repeated fractures result in deformities and skeletal dwarfism. The radiological changes in the bones are characteristic: an almost complete absence of cancellous tissue, and the cortex appears as a faint line.

3. Simple Dwarfism Resulting from Acquired Skeletal Disease

In discussing the factors which limit growth, reference was made to Harris's investigations showing that changes of premature senescence in growth cartilage and temporary arrest of growth result from the infective and metabolic diseases of childhood. In fact, a quick succession of severe diseases during adolescence has been observed to cause permanent arrest of growth with complete union of the epiphyses. Dwarfism then results, but often without infantilism. Apart from these cases there are two types of dwarfs which require consideration in this section: (a) due to simple and late rickets, and (b) to spinal caries or other spinal deformity.

(a) SIMPLE DWARFISM AND RICKETS

The era before the recognition of the cause of rickets was responsible for many examples of rickety deformity and simple dwarfism. The pathological process, as is well known, is due to vitamin D deficiency, and consists of a softening and deformity of the bones from deficient

deposition of calcium phosphate, and especially imperfect calcification and ossification at the epiphyseal line. The cartilage cells proliferate excessively, and become converted in a patchy manner into tissue resembling bone but very deficient in lime salts—the so-called osteoid tissue. The intensity of the process is directly proportional to the rapidity of growth, for the non-growing animal does not develop rickets. A similar vitamin D deficiency in adolescence leads to late rickets, a comparatively rare condition, and, in adults, when all bone growth has ceased, to a general rarefaction and softening of the skeleton—osteomalacia. Owing to the softening of the bones in these rickety conditions, the skeleton, and particularly those parts which are subjected to muscular strain or the influence of posture, may be greatly deformed. In the past rachitic dwarfism with severe deformities was not infrequently seen. Now, since the recognition of the cause of rickets, it is a condition of extreme rarity. A study of a series of children with defective growth, however, does suggest that inadequately treated simple rickets in infancy may produce considerable degrees of dwarfism, noticeable only in later years. With adequate treatment rachitic osteoid tissue is replaced by healthy growing bone in the course of three or four months, and this is the measure of interference with growth. Nevertheless, in my series of cases brought for defective growth alone, 20 per cent. gave a history of infantile rickets as the only likely causal factor. Slight stigmata of past rickets could usually be found, and the skeleton was disproportioned as a rule and shorter in the lower measurement than in the upper. The intelligence and genital sphere developed normally.

Late rickets is also a cause of dwarfism, and must be distinguished from rickets of the coeliac and renal types. Typical rickety changes occur in the skeleton, and stunting of growth may be pronounced. Late rickets, rarely seen now, tends to develop during the active growth periods of the second dentition and puberty.

(b) SIMPLE DWARFISM DUE TO SPINAL CARIES AND OTHER SPINAL DEFORMITIES

For the sake of completeness mention must be made of the forms of simple dwarfism due to spinal caries and other spinal deformities. The skeletal deformities are of the short-trunk type. A good illustration is the hunch-backed dwarf in whom angulation of the spine has resulted from vertebral tuberculosis. The site of the vertebral lesion is usually in the lower thoracic region, and the tuberculous process begins as a rule in a single vertebra, as a nutritional disturbance of the marrow following a tuberculous endarteritis. Absorption and caseation take place, and sooner or later the vertebral shell collapses from the superimposed weight of the vertebral column. A kyphosis and angular deformity then results. Other diseases of the vertebral column occurring during the developmental period may produce dwarfism of the short-trunk type, and occasionally examples have been observed from muscular atrophy following anterior poliomyelitis.

4. Hypergonadal Dwarfism

The occurrence of hypergonadal dwarfism has already been fully discussed under the heading of macrosomia praecox. It was then pointed out that, although these syndromes are associated with transitory overgrowth in childhood, accompanied by sexual precocity, dwarfism ultimately results owing to premature epiphyseal fusion.

Before we pass on to a consideration of the conditions in which dwarfism and infantilism are combined, let us briefly review the main features of simple dwarfism. In

some instances the statural defect is largely due to deformity, as in the dwarfism of osteogenesis imperfecta, spinal caries, and other conditions of the group. Much depends on the extent of the skeletal abnormality, which varies from a generalized disturbance of osteogenesis with multiple fractures in osteogenesis imperfecta to a localized infective process, perhaps in a single vertebra, in spinal caries. In the main varieties of simple dwarfism, on the other hand, the site of the pathological disturbance is the growth cartilage. It varies from mucoid degeneration and scaffolding defects in achondroplasia and allied conditions to cartilage cell proliferation with imperfect calcification and ossification in rickets. In hypergonadal dwarfism, premature senescent changes occur in the cartilages from the action of the sex hormones, and there results early epiphyseal fusion with cessation of growth. Similar changes and dwarfism have been found as a result of a series of infective diseases in adolescence. The growth and differential growth curves in these individuals are determined largely by the age of onset and severity of the pathological process, and by the nature and degree of permanence of the lesion. The most severe statural defects are found in achondroplasia and allied conditions where the degenerative process dates to foetal life, and, in the present state of our knowledge, is permanent and irremediable. In rickets, on the other hand, even when inadequately treated, the cartilage changes are more temporary and the damage done not irreparable. A disproportioned skeleton is liable to result because the long bones at the "springing-up" periods are especially interfered with.

Dwarfism and Infantilism

Conditions in which dwarfism and infantilism are combined may be considered under three main headings: (1) simple cachectic infantilism, in which retardation or arrest of development, general and skeletal, results from some serious chronic wasting disease in childhood; (2) cachectic infantilism complicated by more specific changes at the growth cartilages as in congenital syphilis, scurvy, and coeliac and renal dwarfism; (3) endocrine infantilism where arrest of development results from anterior pituitary, thyroid, or gonadal deficiency.

1. Cachectic Infantilism

Cachectic infantilism has been variously described as Lorain's (1871) infantilism, infantilo-chétivism, or dystrophic infantilism. The essential aetiological factor is a chronic wasting disease occurring during the growth period. This causes retardation of general development and "lines of arrest" in the growth cartilages. The skeletal and other systems of the body are affected proportionately. Any condition of prolonged illness in childhood may produce such a result, but especially important are gastrointestinal diseases in which there is long-standing interference with nutrition. Chronic abdominal tuberculosis and coeliac disease provide the most typical examples. The latter condition will be considered in a separate category owing to the rickety changes which occur. The infantilism of congenital syphilis, scurvy, and chronic interstitial nephritis will also be grouped with coeliac disease, as here also special epiphyseal changes complicate the picture. The nutritional disturbance from dietetic restriction in young diabetics before the introduction of insulin therapy has also been observed to result in considerable retardation of growth and development, amounting in severe cases to infantilism. Other causes of cachectic infantilism are inanition from underfeeding, chronic dysentery, and intes-

tinal worms. In regions where malaria and hookworm disease are prevalent both have been found to lead to serious delay in development. Cases of hepatic and hepatosplenic infantilism have been reported in the literature, but it seems reasonable to classify them under this rather than under a separate heading. The interference with metabolism can be held responsible for the delay in development. Hepatic disease does not appear to be a common cause of infantilism, for the reason probably that, if the hepatic lesion is severe, death occurs before development is interfered with, whilst, if the lesion is slight, liver function is not seriously affected and infantilism does not result.

In cases of cachectic infantilism seen in later years an adult type skeleton is usually attained. Much depends on the severity of the illness and the time at which the pathological process began. The interference with growth at the growth cartilages is not, as a rule, so pronounced as in the rickety and endocrine forms. The nature of the pathological changes and the process of formation of lines of arrested growth have already been considered. Development may be severely retarded, but adulthood is ultimately attained.

In describing the clinical features of cachectic infantilism the original description of Lorain (1871) cannot be bettered. The physique remains feeble and the general bodily form is small; the skeletal proportions are normal; the genitalia appear diminutive, but are in proportion to the bodily size. There is no defect in intelligence, but the mind remains childish. The changes of puberty do not occur at the normal age, and may be delayed until 18 or 20. Even then they may not be normally established, but in later life fuller development is usually reached. This description is very typical of the majority of cases of cachectic infantilism. It need only be added that radiological examination shows that osseous development is not disproportionately retarded if considered in relation to body size. Lines of arrested growth can be demonstrated in the diaphyses.

2. Cachectic Infantilism Complicated by More Specific Changes at the Growth Cartilages

(a) CONGENITAL SYPHILIS

The retardation of growth and development not infrequently observed in congenital syphilis could be reasonably included under the heading of simple cachectic infantilism were it not for the specific changes which may occur in the growth cartilages, interfering with growth in the long bones and so producing a disproportioned dwarfism. These changes are usually of the nature of a syphilitic osteochondritis, as originally described by Wegner (1870), a chronic inflammatory process which destroys the normal arrangement of the cartilage cell columns. The calcifying cartilage is interspersed with fibrous tissue characteristic of chronic inflammation, and the margin of the newly formed bone is irregular. Radiographically, the free edge of the ossification zone resembles a saw in which the teeth are irregular in size. The clinical picture of retardation of physical development in congenital syphilis would not be so distinctive were it not for the presence usually of some of the other well-known stigmata of this disease. Other conditions in this group showing short-limbed disproportion of the skeleton, scurvy and coeliac and renal dwarfism, require differentiation, but the radiological and other signs should make the diagnosis clear. Harris (1933e) has pointed out one important distinction between these conditions: whilst syphilitic osteochondritis especially affects the slow-growing cartilages, those at the proximal end of the femur and distal end of the tibia, the changes of scurvy

and rickets are usually manifested at the fast-growing cartilages in the neighbourhood of the knee.

(b) SCORBUTIC INFANTILISM

Although scurvy from vitamin C deficiency—Barlow's (1883) disease—in its chronic form and of sufficiently long standing to be productive of infantilism must now be a very rare condition, when it has been observed specific changes have been found at the growth cartilages which have seriously interfered with growth. Pathologically, the process in the growth cartilage consists of a haemorrhage from the capillary loops, which disrupts the newly formed bone and calcified cartilages and thus interferes with their nutrition. The haemorrhagic clot may be replaced by fibrous or, in certain instances, bony tissue. Cassidy (1922-3) has reported a striking example of scorbutic infantilism which showed the typical clinical picture. A boy of 18 had never eaten either fruit or vegetables. He first developed scurvy at the age of 13, with haemorrhagic swellings in the muscles of the legs, and was unable to walk. Subsequently, under antiscorbutic treatment, he improved, but soon reverted to his favourite diet. At 18 there was striking retardation of growth and development, and the changes of puberty had not appeared. He was bed-ridden and, although then able to move his feet, his lower limbs were otherwise powerless. Radiological examination showed the presence of remarkable conical structureless opacities at the ends of the diaphyses of the long bones, most evident around the knee. These opacities suggested previous haemorrhages, in which bony changes had taken place. Otherwise there was extreme rarefaction of the skeleton and in both femora evidence of recent fractures. This case illustrates the severe retardation of growth and development which follows prolonged vitamin C deficiency.

(c) COELIAC RICKETS

The arrest of development in long-standing cases of coeliac disease is often very striking. Smallness of stature is associated with general retardation of development. The state of cachectic infantilism, however, is complicated by the presence of rickety changes in the metaphyses which interfere with the growth of the long bones and produce a short-limbed, disproportioned skeleton. Radiologically, the development of the epiphyseal centres of ossification corresponds to the size of the individual. The metaphyses are swollen and cupped as in rickets and the epiphyseal line becomes irregularly serrated. Osteoporotic changes appear sooner or later in the shafts. These rickety changes can be attributed to vitamin D deficiency, the result probably of malabsorption or possibly of re-elimination from the intestines of fat and the dietary restrictions which have to be imposed on the individual.

Clinically the general appearance is striking and very similar to that in hypopituitary infantilism except for the presence of rickety deformities. Dwarfism is pronounced and the appearance of the secondary sex characters may be delayed until the age of 20 or later. In the majority of cases a history of fatty stools provides the clue to the nature of the underlying condition.

(d) RENAL DWARFISM

Renal dwarfism from chronic interstitial nephritis in childhood provides a very similar picture. It differs fundamentally from the other types in being due to the action of retention products—that is, inadequate excretion of phosphates by the kidneys and consequent interference with calcium metabolism. The underlying renal

condition is in many cases congenital. In fact, these children often show signs of renal inadequacy from birth, and as they grow older the bony deformities appear: the enlarged radial epiphyses, deformities of the long bones, and, later, genu valgum are characteristic of this condition. The urinary findings are similar to those of chronic interstitial nephritis in the adult, and at post-mortem examination the kidneys have been found to show varying degrees of granular atrophy.

Growth is slow and development severely retarded, and the presence of "rickety" changes at the metaphyses through early years produces a disproportioned short-limbed skeleton. Radiologically, the metaphyseal changes are difficult to distinguish from those of simple rickets, and a typically cupped and irregularly serrated metaphyseal line occurs. Mental development is unaffected, but the appearance of the secondary sex characters and the changes of puberty are much delayed. Perhaps the majority of cases of renal dwarfism do not survive until the age of puberty.

3. Endocrine Infantilism where Arrested Development Results from Glandular Deficiency

(a) HYPOPHYSEAL INFANTILISM

An immense step forward in our knowledge of hypopituitary dwarfism and infantilism was made when it was shown that the removal of the anterior pituitary in young animals, notably puppies, produced arrest of growth and sex development. The testes remained in an infantile condition and the secondary sex characteristics failed to appear. The arrest of growth was particularly noticeable in the long bones, which were unable to increase their shaft length. Histological examination showed lack of formation of cartilage cells in the growth cartilages. Radiologically, there was retarded osseous development and delayed epiphyseal fusion far beyond the normal time.

The most clearly understood pathological states causing hypopituitary infantilism are vascular lesions or tumours. Little is known, from the pathological aspect, of functional exhaustion of the anterior lobe. Of the vascular lesions Simmonds has described a syndrome, which now bears his name, of infantilism, cachexia, and defective growth from ischaemic necrosis of the anterior pituitary, the result of infarction generally from infective or acute illness. The deficiency syndromes in childhood due to compression from a tumour are those associated with the parhypophyseal and suprasellar cysts. As a rule the clinical picture is characterized by infantilism, dwarfism, and obesity, the syndrome of dystrophia adiposogenitalis described originally by Fröhlich. Whilst there is no doubt that cachexia usually accompanies the lesion described by Simmonds (1918) and adiposity the tumour syndrome associated with Fröhlich's (1901) name, on clinical as opposed to pathological grounds this nomenclature appears to be too rigid, for cachexia is sometimes found with parhypophyseal cysts and typical cases of Fröhlich's syndrome without any tumour. The difficulty in keeping to this nomenclature for clinical cases lies in the varying metabolic disturbances. The relation of the latter to the pituitary is not yet understood, and it would seem better to classify cases as hypopituitarism with or without tumour. In adults compression of the cells of the anterior lobe may be due to another type of tumour, the pituitary chromophobe adenoma; but such adenomata only occasionally develop before the end of the second decade—that is, before the developmental stage has passed—and consequently are not often associated with the hypopituitary syndrome of dwarfism and infantilism.

The growth defects in hypopituitarism are characteristic in cases antedating the second dentition. There is a striking disproportion of the skeleton, and the limbs are short in comparison with the trunk—that is, an infantile type of skeleton is retained. If, on the other hand, hypopituitarism does not develop until after the second dentition, when the skeleton has reached a more mature type, the same degree of disproportion will obviously not be found. There is, however, a tendency for the growth of the long bones to be affected more than that of the trunk, and evidence of this will be found in skeletal measurements. Radiological examination will show characteristic delay in the appearance and development of the ossification centres. The epiphyseal disks remain ununited beyond the usual time. This growth defect, associated with sex infantilism, normal intelligence, and mental faculties, gives the typical picture of hypopituitary infantilism. Adiposity when present tends to be limited to the trunk, abdomen, hips, and proximal segments of the limbs, the subcutaneous deposits in the distal segments being scanty and the extremities spared. In hypopituitarism due to parhypophyseal cysts neighbourhood symptoms are present and radiological examination of the sella turcica gives a characteristic picture. The fossa is usually of the flat and open type, with erosion of the clinoid processes. Suprasellar or intrasellar shadows may be observed, and indicate areas of calcification in the cyst. The calvarium often presents characteristic features—a "beaten-silver" appearance due to convolitional pressure and thinning from increased intracranial tension.

(b) HYPOTHYROIDIC INFANTILISM

Until recent years the thyroid has been regarded as the focal centre of pathogenesis in hypothyroidic dwarfism and infantilism. Varying grades of atrophy of the gland are found. Lately, however, it has been shown that the thyroid is under anterior pituitary control, so that in hypothyroidic states the possibility of a primary anterior pituitary pathogenesis must be borne in mind. It has been shown, for instance, that pituitary ablation is followed by hypothyroidism and that the anterior pituitary secretes a thyrotropic hormone which exercises its function through the mediation of the thyroid. One must bear in mind, therefore, the possibility of an anterior pituitary factor in hypothyroidic infantilism, and that the latter may be due to inadequate supplies of thyrotropic hormone.

In untreated congenital athyroidism dwarfism is pronounced; the long bones are affected more than the trunk and a disproportioned short-limbed skeleton results. The ossification centres are late in making their appearance in the cartilaginous epiphyses, and, when they do, they develop slowly. Growth may come practically to a standstill in untreated cases and the epiphyses may remain largely cartilaginous till the end of life. These changes can be demonstrated radiologically. Histologically, in the growth cartilages cell proliferation is much diminished and there is a want of orderly arrangement of the cells. Rounded groups of cells tend to occur and the normal palisade arrangement is not present. Some calcification of the matrix is seen, but its depth depends on the degree of cell proliferation (Knaggs, 1928-9).

In post-nasal hypothyroidism, a condition to which the term myxinfantilism has sometimes been applied, the resulting skeletal defects depend to a large extent on the age at which the thyroid defect develops. If it antedates the second dentition and the condition remains untreated a disproportioned skeleton with short limbs is found. In cases developing later and after the second dentition

retardation of growth in the long bones and skeletal disproportion are less pronounced. Disturbances in dentition go hand in hand with those of bone growth, and retardation of sex development in untreated cases is striking. The appearance of the secondary sex characters and changes of puberty may be indefinitely delayed. This picture with its associated mental retardation and myxoedematous skin changes is quite characteristic. The rate of growth and development when thyroid treatment is first instituted is often remarkable, but when once leeway has been made up further progress is on more normal lines.

(C) HYPOGONADAL INFANTILISM

(i) *From Pre-pubertal Castration.*—The effects of early castration in young animals are well known, but clinical reports of individuals, at any rate of the female sex, who have been castrated in juvenescence are extremely rare, if any exist. The details of such a case will therefore be of interest. Castration in adolescence results in characteristic changes in the skeleton. The growth of the long bones persists beyond the usual time, resulting in abnormal length of the limbs in comparison with the trunk. Such castrates usually grow tall, but, in the present instance of early castration, striking retardation was found. This patient, aged $19\frac{1}{2}$ at the time of my observations, had had both ovaries removed at the age of $9\frac{1}{2}$ for bilateral multilocular cystadenomata. As a result of the operation the changes of puberty did not occur and the secondary sex characters failed to develop. Her height at $19\frac{1}{2}$ was only 5 ft. $1\frac{1}{2}$ in., the measurement from symphysis pubis to vertex being $27\frac{1}{4}$ in. and from symphysis pubis to soles $34\frac{1}{4}$ in.—that is, the length of the lower limbs was greatly in excess of that of the trunk. A remarkable feature of this case was the very great enlargement of the sella turcica, which was shown by radiological examination to be three to four times the normal size. No changes were found in the fundi or visual fields. Radiological examination of the carpus showed that ossification centres were present and of a more or less normal degree of development, though it should be remembered that the patient was approaching this state at the age the operation was performed. The epiphyseal junctions, however, at the time of observation, had remained unfused. A modified Aschheim-Zondek test gave a positive result, indicating the presence of gonadotropic substances in the urine. Does it not seem reasonable to assume that the growth defect in this patient, unusual after castration, was due to a failure of the normal secretion of growth hormone from the eosinophil cells of the anterior lobe? In view of the enlargement of the sella turcica noted in the radiograph the hypothesis could be advanced that an overgrowth of the "castration" cells of the anterior lobe had interfered with the function of the growth cells. An overgrowth of "castration" cells is consistent with the findings in the pituitary after castration in animals.

(ii) *Hypogonadal or Eunuchoid Infantilism.*—Under this heading I will also quote a case which illustrates another most unusual condition—a combination of eunuchoidism, infantilism, and dwarfism. This patient, a male, was aged 34 at the time of my observations. He was very small, and wizened in appearance. His height was only 4 ft. $3\frac{3}{4}$ in., the measurement from symphysis pubis to vertex being $21\frac{1}{2}$ in. and from symphysis pubis to soles $30\frac{1}{4}$ in.—that is, the length of the lower limbs was greatly in excess of that of the trunk, as in eunuchoidism. Neither testicle was palpable in the scrotum or inguinal canal, and there was a complete absence of facial, pubic,

and axillary hair. The skin changes and distribution of the subcutaneous fat deposits were also typical of eunuchoidism. Radiography showed that the epiphyses of the long bones were still ununited. The clinical features suggest that the primary lesion is gonadal rather than anterior pituitary. The condition differs from hypopituitary infantilism in the excessive length of limb as compared with the trunk, and from true eunuchoidism in the defect in stature. In ordinary forms of eunuchoidism the individual is of normal stature or tall, a feature which is ascribed to the persistence of growth in the presence of ununited epiphyses. The persistence of growth could be attributed to eosinophil activity in the anterior pituitary. It seems likely that in this diminutive and infantile individual the defect of growth may have resulted from a secondary eosinophil hypoplasia. There were no changes in the sella turcica, nor was there any evidence of pituitary tumour as judged by examination of the fundi and visual fields. On the other hand, the general features and bodily dimensions of eunuchoidism strongly suggest a primary hypogonadal aetiology.

Other Forms of Infantilism

From time to time other forms of infantilism have been described, and although, if the above classification is adopted, it seems reasonable to include them in one or other of the categories, they should be mentioned separately for the sake of completeness.

So-called cardiac dwarfism and infantilism may be cited as an example. It would be generally agreed that severe cardiac lesions may cause interference with growth, and congenital abnormalities in particular are sometimes associated with arrest of development. On the other hand, congenital heart disease is known to be consistent with average development, and even when retardation does occur it is difficult to be sure that both cardiac and general maldevelopment are not part of a more widespread abnormality of the germ plasm. Mitral stenosis of rheumatic origin has been regarded as a cause of dwarfism. That it cannot be a common cause is evidenced by the number of children with it who grow and develop satisfactorily. There is certainly little evidence that cardiac lesions of this type produce any serious arrest, for the changes of puberty and secondary sex characters usually develop in a normal way. In fact, in cases which have been reported the defect has usually been one of stature rather than of general and sexual infantilism.

Some reference is also necessary to the condition described by Gandy (1906) as retrograde infantilism, in which there is found a regression of sex characters and functions towards the infantile state. The cases described have almost invariably had their onset in adult life, so that skeletal dimensions and proportions have not been affected. In the light of recent knowledge it would seem that some of these cases, at any rate, fall into the category of late eunuchoidism, in which a late atrophy of the sex glands occurs. Others, judging by the presence of hypothalamic symptoms, have probably been of hypopituitary origin. In some instances thyroid atrophy has been reported, and here the gonadal defects have probably been secondary to hypothyroidism. So-called retrograde infantilism, therefore, seems to include a number of different conditions. In any case, as the skeletal dimensions are normal it does not seem justifiable to place them in the category of true infantilism.

Finally, infantilism in association with Addison's disease has been reported in one striking case by Morlat (1903) and in a somewhat similar one by Apert (1933)

under the tentative heading of suprarenal infantilism. The latter points out that bilateral tuberculosis of the suprarenal cortex causing Addison's disease is rare in childhood, and such lesions, if they do occur, usually prove fatal before infantilism can develop. In the two cases mentioned dwarfism and infantilism were associated with pigmentation, asthenia, anorexia, and wasting. In Morlat's case, though the stature was small, the proportions of the head, trunk, and limbs were those of an adult. In the scheme of classification adopted in these lectures these cases could therefore reasonably be grouped in the category of cachectic infantilism.

A survey of the field of clinical and pathological abnormalities of growth and development shows that we are concerned with the influence and interaction of four groups of factors, nutritional, infective, endocrine, and hereditary, in the production of two different groups of phenomena—alterations in bodily size on the one hand, and variations in the time factor of development on the other. Abnormalities of stature such as gigantism and dwarfism and abnormalities of development, precocity and infantilism, are clearly associated closely, and occur together in a number of different combinations.

Our knowledge of some of the factors which modify growth and development is still very deficient. The influence, for instance, of over-nutrition and hypervitaminosis is not understood, though the effects of under-nutrition, hypovitaminosis, and infective disease in limiting growth and retarding development are becoming more clearly defined.

Our understanding of the endocrine factors has advanced in the last decade. Experimental work with the anterior pituitary growth hormone has shown its capacity for stimulating bodily growth, and its effects on the growth cartilages and general bodily tissues can be appreciated from clinical observations in hyperpituitary gigantism and hypopituitary dwarfism. The equally important influence of the sex hormones in permitting or limiting longitudinal growth by controlling the time of fusion of the epiphyses can also be judged from the various pathological conditions in which sex hormone production is disturbed. The sex hormones appear to produce senescence in growing cartilage, a maturing effect comparable to that seen elsewhere in the body. It is only necessary to remind ourselves that sex hormone production is primarily under anterior pituitary control to realize how important in growth and development is the anterior pituitary body. Further knowledge with regard to the site of production of the anterior pituitary sex hormones, and histological evidence as to the cell counts in the anterior pituitary in different pathological states of growth and development, should do much to help our understanding of these conditions.

The influence of heredity in growth and development is not within the scope of these lectures, but it is evident that hereditary influences have played an important part in a number of the conditions we have been studying. The importance of genetic factors in determining endocrine constitution can only at present be surmised, but it is in this field that there would appear to lie the explanation of many of the variations of growth and development met with in everyday experience.

REFERENCES

- Aldrich, C. A. (1936). *J. Pediat.*, **8**, 381.
 Apert, E. (1933). *Infantilism*, p. 41, London.
 Bailey, P., and Davidoff, L. M. (1925). *Amer. J. Path.*, **1**, 185.
 Barlow, T. (1883). *Med. Clin. Trans. Lond.*, **66**, 159.
 Brailsford, J. F. (1929). *Amer. J. Surg.*, **7**, 404.
 Brissaud, E., and Meige, H. (1895). *Rev. Scient.*, **3**, 330.
 Broster, L. R., and Vines, H. W. C. (1933). *The Adrenal Cortex*, London.
 Cassidy, M. (1922-3). *Proc. roy. Soc. Med.*, **16** (Clin. Sect.), 16.
 Cushing, H. (1912). *The Pituitary Body and its Disorders*, Philadelphia.
 — (1927). *British Medical Journal*, **2**, 48.
 Evans, H. M. (1933). *J. Amer. med. Ass.*, **101**, 425.
 — (1935). *Glandular Physiology and Therapy*, p. 49, Chicago.
 — (1935a). *Ibid.*, p. 47.
 — (1935b). *Ibid.*, p. 53.
 — (1935c). *Ibid.*, p. 49.
 — and Long, J. A. (1921). *Anat. Rec.*, **21**, 62.
 Fell, H. B. (1928-9). *Arch. exp. Zellforsch.*, **7**, 390.
 Ford, F. R., and Guild, H. (1937). *Johns Hopk. Hosp. Bull.*, **60**, 192.
 Fröhlich, A. (1901). *Wien klin. Rundsch.*, **15**, 883.
 Gandy, C. (1906). *Bull. Soc. méd. Hôp. Paris*, **23**, 1226.
 Gilford, H. (1911). *The Disorders of Post-natal Growth and Development*, London.
 Green, H. N., and Mellanby, E. (1928). *British Medical Journal*, **2**, 691.
 Grollman, A. (1936). *The Adrenals*, London.
 Haldeman, K. O. (1927). *Arch. Neurol. Psychiat.*, **18**, 724.
 Hanson, A. M. (1936). *Minnesota Med.*, **19**, 1.
 Harris, H. A. (1926). *Arch. intern. Med.*, **38**, 785.
 — (1933). *Bone Growth in Health and Disease*, p. 89, London.
 — (1933a). *Ibid.*, p. 53.
 — (1933b). *Ibid.*, p. 10.
 — (1933c). *Ibid.*, p. 145.
 — (1933d). *Ibid.*, p. 34.
 — (1933e). *Ibid.*, p. 157.
 Harrison, R. G. (1935). *Heteroplastic Grafting in Embryology*, Baltimore, p. 116.
 Hess, A. F. (1930). *Rickets*, London.
 Hopkins, F. G. (1912). *J. Physiol.*, **44**, 425.
 Horrax, G., and Bailey, P. (1925). *Arch. Neurol. Psychiat.*, **13**, 422.
 — (1928). *Ibid.*, **19**, 394.
 Hunter, J. (1837). *Works*, **4**, 317, London.
 Knaggs, R. L. (1926). *Inflammatory and Toxic Diseases of Bone*, Bristol.
 — (1928-9). *Brit. J. Surg.*, **16**, 370.
 Krabbe, K. H. (1921). *N. Y. med. J.*, **114**, 4.
 Launois, P. E., and Roy, P. (1904). *Étude Biologique sur les Géants*, p. 478, Paris.
 Lee, M. O., and Schaffer, N. K. (1934). *J. Nutrit.*, **7**, 337.
 Lorain (1871). Lettre Préface la Thèse de Paris de Faneau de la Cour, F. V.
 McCollum, E. V., and Simmonds, N. (1927). *Newer Knowledge of Nutrition*, third edition, New York.
 Mann, H. C. Corry (1926). Medical Research Council, Special Report Series, No. 105, London.
 Massalongo, R. (1892). *Rif. med.*, **8**, 74.
 Mellanby, E. (1921). Medical Research Council, Special Report Series, No. 61, London.
 Mitchell, H. H. and Beadles, J. R. (1929-30). *J. Nutrit.*, **2**, 225.
 Morlat, A. (1903). *Thèse de Paris*, No. 194.
 Morquio, L. (1929). *Arch. Méd. Enf.*, **32**, 129.
 Novak, E., and Long, J. H. (1933). *J. Amer. med. Ass.*, **101**, 1057.
 Ollier (1899). *Bull. Soc. Chir. Lyons*, **3**, 22.
 Parsons, L. G. (1936). *British Encyclopaedia of Medical Practice*, London, **1**, 135.
 Robertson, T. B. (1923). *Abt's Pediatrics*, Philadelphia, **1**, 445.
 Rössle, R. (1917). *Ergebn. allg. Path.*, **18**, Abt. 2, 677.
 Rowlands, R. P., and Nicholson, G. W. (1929). *Guy's Hosp. Rep.*, **79**, 401.
 Sacchi, E. (1895). *Riv. sper. freniat.*, **21**, 149.
 Scottish Board of Health (1928). *British Medical Journal*, **1**, 140.
 — (1929). *Ibid.*, **1**, 23.
 Simmonds, M. (1918). *Dtsch. med. Wschr.*, **44**, 852.
 Stewart, C. A., Bell, E. T., and Roehlke, A. B. (1936). *Amer. J. Cancer*, **26**, 144.
 Stratz, C. H. (1922). *Der Körper des Kindes*, Stuttgart.
 Tandler, J., and Gross, S. (1910). *Arch. Entwicklgs-mechn.*, **29**, 290; *ibid.*, **30**, 236.
 Thursfield, H., and Lightwood, K. C. (1930-1). *Proc. roy. Soc. Med.*, **24**, 564.
 Wegner, G. (1870). *Virchows Arch.*, **50**, 305.
 Wright, A. Dickson (1930-1). *Proc. roy. Soc. Med.*, **24**, 283.

The Committee of Award of the Commonwealth Fund Fellowships has made a number of appointments to fellowships tenable by British graduates in American universities for the two years beginning September, 1937. These fellowships are offered by the Commonwealth Fund of New York, of which Mr. Edward S. Harkness is president. Awards in medicine have been made to D. M. Douglas, M.B., Ch.B. (of the University of St. Andrews) to Minnesota University, and to J. D. Spillane, M.B., B.Ch. of the University of Wales) to Columbia University.