in hospital practice. The incidence of rheumatic fever was found by Coombs to be 5 girls to 4 boys; yet the incidence of mitral stenosis in females must be in the region of 5 to 1 male. This is due to the less active life led by girls, who are seldom permitted to be out of range of the maternal eye; they sit and knit and acquire obstructive stenosis and provide most examples of auricular fibrillation in the forties. Her brother with the same diseases escapes to the playground and, despite restrictive advice, ensures that the cusps of the mitral valve are kept well apart by the vigour of his exercise, thereby securing a greater volume in flow and growth of the left ventricle. At the same time he prolongs his life.

Convalescent heart homes are not the nurseries for the cultivation of mitral stenosis that they used to be; nevertheless the introduction of the sedimentation rate has tended to increase bed rest in these cases, and mitral stenosis is actively encouraged once again. Because the sedimentation rate is increased, active endocarditis is presumed with often no justification; it might well be tonsil, sinus, or other infection. Immobilization and fibrosis are inseparable bed-mates and are no companions for growing hearts.—I am, etc.,

London, W.1.

BRUCE WILLIAMSON.

Diet and the Nation's Health

SIR,—As a humble member of the profession I note with dismay that Dr. S. K. Westman (Dec. 6, 1947, p. 926) regards the Hunterian Society and frivolity as being synonymous. A press précis can be misleading; it is probable that he has now had access to the full discussion and is a wiser man. The comparative statistics given in his letter beg many questions, but this outstanding and alarming fact is borne out—namely, the famine babies of 1919–21 responded, in the absence of adequate food for the body, to a dangerous and devastating dietary of distorted mental and moral feeding. They contributed to the downfall of their country and plunged the civilized world into chaos and economic disaster. Was the subsequent prowess of Hitler Youth in war the triumph of starvation or the thrust behind a fanatical and insatiable ambition for power? In their quest for *Lebensraum* was it not perhaps a bigger dining-room they subconsciously sought?

While I admire Dr. Westman's personal victory of mind over matter, I would remind him that his unusual menu for strength is in short supply in this country. It would be regrettable if our rowing blues were deprived of their sport because they had nibbled their oars and sucked the substance of the boats while in training. The whittling down of our larder has already caused deterioration in our morale and bred dishonesty in many a decent home. The man in the street does not need a learned society to point that out to him. The desire for a high standard of food and health is not actuated by selfishness and greed on Britain's part, but in her upholding the terms of the Atlantic Charter—"that all the men in all the lands may live out their lives in freedom from fear and want." We want butter in our guns and so avoid any continental European or Asiatic parallels of hunger and discontent.—I am, etc.,

Dursley, Gloucestershire.

ISABELLE B. S. FAYLE.

SIR,—May I add a note to Dr. F. M. R. Walshe's eloquent letter (Dec. 13, 1947, p. 971) emphasizing the importance of subjective impressions in the formulation of evidence? Sir Jack Drummond's argument seems to me to suffer from a fallacy of misplaced concreteness. He was mistaking the modern concept of nutrition for a fact, and forgetting that it is a theory still to be tested by experience. He maintained that, because we were getting all the items of diet which the modern theory of nutrition held to be necessary for our health, therefore we must be healthy. But this theory may prove to be inadequate, just as those theories of nutrition in vogue before the discovery of vitamins proved to be inadequate. It may be that from these subjective impressions which he scorns to notice new facts will emerge compelling the theory to be recast.

Of course, subjective impressions are dangerous data. The field they grow in may be thickly sown with the tares of wishful thinking. Disapproval of the Government's activities may easily squeeze itself through the gaps in our sentiment of scientific impartiality and emerge as subjective impressions concealing a grudge. This is what happened, no doubt, to those

physicians who felt that their patients were suffering from the effects of calcium in the bread at a time when the calcium was present only in their imaginations. Much laborious and honest threshing is necessary to winnow away the chaff of adventitious percepts from the grain of a new impression. But it is the impressions which supply us with new facts and which therefore ought never to be neglected.—I am, etc.,

Wallingford, Berks.

E. H. HARE.

A Sign of Carcinoma

SIR,—Carcinoma of the breast frequently presents one of the stiffest problems in surgical diagnosis, and all surgeons of experience approach these cases with humility, which is understandable when pathologists tell us that never a year passes by but they see at least one breast removed in the Halsted fashion for mastitis and more than one local removal of a carcinoma in error for a benign condition.

The simple manoeuvre employed to obtain the accompanying photograph I have found after long experience to be a great help in diagnosis. If the arms are raised above the head as high as possible all the usual signs of carcinoma are increased and, if not already present, are frequently made manifest in the most surprising manner. This applies to the level of the nipples, retraction of the nipple, puckering of the skin, and even peau d'orange. In addition the palpability of the tumour to the flat hand is easier to elicit when the underlying muscles are on the stretch.

The case illustrated was referred to the out-patient department as a case of chronic mastitis, and with the arms hanging by the side the breasts were indistinguishable on inspection and the tumour was vague to the finger and impalpable to the palm of the hand. The raising of the arms above the head instantly made the diagnosis of malignancy indubitable. With chronic mastitis some puckering of the skin may be produced, but it is generally a diffuse rippling over a fair area rather than a well-defined depression such as is seen in the photograph. Feeling sure that others have made this discovery before, this simple test is nevertheless put forward in the hope that wider publicity for it will help in what must be regarded as one of the most responsible and difficult diagnoses which come the way of the clinician.—I am, etc.,

London, W.1.

A. DICKSON WRIGHT.

Staphylococcal Infection due to Penicillin-resistant Strains

SIR,—In her very interesting article (Nov. 29, 1947, p. 863) Dr. Mary Barber draws attention to the increasing incidence of strains of *Staphylococcus pyogenes* that are grossly resistant to penicillin. We have been testing between April and December, 1947, all cultures of *Staph. pyogenes* for penicillin-sensitivity, using the penicillin-ditch method with 1, 5, and 10 units of penicillin per ml. Secondary cultures had been tested throughout. Out of a total of 115 cultures of *Staph. pyogenes*, 9 (7.8%) were penicillin-resistant.

Type of Infection.—Axillary abscess, 1; otitis media, 2; furunculosis, 1; dermatitis, 1; aplastic anaemia, 1 (blood culture). The 3 other penicillin-resistant strains were grown from the throat (2) and nose swabs (1) of carriers. Considering only the strains from infected cases, 6 (5.2%) were penicillin-resistant.

In Czechoslovakia, owing to the comparative shortage, the use of penicillin, especially in the form of ointments, lozenges, etc., has not been as widespread as in Great Britain. None of the cases examined had received penicillin prior to the bacterio-