The quantitative and qualitative similarity of this phenomenon in both conditions suggests a common cause for the eosinopenia.

Adrenaline does not affect the eosinophilic content of bone marrow, and the latter accordingly is not the site of eosinophilic retention.

Eosinopenia due to sufficiently high insulin dosage indicates an ability of the organism to meet the emergency of hypoglycaemia by secreting adrenaline, and this, if repeated several times, might explain the improvement reported as being effected in allergic conditions through treatment with insulin hypoglycaemia.

REFERENCES

- Bertelli, G., Falta, W., and Schweeger, O. (1910). *Z. klin. Med.*, **71**, 23.
 Cannon, W. B., McIver, M., and Bliss, S. W. (1924). *Amer. J. Physiol.*, **69**, 46.
 Eppinger, H., and Hess, L. (1909a). *Z. klin. Med.*, **66**, 345.
 (1909b). *Ibid.*, **68**, 205.
 Godlowski, Z. (1946). *British Medical Journal*, **1**, 717.
 Heilbrun, G., and Libert, E. (1939). *Endocrinology*, **25**, 354.
 Naegeli, O. (1931). *Blutkrankheiten und Blutdiagnostik*, 5th ed. J. Springer, Berlin.
 Randolph, Th. G. (1944). *J. Allergy*, **15**, 89.
 Spangler, R. H. (1925). *Arch. intern. Med.*, **36**, 779.

THE CONVULSIVE PROPERTIES OF THIOPENTONE

BY

R. L. WYNNE, M.B., M.R.C.S., D.A.
Anaesthetist, Birkenhead General Hospital

The case of frank convulsions under thiopentone anaesthesia recorded by Bodman and Farr (1947) is of interest inasmuch as such phenomena in slighter form must have been widely observed but hitherto seem to have excited little or no remark in the literature other than an occasional reference to "tremors."

Thiopentone is a drug not distantly related to others—e.g., 1-3-dimethylbutylethyl barbituric acid, and 2-1-methylcrotyl-2-methylallyl-, 1-methylbutyl-2-methylallyl-, and crotyl-*n*-butyl thiobarbituric acids—which have been

described by Swanson and Fry (cited by Adams, 1944) as convulsants having no sedative properties.

The "tremors," or minor convulsive manifestations, of thiopentone occur shortly after consciousness is lost, and consist of: (1) Pronatory spasm of the arm, either spontaneous or readily elicited by pricking, or even simply by withdrawal of the syringe needle. (2) This in some cases progresses to strong clonic contractions of the pectoral and neighbouring muscles, a sign which might perhaps be descriptively named the "shudder reflex." (3) Other muscles, especially those of the glottis and diaphragm, may be involved. The phenomena are in all cases most strongly grouped round the musculature associated with respiration, normal or forced. (4) In some cases the spasm is of tonic rather than clonic nature. These are harder to recognize, since the predominant sign is a cessation of respiration, which more often occurs as indicating a relaxation of the musculature consequent on a momentary overdose. The spastic type of respiratory arrest may, however, be differen-

tiated by its occurrence after a relatively small dose and almost invariably in the muscular type of patient, by observation of the rigidly held chest, and by the fact that it often disappears on rapidly administering a further small dose of the drug (though it is not recommended that this last feature be adopted as a routine test).

The following case is cited in illustration.

Case Report

A healthy gunner aged 33 was presented for anaesthesia following a fracture of both bones of the left leg, the date being Oct. 11. After "omnupon"—scopolamine he was given 1 g. of thiopentone intravenously, combined with gas-oxygen, for skeletal traction by a pin through the os calcis, and plaster. No untoward symptoms were noted. On Oct. 29 he was anaesthetized for replastering, again receiving omnupon—scopolamine and 1 g. of thiopentone. This time no inhalation was given, and a further injection of 0.5 g. of thiopentone was found necessary; otherwise there were no unusual features.

On Nov. 2 further manipulation was required and he was once more anaesthetized, on this occasion for some reason without any premedication. He was given 1.5 g. of thiopentone, and definite convulsive phenomena at once developed. Pronatory movements were followed by the "shudder reflex" described above, and this by convulsive inspiratory movements of the respiratory musculature. Respiration became shallow and rapid, the movements being about 60 a minute, and thus frequent enough to ensure adequate oxygenation. During the next five minutes the signs died away, and the patient thereafter progressed without further unusual signs.

On Nov. 5 another anaesthetic was required, and this time, being forewarned, he was given omnupon—scopolamine, and 0.5 g. of thiopentone was injected to see what the response would be. As soon as he became unconscious the same strong convulsive movements were once more observed, and no time was lost in administering gas-oxygen with ether. The administration was rendered difficult by laryngeal spasm, and not until a tracheal tube was passed was the anaesthesia free from difficulty. The patient thereafter progressed normally, and was not seen again.

Comment

This case illustrates other features associated with these properties of thiopentone: (1) The untoward signs were not at first elicited, but developed and got worse after several administrations of the drug during a comparatively short period, presumably indicating a progressive intolerance which might possibly be seen in other cases were their apparent tolerance subjected to the same test. (2) Though premedication was not in itself sufficient to prevent the signs, its absence on one occasion was associated with their appearance in fully developed form. (3) The condition was readily controlled by combination with other anaesthetic agents. (4) The condition each time occurred so early that it must be regarded as produced by the direct effect of the drug on the nervous system, and not by ketosis, carbon dioxide excess, or any of the other metabolic factors which have in the past been suspected as causative in other forms of convulsions.

My records contain several cases in which so similar a procession of events occurred that it seems needless to recount them in detail. Army experience taught me to regard the occurrence of any degree of spastic pronation during administration as a warning that the subject was unsuitable for uncombined thiopentone anaesthesia, and to add gas-oxygen inhalation in 25% mixture at the earliest opportunity. I always found that two to five minutes of this would ablate all convulsive phenomena and produce an even anaesthesia which could be managed by ordinary methods. Laryngeal spasm and the "total" respiratory spasm described above can usually be ablated by a further dose of thiopentone, but the "shudder reflex" is never thus amenable without the addition of other agents.

Cystoscopy or the passage of bougies under uncombined thiopentone anaesthesia has in my experience often produced difficulties resolved by the addition of gas-oxygen, and the inference seems to be that the sphincters may also be involved in a spastic process during such anaesthesia, which in these cases is also best combined with inhalation agents.

It is curious that these by no means rare signs should occur with a drug often used to combat convulsions; and in this connexion the hypothesis of Williams and Sweet (1944) seems to be pertinent. In considering anaesthetic convulsions in general, they adduce the probability of a constitutional factor, and suggest that in every case a convulsion is "primarily due to an inborn but latent epileptic liability . . . factors which arise during anaesthesia being merely precipitants of the convulsions." Encephalograms of patients known to have had convulsions during anaesthesia show irregularities supporting this hypothesis. Given the existence of such a type of patient, it should not be too much to suppose that the sudden addition of any potent drug to the blood stream might well act as a "precipitant."

Summary

The convulsant possibilities of thiopentone might be suspected, having regard to its near relationship to other strongly convulsant drugs.

The most common phenomena associated with these properties of the drug are: (a) pronatory movements of the arm, sometimes progressing to (b) strong shuddering movements, and even to (c) clonic spasm of other muscles, mostly those associated with respiration, either normal or forced. Tonic spasm of the glottis or of the entire thoracic musculature is not uncommon, and the sphincters also sometimes seem to be involved in the process.

The time factor suggests that no other exciting agent than the drug itself is responsible for their occurrence; but there is a possibility that an inborn convulsive tendency in the patient may predispose to their appearance.

The onset of convulsive phenomena is favoured by absence of premedication, uncombined thiopentone anaesthesia, and repeated thiopentone anaesthesia.

Convulsive signs are most likely to occur in muscular highly trained males leading an open-air life.

They are easily prevented or ablated by omnopon-scopolamine premedication and early addition of gas-oxygen in 25% or other suitable mixture.

REFERENCES

- Adams, R. C. (1944). *Intravenous Anesthesia*, p. 139. New York.
Bodman, R. I., and Farr, A. (1947). *British Medical Journal*, 2, 175.
Williams, D., and Sweet, W. H. (1944). *Lancet*, 2, 430.

GRANULOSA-CELL TUMOUR OF OVARY AS ACUTE ABDOMINAL EMERGENCY

BY

A. C. BREWER, F.R.C.S.

Assistant Honorary Surgeon, Liverpool Royal Infirmary

Granulosa-cell tumours of the ovary are usually recognized as a result of their effect on the endometrium, but like other ovarian neoplasms they may give rise to acute surgical complications.

Case Report

A female patient aged 48 years was admitted to hospital complaining of acute abdominal pain of some twelve hours' duration. She had had no previous illnesses. For the last three months she said she had "not been feeling too well," and for the last twelve days she had had a diffuse lower-abdominal pain of a moderately severe nature. There had been no definite colic. During the past five days the pain had become localized in the right iliac fossa and had become more constant; it radiated into both loins. Lying in bed

gave some relief. There had been no nausea and no vomiting. The bowels were regular and micturition normal. The menopause had occurred at the age of 44, and since that date the patient had had no vaginal bleeding. She had two children aged 18 and 12 years, and there had been one miscarriage thirteen months before the birth of her eldest child.

On examination the tongue was moist and had a white fur. The abdomen moved freely on respiration. Tenderness and muscle-guarding were present in the right iliac fossa. Vaginal examination revealed tenderness in both fornices, more marked on the right side. Pain was complained of on moving the cervix. The uterus was slightly increased in size. The condition was diagnosed as subacute appendicitis and operation advised.

A split-muscle incision was made. On opening the peritoneal cavity a considerable quantity of blood was evacuated. The exposure was considered to be unsatisfactory, so a further midline incision was made. About 1 pint (568 ml.) of blood and clot was evacuated from the pelvis. An ovarian neoplasm of moderate size was delivered, with blood clot extruding from two places in its capsule. Salpingo-oophorectomy was performed. The uterus and left ovary appeared to be normal. The patient had an uneventful convalescence.

Comment

The specimen consisted of a tumour measuring $9 \times 7 \times 5$ cm. and having a thick whitish capsule, which in places showed a yellowish colour owing to the tissue deep to it. It had ruptured at two points, and blood clot was protruding from these ruptures.

A portion of Fallopian tube was adherent to the tumour at one point. The cut surface showed a uniform whiteness, but with areas of haemorrhage. On section a dense fibrous capsule resembling ovarian stroma was seen. No follicular elements were present, the tumour consisting of sheets of oval cells lying in a loose stroma rich in blood vessels. In places the cells had an acinar arrangement, but nowhere was there follicular formation. In a few areas there were vacuolated cells resembling Call-Exner bodies. There was a considerable degree of necrosis and interstitial haemorrhage. The appearances were those of a granulosa-cell tumour of the diffuse type.

The granulosa-cell tumour is one of the solid gonadal tumours of the ovary in which the cell elements resemble the granulosa cells of the normal Graafian follicle and has the ability to secrete an oestrogenic hormone. The tumours are almost invariably unilateral, maintain the shape of the ovary within a firm fibrous capsule, and vary in weight from a few grammes to several kilograms. The cut surface is a soft whitish pink with yellow zones, sometimes partly cystic, and occasionally there are areas of haemorrhage. Histologically three main types are recognized according to the degree of glandular-tissue differentiation—follicular, trabecular, and diffuse—and as a general rule the least differentiated tumours show the least hormonal activity. In those tumours with oestrogenic properties the uterine endometrium is of the hyperplastic, non-secretory type with cystic dilatation of the gland tubules. Tumours showing evidence of malignancy are extremely rare. The pathology has been well summarized by Harvey, Dawson, and Innes (1939).

Although haemorrhage into a granulosa-cell tumour of the ovary is common, I cannot find any previous report of a case associated with so severe a degree of haemoperitoneum. The fact that the haemorrhage had been severe enough to burst through a capsule so thick as the one in this case shows the severity of the bleeding. There was no evidence of torsion, a complication that has been reported.

REFERENCE

- Harvey, W. F., Dawson, E. K., and Innes, J. R. M. (1939). *Edinb. med. J.*, 46, 256.