WEIGHT AND BLOOD PRESSURE

BY

R. A. MURRAY SCOTT, M.B.

Though information about the state of health of any section of the community is of particular value at the moment, few surveys of the nutrition of adults have been recorded. Some statistical records of haemoglobin levels are available, but little is known of the comparison, both in weight and in blood pressure, between age groups of people before the war and to-day.

A review has been made of the weights and blood pressures of the male members of an office staff spread over the North of England. The table shows the age distribution of the men examined in 1938-9, 1943, and 1946-7.

Age Distribution

Age Group	1938-9	1943	1946-7	
			Ex-Service	Non-Service
16-20	38	29		104
21-25	152	8	35	24
26-30	302	11	32	28
31-35	291	72	70	7
36-40	192	128	38	55
41-45	139	60	10	81
46-50	76	69		52
51-55	46	28		42
Totals	1,236	405	185	393

The weights were taken fully clothed, but as the examinations were evenly distributed through the years seasonal variation in the weight of clothes has been ignored. The measurements of blood pressure before the war were taken by one doctor, while those taken in 1943 and 1947 were measured on the same instrument by two doctors, one of whom examined over three-quarters of the men. The measurements made by these two doctors have been several times compared and found to be similar. In the 1938-9 group the average weight was 155 lb. (70.3 kg.). The average weight in 1943 fell to 151 lb. (68.5 kg.), and in 1946-7 it rose to 156 lb. (70.8 kg.). These averages do not in themselves give a true picture of what took place, as the proportion of men in the several age groups varied in each period—e.g., the largest number of examinations in 1938-9 were in the 26-40-year groups, while in 1946-7 the largest group is 16-20 years.

A more detailed exposition of weights by age groups is shown in Graph 1. The average weight in pounds is plotted against

GRAPH 1.—Average weight according to age groups.

five-year age groups. The number of men in each group can be seen from the above table. From the age of 21 to 40 the average weight in each 1946-7 group exceeds that in each comparable 1938-9 group, showing a most marked divergence of about 8½ lb. (3.8 kg.) at 26-30 years. The 16-20 age group is static in each period; so is the 41-50 group. In 1946-7, the 51-55 group is 5½ lb. (2.5 kg.) less than the 1938-9 counterpart.

Graph 2 shows the effect of splitting the 1946-7 series into ex-Service and non-Service components and leaving the 1938-9 series for comparison. This shows that from 21-35 years the ex-Service men give a higher average weight than the non-Service men, while the 36-40 group of non-Service men is higher than the ex-Service group. The number of men in each group is now too small to be completely significant, but the steady trend of ex-Service men in three consecutive age groups to be heavier than the non-Service men is suggestive. It is

interesting to note that of 52 men aged 46-50 years examined in 1943, the average chest measurement was 37 in. (92.5 cm.) and the waist 32½ in. (81.25 cm.), while of 52 men of this age group examined in 1947 the average chest measurement was the

GRAPH 2.—Average weight of ex-Service and non-Service men of the 1946-7 group compared with that of the 1938-9 group.

same, but the waist measurement was 1½ in. (3.75 cm.) greater. This can be compared with an increase in average weight of 7 lb. (3.2 kg.) of the 1947 group over that of 1943.

Graph 3 shows the rising lines of diastolic pressure of the increasing age groups of the three series of examinations. Through the whole of its length the 1938-9 line lies below those of 1943 and 1946-7. The 1943 line similarly lies above the other two. This grouping at once suggests that the war

GRAPH 3.—Average diastolic blood pressure.

strain of 1943, due to extra work and additional employment in A.R.P., fire-watching, and so on, coupled with the worries of bombing and the fear for relatives in the fighting line, caused a comparative rise of diastolic blood pressure at all ages. This increase has now fallen to a level roughly half-way between the wartime and pre-war levels. There is an interesting but unexplained dip in the 1938-9 and the 1946-7 lines at the 41-45-year group. Of more practical importance is the fact that the 46-55 groups of 1946-7 have a diastolic pressure averaging about 88 mm. Hg, which is some 5 mm. higher than that of the 1938-9 group.

Graph 4 records the average systolic pressures of the age groups. It shows positively the great average increase in pressure in those over 40 years in the wartime and post-war groups compared with pre-war. The post-war heights show

GRAPH 4.—Average systolic blood pressure.

little decrease on the 1943 range. The rise of average systolic pressure in the 21-25 group in 1946-7 compared with the 1938-9 group is thought to illustrate the nervousness of men at their first examination, as no rise occurs in the diastolic pressure.

Conclusions

Up to the age of 40 years the men examined in 1946-7 are on the average heavier than men of the same age in 1938-9. From 41 to 55 years there is little difference between the two periods.

The men between 21 and 35 years who have been in the Forces are on the average heavier than those who have not.

The average diastolic pressures of all 1946-7 age groups are above those of 1938-9, the maximum difference being 12 mm. Hg at 41-45 years.

The average systolic pressures of all 1946-7 age groups are above those of 1938-9, the maximum difference being 27 mm. Hg at 46-50 years.

The rise in blood pressure can be correlated with a recent increase in neurocirculatory disorders in the male staff over 50 years of age.

This sample survey of a section of the community is interesting and provides nutritional implications. Additional information of this type from other sections of the community would help to paint more clearly the picture of the health of the people of this country to-day. Perhaps it is available, but not published.

ROYAL SANITARY INSTITUTE CONGRESS INDUSTRIAL HEALTH DISCUSSIONS

A Health Congress was organized by the Royal Sanitary Institute at Harrogate from May 24 to 28. Delegates were present from several hundred municipal authorities and voluntary bodies, as well as from Government Departments, home and overseas. Dr. James Fenton and Dr. J. A. Ireland represented the British Medical Association. The Congress, over which Lord Inman presided, met in eight sections and at four special conferences. The president of the Conference of Medical Officers of Health was Dr. F. Hall, County Medical Officer, Lancashire. Dr. G. S. Wilson presided over the Section of Preventive Medicine and devoted his address to a description of the Public Health Laboratory Service, now established on a permanent footing under the National Health Service Act; on this subject he was followed by Dr. J. Greenwood Wilson and Dr. A. Leslie Banks. Dr. Andrew Topping presided over the Conference of Health Visitors, and urged, though with an awareness of the nursing shortage, that the health visitor should be a State-registered nurse.

Pneumoconiosis among Miners

One of the best sustained discussions in the Section of Hygiene in Industry, presided over by Prof. R. E. Lane, of the University of Manchester, was on the health of the coal-miner. Dr. J. B. Atkins, regional mines medical officer, Ministry of Fuel and Power, dealt with the problem of pneumoconiosis. In South Wales during the last two years there had been a steady improvement in incidence. In that area the problem had been so great as to strip the industry within three or four years of the best of its young man-power. By the end of 1945, 12,000 men, mainly skilled coal-face workers, out of a labour force of some 120,000, had been forced to leave the mines on account of this disease, thereby not only depleting the industry but seriously handicapping recruitment.

The general view was that the exceptionally high dust concentrations met with in South Wales, particularly in the mechanized mines before dust-suppression measures were taken, accounted for the wave of cases during recent years. Some concentrations were described by those who worked in them as being so thick that a man's lamp could not be seen at a distance of a few feet, and the choking effect was such that men frequently vomited their breakfast on starting work. The most effective attack on the problem had been made by ventilation, water infusion of the seam before mining, and the modification of mechanical mining to reduce the dust carried, for example, by wet boring and cutting and the incorporation of water sprays with pneumatic picks. Another line of investigation was by the finer analysis of jobs from the point of view of dust hazard. Regular x-ray examination of all men exposed would enable precautions to be taken when radiological signs of nodulation had developed, though Dr. Atkins pointed out that a single x-ray examination might be misleading if other clinical findings were not considered, because the appearance of x-ray changes due to dust might be masked by the development of focal emphysema. The first of a number of special light industry factories for the employment of men suffering from pneumoconiosis had recently been opened.

National Coal Board Medical Service

Dr. C. L. Cope, of the Scientific Department of the National Coal Board, described how research on pneumoconiosis was being actively pursued by the Medical Research Council unit at Cardiff, the team numbering nearly sixty. In this connexion reference should be made to the Goulstonian Lectures published in our last two issues by Dr. C. M. Fletcher, Director of the Pneumoconiosis Research Unit of the M.R.C.

Until the National Coal Board established an x-ray plant of its own, Dr. Cope continued, the contribution of the Scientific Department could only be the collection and co-ordination of information obtained from surveys and from records of dust concentrations. An advisory committee on nystagmus among miners which was formed originally by the British Colliery Owners Research Association was being actively maintained. Negotiations were proceeding for the appointment of a medical officer in the area of the small Kent coal-field to investigate miners' dermatitis. For some reason the incidence of dermatitis among miners in Kent was much higher than in any other part of the country.

A more difficult problem to tackle was the group of diseases not peculiar to miners. It was uncertain how miners' morbidity compared with that of the rest of the community. Chronic rheumatism was a good example of the type of condition on which more morbidity statistics were needed. It was held by some to be more common among miners.

Finally Dr. E. H. Capel, principal medical officer of the Board, discussed present and future arrangements. Under private ownership a few mines provided good medical services, but in general the industry lagged behind modern industries in this respect. Coal-mining was well known to be a dangerous occupation. During 1945 more than one miner out of every four sustained an accident or contracted an occupational disease necessitating more than three days' loss of work, a rate about seven times higher than that for industrial workers generally. During the past ten years the incidence of fatal accidents had fallen by about 30%, serious accidents had fallen slightly, but less serious accidents had risen by nearly 40%. Although the incidence of pneumoconiosis had fallen in South Wales since 1945, it had risen in other areas. Of other diseases, between 1,000 and 2,000 cases of nystagmus were certified annually; the incidence of occupational dermatitis was rising steadily, and chronic infections of the knee and elbow—the so-called "beat" diseases—remained at a high figure.

The medical service of the mines, as Dr. Capel saw it, would devote itself to selecting workers and assisting in their placement, controlling environmental conditions from the health point of view, and providing first treatment in accident and illness arising at work. One difficulty in organizing a medical service was that mines were small units. Of 956 mines, only 532 employed more than 500 men; 80 employed fewer than 100. If as a result of health measures absenteeism could be reduced from its figure of 12.5% last year to 11.5% it would mean an annual increase of about two million tons of coal.

Purity of Food

Several papers bearing on food inspection were read to the Congress. Mr. E. Dodsworth, chief sanitary inspector, Harrogate, discussed the adulteration of milk. He said that the Ministry of Health had suggested that out of every three samples of food and drink tested one should be of milk, milk being the ideal substance for the practice of adulteration. The familiar plea of defendants in milk cases that the appearance of the milk was due to the insufficient quantity and quality of the food given to the cows was contrary to the opinion of research workers hitherto, though worth re-examining, Mr. Dodsworth urged sharper penalties for offenders. The figures for adulteration remain obstinately high.

Ice-cream was also debated, and Mr. Morley Parry, chief sanitary inspector for Northampton, declared that owing mainly to the 1947 regulation which stated that all apparatus for the manufacture of ice-cream should be installed, maintained, and operated to the satisfaction of the local authority, the days of back-scullery manufacture of ice-cream were over. Mr. Parry was of opinion that there was no satisfactory equipment capable of complying with the regulations for heat treatment in their full form for those who manufactured less than