BRITISH MEDICAL JOURNAL

LONDON SATURDAY MARCH 20 1948

EPILEPSY*

BY

Sir CHARLES SYMONDS, K.B.E., C.B., D.M., F.R.C.P.

Physician for Nervous Diseases, Guy's Hospital

Now and again in the history of medicine the invention of a new instrument suddenly lights up the path of clinical observation so that many things which have been seen dimly become clear. Speculation is replaced by knowledge, which in turn shows the way to further speculation. Thought advances at a bound. At such a moment it may be forgotten that had it not been for all the patient and careful examination of what had already been seen this advance could never have been made, and that further progress must still depend upon clinical observation, helped, but not replaced, by the new method. The electro-encephalogram (E.E.G.) has thrown a light into many dark places on our way to the knowledge of epilepsy, but I would remind you at the outset that although the existence of an abnormal discharge of energy in the cerebral cortex coincident with epileptic attacks has only recently been proved by electrical methods, Hughlings Jackson seventy years ago conceived the idea that such a discharge must be the basis of epilepsy, and his views have been accepted by neurologists since that time as the basis of further speculation.

One of the ways in which the E.E.G. has helped the neurologist is in testing clinical hypothesis. For example, a man aged 30 came to me in 1947 with the story of two recent generalized epileptic attacks. There was a family history of epilepsy. The patient also complained of twitching in his eyes, which had gone on continuously since he could remember. This symptom was easily observed, for it occurred two or three times in every five minutes. The eyelids closed briskly but not so completely as to prevent one seeing an associated upward movement of the eyeballs. The whole episode lasted about a second. This complaint had been diagnosed as a tic or habit spasm, but it was not quite like a voluntary movement. Incomplete closure of the eyes as a voluntary movement is not associated with visible upward movement of the eyeball. Considered in its setting the symptoms suggested an epileptic discharge for this reason, and I made a note that I thought it must be epileptic though I had never seen a case with such frequent, circumscribed, and stereotyped motor expression. There, before the advent of the E.E.G., the observation would have ended in suspicion without proof. The electrical record showed a typical epileptic discharge with each episode of blinking. Confirmation in a case of this kind is of great value, for we have all many times seen little attacks in which the diagnosis has been doubtful, and with increasing experience we have all probably become less inclined to exclude epilepsy as the cause, because we have continued

to observe new and sometimes quite peculiar variants happening, at first alone, but later as the aura of an undoubted fit. Now we have at least the chance of getting confirmation, though a negative record would still leave doubt, for in these very brief myoclonic episodes there may be no detectable electric discharge.

Epileptic Variants

There are many variants of epilepsy of whose nature we can be sure without the aid of the E.E.G., but which because they depart so far from the "ordinary" attacks are apt to escape recognition. I suppose that loss of consciousness with or without convulsive spasm is the criterion most widely used in practice for the diagnosis of epileptic attacks. It is not sufficiently appreciated that in many attacks consciousness is not lost but disturbed. The sufferer may be aware of all that is going on around him, may hear and understand what is being said, but is for the time being unable, as he says, to collect his thoughts or reply to a question. There is in fact not loss of consciousness, but confusion, which may be of the mildest degree. As Gowers put it, the effect of an epileptic attack upon the stream of consciousness may vary from a complete interruption of its flow to a mere ruffling of the surface.

Apart from convulsive spasm there may be a momentary loss of power, and on the sensory side a great variety of abnormal feelings constituting the epileptic aura, which may occur without disturbance of consciousness. The chief characteristic of all these slight attacks is their suddenness and brevity. It would be impossible here to attempt any classification of epileptic variants. I shall select one or two for example.

First, the attacks called myoclonic. Anyone who has witnessed a generalized convulsion will have observed that sometimes after this is over there is occasional twitching of the limbs, which may continue for a while after the patient has regained his senses. Myoclonic twitching of this type often precedes an epileptic attack, and it may be the only symptom of an attack. Usually it is confined to the arms, and is especially apt to occur during the first hour or two after waking. There is no disturbance of consciousness, but sudden jerking movements of one or both arms which may interfere for the moment with dressing or shaving. The experienced patient knows when this happens that he may have a fit, but quite often the myoclonus passes off without this. The movements are fractions of a generalized convulsion. In rare instances such myoclonic twitching may be the only evidence of

*Annual oration to the Harrogate Medical Society, 1947.

epilepsy in a patient who never has any generalized convulsion or disturbance of consciousness, and may occur with great frequency

Next let me take a rare form of attack. Ramsay Hunt (1922) noted many years ago that a patient might sometimes fall without any disturbance of consciousness at all. Recognizing the epileptic character of these attacks, he called them "drop seizures." I have met with several examples of this kind of attack, and two families in which several members were afflicted. I recently saw a married woman of 62 whose only complaint was of falling attacks. These she had had since the birth of her first child when she was 26. They tended to occur in series at long intervals, and she had had five in as many weeks before I saw her. All were exactly the same. They occurred only when she was standing; without warning and without any impairment of consciousness she fell to the ground, as a rule forwards on hands and knees, often bruising herself. She was immediately able to get up without assistance. These attacks were never precipitated by laughter or other emotion, being thus distinguished from the well-known phenomenon of cataplexy. Her mother, who died at 65, had exactly similar attacks during the last ten years of her life, and an older sister also has them.

Focal Symptomatology

All these peculiar phenomena of the epileptic attack should, I think, be classified under the heading of focal symptoms because symptoms which are so specific must be related to the site of maximal epileptic discharge. This was the conception of Hughlings Jackson, derived from clinical observation long before modern methods had provided opportunity for precise observation. Seeking proof to support his theory, he found a case in which convulsive spasm in a single limb was associated with a scar in the appropriate area of cortex due to a depressed fracture. Hence the extraordinary notion that Jacksonian epilepsy and traumatic epilepsy were the same thing. Jackson used a case, which happened to be traumatic, to prove his argument that a lesion of a particular part would cause a particular kind of attack. To anyone who reads his papers it must be perfectly clear that this was all he meant. He was concerned with the localization of the lesion, not with its pathology. He was well aware that focal symptoms might occur in cases of brain tumour, or in what is called idiopathic epilepsy. Dr. James Collier used to say that the commonest cause of a Jacksonian attack was idiopathic epilepsy—a broad generalization which I believe is true when the idiopathic attacks are accurately observed, for there is very often an initial symptom of the attack which indicates a focal origin.

Penfield's (1946) observations in recent years have enriched our knowledge of focal symptomatology. With painstaking accuracy he has recorded the effects of electrical stimulation of different areas of cortex in patients subjected to craniotomy under local analgesia. His experiments have confirmed in the most striking way many of the conclusions reached by Hughlings Jackson from clinical observation. Jackson, for example, first described two kinds of attack associated with lesions of the anterior temporal lobe—the hallucinatory attack of taste or smell, and what he called the dreamy state. Penfield was able to produce both kinds of attack by stimulation of the anterior temporal cortex and of no other part of the brain. These attacks therefore have great localizing value, and the descriptions given by patients are characteristic. The smell or taste is always the same, always unpleasant, never quite like any normal taste or smell. The dream, which may include both hallucinations and illusions, has the dreamlike com-

bination of unreality and familiarity. In many cases both of organic disease and of idiopathic epilepsy Penfield was able to reproduce by stimulation attacks which the patient recognized as the same as those which happened spontaneously.

From what I have said you will appreciate that when we recognize a focal origin for the attack this does *not* imply an organic basis. Nevertheless the story of focal attacks in a patient with idiopathic epilepsy is usually different from that of a patient with a cerebral tumour, and the difference seems to depend upon the existence in idiopathic epilepsy of an abnormal state of the cortex as a whole, which makes it easier for the epileptic discharge to spread. The focal attack of idiopathic epilepsy may remain focal if the discharge is a slight one, but a more severe discharge will tend to spread. Consequently the patient often has minor attacks which consist of nothing but what we call an aura; but if there is a major attack consciousness is lost and convulsion, if it occurs, is generalized. By contrast, the discharge from an organic lesion is often restricted whatever its degree. The patient may have focal convulsions or sensory phenomena of considerable extent and rapid recurrence with no loss of consciousness or generalized convulsion. Such attacks recurring in stereotyped form usually indicate a coarse lesion—most often, of course, a tumour.

Modern Methods of Pathological Diagnosis

The preliminary sifting is made from the clinical data. Fits beginning under the age of 20 in those with a positive family history are idiopathic unless there is some strong clinical evidence to the contrary. The same is true of fits beginning in this age period which have continued for several years even though the family history is negative. But when we meet with a recent history of fits in an adolescent with negative family history and clinical examination, a radiograph of the skull is worth while. It is a simple procedure, and now and again will reveal some gross abnormality. For example, an otherwise healthy girl of 16 had a single fit with generalized convulsion and loss of senses. The onset was with spitting as if she wanted to get rid of something unpleasant—suggestive of a temporal lobe focus. Clinical examination was negative, and there was no family history of attacks. The radiograph showed a calcified tuberculoma in the right temporal lobe.

In adults the onset of the fits demands more detailed investigation. Past head injury, middle-ear or sinus infection, encephalitis, and cerebral syphilis are uncommon but important causes, and anyone who has lived in India should have the limbs x-rayed for evidence of cysticercosis. Radiographs of the skull should be done as a routine—calcification is quite common in the gliomas, which are the commonest of cerebral tumours. The E.E.G. is useful, and doubtless when apparatus is available will be considered indispensable. In 50% of idiopathic epileptics the record is of positive diagnostic value, and in another 25% it shows abnormalities which are suggestive. An organic lesion if fairly near the surface shows focal slow wave activity. An air encephalogram completes the investigation. Useful information can be obtained by the injection of 30 c.cm. by lumbar puncture—a procedure which as a rule upsets the patient only for a day or two. If epilepsy is symptomatic of a removable tumour (meningioma) there will very probably be a filling defect, though normal pictures are compatible with an infiltrating glioma. No one quite knows how many of the patients without apparent hereditary or other cause starting to have fits in adult life eventually prove to have tumours. It is commonly estimated at 50%.

Natratss (1943) has questioned this belief on the basis of a follow-up of 81 cases whose first attack was after the age of 40, finding that of 57 who could be traced 16 had died in less than twelve years, but only nine of proved organic cerebral disease. The course in the others, both dead and survivors, had been that of idiopathic epilepsy. My impression is that his figures, though small, are near the truth. A considerable number of idiopathic epileptics have their first attack after 30, and idiopathic epilepsy is a very common disease.

Causation

About the causation of idiopathic epilepsy we have much still to learn, but the E.E.G. has taught us a good deal about the hereditary factor. Its familial incidence has of course been long recognized, though in only about a quarter of the cases, and then curiously enough more often in siblings, aunts, and uncles than in either parent; but Lennox and the Gibbsses (1940) have recorded the following observations. They examined E.E.G. of 183 first-degree relatives of 94 patients with clinical epilepsy and found abnormal E.E.G.s in 60% as compared with 10% of 100 persons with no personal or family history of epilepsy. In 53 instances records were obtained from both parents, and in only 5% were both normal. Even more startling was the finding that in 35% both parents showed an abnormal E.E.G. Of the relatives with abnormal E.E.G.s only 1 in 3 had clinical epilepsy. Lennox and his collaborators conclude that the abnormal E.E.G., presumably associated with an epileptic tendency, may be inherited as a dominant trait. The risk of an epileptic offspring having epileptic children is great only if the mate also carries the trait. Thus if a person with epilepsy will choose a mate who has a normal brain rhythm his chance of having any offspring with epilepsy is greatly reduced. His chances of normal offspring are greater than those of parents both of whom have no present or family abnormality but have brain waves of abnormal frequency.

The E.E.G. has therefore become an indispensable aid to the doctor who is asked for advice about the risk of inheritance by a person about to marry who has a personal or family history of epilepsy. The answer is that if the proposed mate has no personal or family history of the disease and has a normal E.E.G. the risk is negligible.

It is fairly safe to assume from the available data that the most important cause of idiopathic epilepsy is an inborn liability to abnormal cortical discharge. If this is granted we still want to know what are the conditions which from time to time precipitate this discharge. We know that most people may have epileptic attacks under certain conditions—for example, extreme degrees of cerebral anoxia or hypoglycaemia—but in idiopathic epilepsy no such conditions are required. The attacks as a rule occur without any apparent cause. In a very small proportion of the cases the opposite is true—the attacks occur only with a specific precipitant. These cases of so-called reflex epilepsy form an interesting group. The best-known example probably is acousticomotor epilepsy, in which a sudden noise precipitates an attack, but there are many others. Coughing, micturition, and sexual orgasm, for instance, may be specific precipitants, or sudden contact of particular parts of the body. In this last group sometimes local contact will provoke a focal attack in the part touched, and electrical records can show the arrival of the sensory impulse in the cortex, the local epileptic discharge, and the resulting muscular contraction occurring in a time sequence corresponding with the known rate of conduction of the nervous impulse.

The artificial production of attacks in epileptics is often possible by alkalosis induced by over-breathing, or by

extreme hydration brought about by a high fluid intake and injections of "pitressin," but there is no evidence to suggest that these causes are commonly of any clinical importance.

Age is certainly important. Infants are peculiarly susceptible to fits, but it is certain that in many cases infantile convulsions are evident of an abnormal liability in this direction. The proportion of infants with convulsions who later develop epilepsy is higher than of those who have not had them: a history of infantile convulsions is commoner in epileptics than in non-epileptics; and a history of infantile convulsions is more often obtained among the relatives of epileptics than of non-epileptics: all these differences are significant and the point has an important bearing upon treatment. An infantile convulsion, especially if without adequate precipitating cause, should be the occasion for inquiry into the family history, and if this is positive regular and protracted sedative treatment should be given.

A striking feature in many cases of epilepsy is the time at which the attacks occur. They may occur only during sleep or between certain hours of the day, often during the first hour or two. A monthly rhythm is frequently noticed in women on account of the relation to menstruation, but is also to be observed in males. Periodicity of the attacks, however, may be seen at much longer intervals of months or years (Griffiths and Tylor Fox, 1938). Such periodicity must always be looked for and taken into account in treatment. Without this it is impossible to estimate the efficacy of drugs.

Diagnosis of Epileptic Attacks

I shall confine myself here to the distinction between epilepsy and syncope, which I think may be a very difficult problem. Syncope as a rule has an obvious cause—for example, long standing, especially on an empty stomach, sudden emotion, or intense apprehension—and it is fairly safe to say that a sudden attack of falling with loss of consciousness without such a cause is not syncopal. It is not so safe to assume that an attack of this kind provoked by emotion is not epileptic. In some persons with reflex epilepsy emotion may be the precipitant, and it may be a particular kind of emotion. In a syncopal attack there is usually some warning—dimness of vision, weakness, uncertainty of balance. But this is not an absolute rule. A person may fall in a faint with little or no warning; and, conversely, a minor epileptic attack may be preceded by similar sensations. If the epileptic attack is closely observed twitching of face or hands is commonly seen—symptoms which at first might be considered diagnostic of epilepsy. But this is not true. If the cerebral anoxia resulting from syncope is severe enough an epileptic discharge with twitching is the result. The following case illustrates the difficulties of differential diagnosis.

In 1937 I saw a boy aged 13 with the history that in the previous six years he had had five attacks of unconsciousness. The first occurred when he had a tooth out with local anaesthesia, the second on getting up after influenza, when at lunch he suddenly slid unconscious under the table. The third happened when he was standing in a row of boys at school waiting to be examined by a doctor, the fourth also at school while waiting to be inoculated, the fifth when because of the recent attack he was taken to see his family doctor, who gave the following account. "While I was listening to his heart and chest I looked up at his face and noticed he was very white. He then fell. At first he was pale; his lips then became cyanosed, and at the same time I noticed twitching of fingers and hands. This lasted only a few moments; his colour then immediately returned and he got up and said he was all right." The only relevant fact in the family history was that a maternal uncle had from childhood

always fainted easily when frightened—if, for example, he saw an accident. The patient himself had had a serious head injury in a car accident at the age of 3, being severely confused for a fortnight. I decided in favour of syncope, and heard no more of him till he was 21 and an officer in the Guards.

The story then was that he had had no further attacks till he was at Sandhurst, when while listening to a lecture on first aid he had suddenly lost his senses and fallen from his seat on to the floor. A year later he had a similar attack while being medically examined. The medical officer was convinced from his observation that this was an epileptic attack and sent him to a neurologist with this diagnosis. While the neurologist was listening to his heart it suddenly slowed to a rate of about 20 a minute and then stopped for an estimated period of 45 seconds, during which the report stated "he had a fit, rolling eyes, rigidity, and a few convulsive movements." The neurological diagnosis was Stokes-Adams disease, and he was recommended for invaliding. The E.C.G. and E.E.G. were normal.

Dr. Denis Williams then staged an attack. Preparations were made for simultaneous recording of E.E.G. and E.C.G.; a rather large needle was produced and the patient was told that it would be stuck into his arm and would be painful. He at once became pale, with marked slowing of the heart beat; his lips became cyanosed and he was momentarily confused. There was no abnormality in the E.E.G. My conclusion was that these attacks were neither epileptic nor symptomatic of heart block, but severe syncopal attacks with a specific emotional precipitant. I gave considerable weight to the family history. However, I advised that he should see Dr. Parkinson, who was confident that they were not Stokes-Adams attacks but gave his opinion against the diagnosis of syncope. His arguments, which he has kindly allowed me to quote, are of great interest. "I know that in extreme instances of fainting convulsions can occur, but rarely. Ordinary faints seldom persist from age 8 to 21 years; the warning is longer, and lesser degrees are common in the same individual. You will gather that in my opinion they might be severe faints, but I incline personally to regard them as of an epileptic nature."

It is my own opinion that we have here an example of what might justly be called reflex syncope. I have met with several comparable examples, and all have had these features in common: (1) a specific liability to faint at the sight of blood, or an accident, or the account or thought of injury or illness; (2) a tendency for this liability to begin in youth and persist into adult life; (3) the occurrence in some attacks of epileptiform convulsions; and (4) a family history of similar liability.

Anticonvulsant Drugs and E.E.G.

The E.E.G. has been of a certain value in providing scientific proof of the efficiency of anticonvulsant drugs. Many patients who are subject to repeated attacks show between these attacks very frequent brief discharges on the E.E.G. which are characteristic of an epileptiform explosion. These are known as subclinical attacks, and it can be shown that their frequency can be diminished or even abolished by sufficient doses of phenobarbitone. Another interesting demonstration is the effect of concentrated mental activity in inhibiting these subclinical attacks, confirming the impression of all experienced clinicians that epilepsy is most apt to occur when the mind is empty and least when it is active and busy. Unfortunately the impression of many lay persons is the opposite, and the doctor is often expected to prescribe prolonged absence from school at the onset of epilepsy—advice which is sometimes given with harm to the patient.

The most important contribution of the E.E.G. to treatment, however, is in connexion with the new drug "tridione." The normal E.E.G. consists of a fairly regular 10-a-second rhythm of low-voltage discharges. Of all patients with clinical epilepsy 25% show rhythms within

normal limits; 50% show abnormality of an irregular kind—usually medium-voltage slow-wave activity; 25% show runs of slow- or fast-wave activity which is characteristic of epilepsy. In this group—the epileptics with epileptic E.E.G.s—there are different kinds of abnormality in the E.E.G. which corresponds roughly to different varieties of attack.

In what is clinically called grand mal—loss of senses with generalized convulsions—the E.E.G. shows bursts of high-voltage fast waves or spikes. Attacks of similar pattern but lesser degree, which the clinician might well call petit mal—brief loss of senses with widespread but slight and momentary tonic and clonic spasm—are associated with the same type of E.E.G.—i.e., the grand mal type of E.E.G. A new name is needed for these attacks, and, based upon the E.E.G. findings, the somewhat ambiguous title "minor grand mal" has been used to describe them. The term has some practical value because attacks with this type of E.E.G., whether grand mal or minor grand mal, respond on the whole better to phenobarbitone, with or without "epanutin," than to other drugs.

Another type of E.E.G. is that in which there is a series of medium-voltage flat-topped slow waves. This is found to be associated with attacks in which confusion and disordered behaviour are the outstanding features, with little or no muscular twitching. These so-called psychomotor attacks respond best to epanutin.

The third type of E.E.G. is characterized by alternating slow and fast waves of rather high voltage—the so-called wave-and-spike complex. This again is subdivided into a round wave followed by a sharp spike—the so-called dome-and-dart E.E.G.—which is characteristic of true petit mal, and a more rapid alternation of flat-topped wave and spike known by electro-encephalographers as the petit-mal variant. Of practical importance here is that if attacks are associated with the dome-and-dart E.E.G. they are likely to be controlled by tridione and are equally unlikely to be influenced by any other anti-epileptic drug. The dome-and-dart E.E.G. is invariably found in what at one time was called pyknolepsy. The main characteristics of this variety of petit mal are an onset in childhood; the daily occurrence of frequent, brief, stereotyped attacks of loss of consciousness seldom accompanied by muscular twitching, except in the face; preservation of the intellectual functions; and a relatively good prognosis. It was recognized before the advent of tridione that the attacks not infrequently ceased at puberty never to return. Other clinical varieties of what the electro-encephalographer calls a true petit mal because of the association with a dome-and-dart complex are brief myoclonic attacks and attacks in which there is sudden falling with loss of consciousness but without twitching. Lennox (1947a) has reported the effects of treatment with tridione in 166 patients with true petit mal, having the characteristic E.E.G.: 31% became entirely free from attacks, 32% had less than one-fourth the previous number; 20% improved to a lesser extent, and 17% are unimproved. By contrast, tridione given to patients with uncomplicated grand mal brought about improvement in only 20%, and actually made 50% of the patients worse. Patients with psychomotor attacks are not so often the worse for tridione, but they do not do so well as with epanutin. I have mentioned that the clinical pattern of true petit mal varies, including very brief loss of consciousness with twitching, if any, confined to the face; myoclonic twitching without loss of consciousness; and attacks of falling with momentary disturbance of consciousness. All three of these clinical variants are equally amenable to tridione. The shorter the history and the younger the patient the better the result of treatment.

Tridione is dispensed in 0.3-g. capsules. The initial daily dosage for infants is one capsule a day, for those aged 2–4 years two capsules a day, for those 5–16 years three capsules a day, and for adults three or four capsules a day. If after one month the attacks are not controlled the dose is increased by one capsule at monthly intervals until relief is secured, toxic symptoms occur, or the dose is doubled. The maximum improvement is usually in the first month, but Lennox reports the case of a child of 8 having 50 to 70 attacks a day in whom at the end of six months the dose was raised to seven capsules daily with final disappearance of the attacks.

The commonest toxic symptom is photophobia, which occurs in about half the cases. It is very seldom severe enough to require cessation of treatment, though sometimes the patient may have to wear dark glasses in sunlight. Rashes have been observed in 14% of cases treated, usually acneiform and confined to the face, sometimes generalized and like that of measles. This calls for cessation of treatment, which is begun again when the rash has subsided, starting with one capsule every other day and gradually increasing the dose to the required level. Most patients acquire tolerance. Much more important, but apparently rare, is the danger of agranulocytosis. Lennox reports three deaths from this cause, and it follows that a fortnightly white blood count is essential for all patients taking tridione. Lennox recommends that a leucocyte count below 4,000 together with polymorphonuclears less than 40% should be an indication for at least temporary discontinuance.

Social Problem of Epilepsy

The epileptic in society has not hitherto been given much sympathy or help. He is obviously disqualified from many professions and trades, but most epileptics are intelligent and normal persons apart from their attacks, and quite able to earn their living. In the U.S.A. it has been estimated that 1 in 200 of the population are epileptic, and there is no reason to suppose that the incidence in this country is less—which means that there are something like a quarter of a million persons suffering from this malady in Great Britain. Supposing half of these to be educable and employable the problem deserves special consideration if only on account of its size. A good deal has been done in America to bring the public to a sense of their responsibility in this matter. There is, however, an instinctive repugnance among healthy people not so much for the disease as for the sight of an attack, and it is for this reason that I believe schoolmasters and parents on the one hand and employers on the other may successfully resist the humanitarian appeal for the epileptic to be admitted more freely to the normal social group. More epileptic schools and self-supporting colonies therefore seem to be the answer.

One problem which most of us must have encountered is that of the epileptic who wants to drive a car, and often does so whatever medical advice may have been given. The applicant for a licence has to answer the question "Do you suffer from epilepsy?" and is told that if in doubt regarding the answer he should get professional advice. The responsibility for declaring a liability to fits clearly lies with the applicant, but it is presumably the responsibility of the applicant's medical attendant to inform him that he has such a liability. Many patients have consulted me who have had one or more epileptic attacks but as the result of treatment have been free from attacks for so long that they claim, not unreasonably, that they should no longer be considered liable to fits as applicants for a driving licence. I have sought from the Ministry of Health and from the Home Office a definition of epileptic liability for purposes

of the Road Traffic Act with no very satisfactory result. My inquiries were related to two separate cases—one that of a man whose fits had been traumatic and who after surgical removal of a bone fragment which had penetrated the dura had remained free for two years. The other patient, a young farmer, had had two attacks of the idiopathic kind. I advised regular dosage with phenobarbitone and told him on no account to stop this without seeing me again. He returned six years later, having continued regularly with his treatment, with no further attacks. Owing to the death of his father he was now in a position in which inability to drive a car was a very serious handicap.

I am informed that the Ministry of Transport in drawing up the regulations for driving licences were advised by a committee of experts appointed by the B.M.A. and were advised that "once an epileptic always an epileptic," and that any person who has at any time suffered from fits should be refused a driving licence. No legal ruling, however, was made, and the answer to my specific question in 1946 was that there is no provision in the law which defines the time which must elapse from the occurrence of a seizure before an applicant for a driving licence may be certified as not being liable to fits. "The legal department takes the view that subsequent liability under the circumstances is entirely a medical question." This to my mind is an unsatisfactory state of affairs, placing as it does an unfair onus upon the practitioner. For his own protection he would probably be wise to inform his patients that the occurrence of a single epileptic attack is a permanent disqualification for a driving licence. This also may be the ideal advice for the protection of society, but is it practically fair? As Lennox (1947b) points out, driving accidents must be at least a hundred times more often due to chronic alcoholism than to epilepsy, but the applicant is not required to state that he has never been under treatment for alcoholism. Nor are patients with known coronary disease debarred. I have known two who have had coronary occlusion when driving, in each case resulting in a serious accident. I think it is time that legal machinery should be set up which would enable the epileptic who has been free from attacks for a specified period to state his case before an impartial tribunal. This would not only be fairer to the epileptic but would also be in the public interest. I have no doubt that at present many epileptics are driving to the danger of the public who might be deterred from doing so by a law providing a specified period of freedom from attacks.

REFERENCES

- Griffiths, G. M., and Fox, J. Tylor (1938). *Lancet*, 2, 409.
Hunt, J. R. (1922). *J. nerv. ment. Dis.*, 56, 351.
Lennox, W. G. (1947a). *J. Amer. med. Ass.*, 134, 138.
— (1947b). *Canad. med. Ass.*, 56, 638.
— Gibbs, E. L., and Gibbs, F. A. (1940). *Arch. Neurol. Psychiat.*, Chicago, 44, 1155.
Natrass, F. J. (1943). *British Medical Journal*, 2, 481.
Penfield, W. (1946). *Ibid.*, 2, 639.

The annual report of the Institute of Child Health of Birmingham University has recently been published. The Institute attaches great importance to postgraduate training, dividing it broadly into three main types: (1) for those doctors who intend to specialize in paediatrics; (2) for general practitioners who want a refresher course; and (3) for medical men who want to specialize in administration. The Institute has received large numbers of applications from postgraduate students in Britain and abroad, and during 1947 received students from India, Palestine, Italy, Poland, Malta, and South Africa. Some of the main lines of research carried out at the Institute are outlined in the report as follows: (1) The nutrition of premature infants. This work is sponsored in part by the Medical Research Council and is being carried out at the Premature Baby Unit at Sorrento. (2) Protein digests in premature, wasted, and marasmic infants. (3) Infections in infancy—nature, reactions, and treatment. (4) Protein and fat metabolism in coeliac disease. (5) Aplastic anaemia, kernicterus, and other haematological problems. (6) Causes of stillbirth and neonatal death.