cholesterol estimated at 400 mg. per 100 ml., was demonstrated by Dr. H. J. Anderson in 1943 (*Proc. R. Soc. Med.*, 36, 179). Effort angina was first complained of at the age of 52. A sister was known to have had similar "nodules" and angina of effort. There may also be a connexion between arterial atheroma and gall-bladder stones, especially solitary cholesterol stones (D. M. B. Gross, *J. Path. Bact.*, 1929, 32, 503). This would serve to illustrate the whole subject of the occasional association of different types of local or systemic errors of lipid (notably cholesterol) metabolism.

A man with hypercholesterolaemic xanthomatosis told me that a brother of his had recovered from similar lumps on his elbows without medical treatment; but I understood that that brother had recently had to have one leg amputated for what seemed to be ischaemic gangrene—very likely connected with atheromatous disease.—I am, etc.,

London, W.1.

F. PARKES WEBER.

Intestinal Lesions in Hypertension

SIR,—Dr. George J. Cunningham (June 5, p. 1075) has described the morbid anatomy of intestinal lesions due to malignant hypertension. In the past two years I have had three cases showing the clinical counterpart of this condition, which is not uncommon.

Case 1 was a woman of 37 with a B.P. of 280/170. There were marked retinal changes but no albuminuria. She was struck down one day by severe epigastric pain. This made her roll about in agony and she was so ill she could not attract help from her neighbours. Her attack lasted for four hours and went as suddenly as it came without any treatment. She had a similar attack 18 months later, and the pain was abated by morphine. I saw her after her first attack. The pain was not a colic. There was no evidence of either gall-stones or coronary thrombosis, and the physician who saw her a week later agreed with my diagnosis of "visceral angina." Vaso-spasm can occur in any part of the body and be accompanied by excruciating pain. The second case was similar in character, the pain being referred to the right femoral region. In the third case the pain was in the lumbar region.

In each case there was severe pain not fitting into any usual clinical picture. Each case had a marked hypertension. Morphine relieved the symptoms, which did not recur immediately. Two cases have had recurrent attacks several months after the first. The pain is so severe as to make it a first-class medical emergency. Amyl nitrite had no dramatic effect on the pain, although it appeared to give some relief while the morphine was doing its work.—I am, etc.,

Leicester.

C. A. H. WATTS.

The Spread of Q Fever

SIR,—Your annotation (June 12, p. 1145) prompts me to add Iraq to the list of countries in which Q fever has been found.

In May, 1945, a British surgeon in that country felt out of sorts for a few days and thought he was developing a severe cold. This did not take the usual course; instead he lost appetite and had intense headache.

He was admitted to hospital with the provisional diagnosis of typhoid or typhus. Mental disturbance, familiar in typhus but which I have not seen recorded in Q fever, became prominent. He was a "difficult" patient, unwilling to take food or stay in bed and not fully aware of what he was doing. Later he became weaker, confused, and his talk became irrational. The severe headache continued. Subsequently he had no recollection of being taken to hospital or of events for the first nine days of his stay there.

In addition to his mental symptoms the condition of his circulatory system gave cause for alarm. His blood pressure fell markedly and his pulse was irregular. This was associated with missed heart beats and accompanied by oedema of the ankles. Congestion at the bases of his lungs was revealed by percussion and auscultation. No x-ray examination was done.

Fever was high during his first eight days in hospital. His maximum temperature on the day of admission was 101.5° F. (38.6° C.); this rose to 104° F. (40.2° C.) on the fourth, fifth, and sixth days in hospital. On the ninth day in hospital and twelfth day of illness it fell to near normal.

Twenty-five days after entering hospital he returned home very weak and having lost much weight. The best way to hasten his convalescence appeared to be to get him out of the heat of Iraq, so, as soon as he was fit to travel, which was two weeks after he

left hospital and six weeks after the onset of illness, he was taken by air to England. He was still weak, had difficulty in accommodating and occasional double vision, and it was not till two months later that he felt fit to journey by himself to Canada to join his family.

Complete Widal and Weil-Felix reactions carried out during his illness gave negative results, as did blood culture. It was not till he arrived in Canada that a retrospective diagnosis of Q fever was made on a sample of serum sent to Dr. Norman H. Topping, of the National Institute of Health, U.S. Public Health Service, who reported positive complement fixation to an end titre of 1:1,024 with Q fever antigen.

The source of infection is unknown. There was no contact with cattle, but seven days before the onset of his own illness the patient examined an Iraqi soldier severely ill with an undiagnosed febrile condition. While doing so he had a distinct sensation of an insect crawling on his knee (he was wearing shorts at the time).

During the six months I remained in Iraq I looked for further cases of Q fever both by testing sera from suspects against an agglutinating antigen kindly supplied by Dr. F. M. Burnet and by sending sera to Dr. Norman H. Topping. No more were found, but I cannot believe the case now reported to be unique. Obscure febrile illnesses are common in Iraq and some may well prove to be Q fever.—I am, etc.,

University of Sheffield.

C. P. BEATTIE.

Pain in Childbirth

SIR,—I read with interest the quotation from *The Soul of a White Ant* which appeared in Dr. J. A. Hadfield's letter (June 12, p. 1155). I do agree that scientific investigation of the matter of pain and anaesthetics in childbirth is called for. Eugene Marais's hypothesis assumes that the half-wild Kaffir buck did indeed suffer pain in labour. Can he disprove that the administration of anaesthetic relieved them, not of pain, but of the satisfaction of completing a natural function? I have known many human mothers who, having been anaesthetized during labour, said afterwards that they found it hard to believe that the baby was theirs, or that they had really had a baby. What chance then for a half-wild Kaffir buck?

The difficulty of assessing the intensity of pain, particularly in animals, is almost insurmountable. During labour my pulse rate increased and I perspired, but I was in no more pain than when I lift the end of a heavy sideboard at spring-cleaning time. Since reading Dr. Grantly Dick Read's *Revelation of Childbirth* (which I commend to Eugene Marais and to Dr. Hadfield) I gave birth to a daughter and later to a son, in each case without pain, though at the first confinement a kindly G.P. thought I was suffering and I was very briefly anaesthetized, but not until after the head was born. My son was born before the doctor arrived. It was a natural and painless labour such as a half-wild Kaffir buck might be expected to enjoy. As he was born—and ever since—I loved him. I submit that if mother love is brought about by a release of hormones—and it seems to me to be very likely—then it is the completion of normal, painless labour, and awareness of bringing forth a child, that are the stimuli.—I am, etc.,

York.

GWENDOLINE ROWNTREE.

SIR,—Dr. J. A. Hadfield (June 12, p. 1155) suggests that, as pain in childbirth is necessary for the development of the maternal instinct in animals, anaesthesia during labour may inhibit the development of this instinct in women. It may be true that the stimulus of pain is necessary in animals, but before we deprive women of the great benefit of painless labour it is necessary to assess the value of this "maternal instinct" which would be so dearly bought.

In the first place, are instincts necessarily good? They are essential to the maintenance of life among animals, but the whole progress of man is a struggle to tame and sublimate his instincts, and where possible to substitute reason for instinctive reaction. Even if we were to accept the necessity for reliance on such reactions, we should hardly try to emulate the animals. Consider the duration of maternal feeling in a bitch, a cat, or a vixen: after a few weeks of lactation the young ones are more and more neglected, and soon the mother is unable even to recognize her offspring. If this is the "maternal instinct" which women are supposed to lose by avoiding the pains of labour, I do not think that the human race will be very much worse off without it.