shrubs and small trees. The contamination was not confined to one genus, but *Trichophyton* species predominated. The Petri dishes, on being opened, would give an acrid smell produced by these fungi strong enough to check the breath for a time. The frequency of this infection was such that I had to seek the advice of the late Prof. Bruhns and of his colleague Dr. Alexander, joint authors of *Allgemeine Mykologie*, in Jadassohn's *Handbuch der Hautkrankheiten* (1928).

Dr. Alexander kindly told me that they had found woodcutters and similar labourers who frequently came in contact with wood and timber most liable to such infections. They had presumed that the natural habitat of pathological fungi must be wood or freshly cut timber. The place where I used to collect the insects was exactly such a habitat.

In Europe mycotic infection of the feet, particularly of the toes, is rather common. It is often believed that the infection is derived from public bathing institutions. There wooden planks are provided for bathers to stand on immediately after leaving the bath. Wood, thus constantly moist, would appear to form an artificial habitat for such germs. This explanation brings in line with the main finding communicated above.—I am, etc.,

Bombay, India.

S. MAHDIHASSAN.

Milk-borne Tuberculosis

SIR,—We read with interest the annotation (Feb. 14, p. 303) of two articles which appeared in the Swiss journal *Gynaecologia* (1947, 123, 265). We draw attention to the last sentence, which reads, "Many of these infections are undoubtedly milk-borne and serve to emphasize yet again the importance of providing a safe milk supply."

There is only one way of knowing whether an infection is of bovine origin or not, and that is to isolate the organism and determine its type. Our protest is against the subtle insinuation that "many are undoubtedly milk-borne" without any proof to support such a statement. Our own experience is that bovine infections are not so common now as some people suggest, and recent surveys also support our view. The dangers of contact infections with tubercle bacilli of human origin are greater than infections from unsafe milk, even in those forms of tuberculosis so persistently suggested as being milk-borne.

We welcome sensible efforts to provide a safe milk supply and believe that this can be achieved without ignoring facts or having recourse to misrepresentation or to exaggeration.—We are, etc.,

PETER W. EDWARDS.

LESLIE J. CUTBILL.

Market Drayton, Salop.

Relief of Pain in Midwifery

SIR,—The subject of analgesia in labour has recently been discussed in your columns. In this connexion it must be remembered that there are two components in the pain of labour. (1) First stage—pain due to uterine contractions. These should, like gastric contractions, be painless, and with proper training in relaxation, as advocated by Dr. Grantley Dick Read, can become so. Nevertheless, just as gastric contractions may be rendered painful by underlying pathology such as an ulcer, so may uterine contractions also in a few cases cause pain and require pethidine. (2) Second stage—pain due to distension of the perineum. Now if the skin of any other part of the body is stretched it is certainly painful; and how in the world is Dr. Read to persuade us that the perineum is less sensitive than any other portion of the anatomy? No. These pains are very real and severe and in most cases call for anaesthesia.

We must remember that 80% of women in this country are delivered by the midwife alone, so the most important thing is to find an anaesthetic which she can safely administer without medical aid. Only the Minnitt fulfils this. Striking success is obtained with this machine *provided the patient is properly trained in its use beforehand*. May I issue a plea to all doctors and midwives who have lost confidence in the Minnitt to try it again, this time giving their patients a good half-hour's antenatal introduction to it and instruction in its use? This could quite easily be done in groups at clinics. I am sure those who take the trouble to do this will find themselves amply rewarded by results.—I am, etc.,

Leek, Staffs.

ANNE M. TOMS.

The Problem of Chilblains

SIR,—Dr. R. John Gourlay's comprehensive treatise on chilblains (Feb. 21, p. 336) commends itself to all interested in the study of that and other vascular disorders.

I would draw his attention to an inaccuracy in his description of cyanosis. He states that carboxy-haemoglobin is responsible for the cyanosis and that cold interferes with the formation or causes the breakdown of carboxy-haemoglobin and thus prevents the elimination of the waste products of cellular metabolism. Carboxy-haemoglobin is a relatively stable compound of carbon monoxide and haemoglobin, and as such it does not exist in normal blood.

The cyanosis is due to the presence of reduced haemoglobin, and depends fundamentally upon the absolute amount of reduced haemoglobin in the capillary blood and not upon the relative proportions of reduced haemoglobin and oxyhaemoglobin. The capillary blood must contain approximately 5 g. of reduced haemoglobin per 100 ml. blood before cyanosis will appear (Lundsgaard, C., and Van Slyke, D., *Cyanosis*, 1923, Baltimore). Carbon dioxide can form a loose carbamino compound with haemoglobin which although small in amount is important in CO₂ carriage, but which as far as I can ascertain does not contribute to the cyanotic colour of blood.—I am, etc.,

Dublin.

WILLIAM P. MCKEEVER.

Dicoumarol

SIR,—In view of the recent correspondence on the control of dicoumarol therapy by prothrombin-time estimation I should like to make three points which have not hitherto been mentioned and which have been noticed in a series of 50 cases, using Fullerton's method.

1. The variability of individuals. In 40 normals, prothrombin times of 10–22 seconds were obtained (av. 16.2), but the individual himself remains fairly constant and any departure from this time due to operation or dicoumarol therapy is related to his own base line, and to it he returns in course of time. Consequently when testing reagents or when comparing abnormals with normals at least 5 normals must be taken to avoid gross errors (Aggeler *et al.*, *Blood*, 1946, 1, 220), and for each new patient a base line should always be obtained if possible and all subsequent readings related to this.

2. In the standardization of technique, I found that I could not get reproducible results if the tubes containing plasma and "stypven" were in the 37° C. water-bath for less than 3–6 minutes—after a half-minute the shorter the time at 37° C. the quicker the reaction. At 1/2 or 1 minute one could get times of 11–12 seconds or less, but they were not constant. The prothrombin times increase with the length of time in the 37° C. water-bath till 3–6 minutes, and thereafter remain relatively constant till 12 minutes, when they again increase slightly.

3. The Wintrobe anticoagulant was found to be quite suitable for the collection of specimens, but it was most important to observe that the volume of blood added was in accordance with the amount of oxalates which had been dried into the collecting tube. If smaller volumes of blood were added to the Wintrobe anticoagulant falsely long prothrombin times were obtained.

Even if all these points are observed patients will occasionally bleed slightly with prothrombin times only 1½ times their normal, but in these cases the clotting time is prolonged in spite of a Quick's Index in the region of 40%.

It is hoped to publish a more detailed report some time in the near future.—I am, etc.,

Sector IV Laboratory,
Watford, Herts.

G. A. JAMES.

Juvenile Delinquency

SIR,—In your leading article entitled "Juvenile Delinquency" (Jan. 31, p. 203) you emphasize the urgent necessity for further investigations into "the efficacy of different methods of treatment as judged by the after-histories of cases of different types." There are, I suggest, several more urgent measures, solidly founded in psychology and calculated to produce practical results far more quickly.

Virtually all juvenile delinquents are made, not born, and the principal factors in their making are the parents' attitude and the child's economic and social environment. Unquestionably the first factor is the more important. Good parents can do much to