Comments

It would appear from these results that a year is a suitable interval between surveys, thus supporting Reid's findings in America. However, the public must be educated in the advantage to be obtained from a periodic x-ray examination of the chest, as only by an adequate response can the scheme become fully effective. The fact must be stressed that a negative fluorograph is no guarantee that an abnormality will not develop at some future date.

The considerable fall in the number of cases of pulmonary tuberculosis found seems to be due to at least two factors; (1) the discovery of the early observation lesion with the chances of its subsequent control; and (2) the diminution of the infector pool. The early removal of the infective case must greatly lessen the incidence of disease among its contacts of all ages is generally admitted, and has been demonstrated in contact work at the Ealing Chest Clinic (Dick and Thompson, 1946). This preventive aspect of mass radiography is probably not fully appreciated.

Also, by finding cases *before* they have become sources of infection to their fellows and not, as is more usual if diagnosis has to wait on symptoms, after they have broadcast bacilli for many months or even years, the infector pool is prevented from being restocked. In this light the importance of mass radiography as a preventive measure is truly seen.

REFERENCES

- Dick, W. Pointon (1945). *British Medical Journal*, **2**, 568.
— and Thompson, B. C. (1946). *Lancet*, **2**, 791.
Medical Research Council (1945). *Spec. Rep. Ser.*, No. 251.
National Tuberculosis Association of America (1940). *Diagnostic Standards*.
Reid, Ada C. (1940). *J. industr. Hyg.*, **22**, 303, 408.
— (1941). *Ibid.*, **23**, 35.

Medical Memoranda

Transmission of Syphilis

In a previous communication it was shown that the transmission of syphilis was possible during the incubation stage in the male (Ross, 1939). It was suggested that the semen was infectious and that seminal transmission of the disease had occurred in the case quoted. A case has recently occurred in which the male was suffering from sero-negative primary syphilis and according to some standards was adequately treated. It is again suggested that the method of transmission was by the seminal fluid.

CASE RECORD

On April 1, 1947, a married woman aged 31 reported at a clinic complaining of discomfort in the genital area. She stated that her husband was home on leave in July, 1946, and led a normal married life. In November he was demobilized, but did not resume marital relations. She was of the opinion that her husband realized that she was fatigued by the attention she had to give to a child of the marriage 18 months old who was in indifferent health. However, during the New Year celebrations, while he was partially intoxicated, intercourse occurred on two occasions, and thereafter was discontinued. At the end of January she experienced some discomfort in the genitals, but as this passed off in a few days she did not seek advice. Towards the end of March the discomfort returned and she reported at the clinic. No extramarital intercourse was admitted.

On examination numerous small superficial circular ulcers were found on the labia minora and perineum. Inguinal lymphadenopathy was marked but no other signs were present. Dark-ground examination showed *Treponema pallidum* in serum from the ulcerated areas and the blood Wassermann reaction was strongly positive. A diagnosis of early syphilis was therefore proved, but it was difficult to decide whether it was a case of early secondary syphilis or of

sero-positive primary syphilis with multiple lesions. The date of infection must have been about Jan. 2.

The husband was found to be attending a male clinic, and his history was of illicit intercourse in August, 1946, followed three weeks later by a primary lesion of the penis. *T. pallidum* was found in the lesion, but the blood Wassermann reaction and Kahn tests were negative. He was treated in a military hospital for ten days and received 30,000 units of penicillin in aqueous solution every three hours to a total of 2.4 mega units. Thereafter weekly injections of neoarsphenamine and bismuth were continued till Oct. 31, when the total dosage was 2.85 g. of neoarsphenamine and 1.2 g. of bismuth.

He came under the care of the civilian clinic on Dec. 18 and presented no signs of syphilis. The blood Wassermann and Kahn tests were negative. On that day and on Jan. 2 and 8, 1947, he was given 0.6 g. of neoarsphenamine and 0.12 g. of bismuth. He defaulted from treatment, and except for one visit on Feb. 21 was not seen till April 8, when he was clinically and serologically negative.

Thus it would appear that the male patient, infected in August, whose treatment was begun during the sero-negative primary stage in September and continued until Oct. 31 and whose blood Wassermann and Kahn reactions were negative on Dec. 18, was still capable of transmitting the disease during sexual intercourse on Jan. 1 or 2. Further, in spite of the lack of treatment between Jan. 8 and April 8, he was clinically and serologically negative on the latter date.

Therefore the question arises, When is a male who has suffered from syphilis non-infectious? From occasional cases seen in the past twenty years one is confirmed in the belief that the time factor is of more value than the serological test for syphilis. In support of this belief another conjugal case is quoted.

In March, 1936, a woman aged 32, married some eight months, reported with periostitis of both tibiae of the type associated with early syphilis. She gave no history of extramarital intercourse. Her blood Wassermann was strongly positive, and it was thought advisable to ask for the attendance of the husband. He proved to be a defaulter from treatment, but all tests of blood and cerebrospinal fluid were normal.

He was first treated in London in February, 1932, for secondary syphilis. In July of that year his blood Wassermann was negative and a second course of treatment was begun, but he defaulted and received only 2.25 g. of neoarsphenamine and 1.2 g. of bismuth. In August, 1934, he returned with acute gonorrhoea, and, since there were no signs of syphilis and his blood Wassermann was negative, no treatment for syphilis was given.

If the facts and implication of this case are accepted then a three-years history of negativity is not sufficient to exclude the possibility of the infectivity of the semen in a syphilitic. On the other hand, it is a matter of everyday experience that a male whose infection is of more than five years' duration, whether his serological tests are negative or positive, is extremely unlikely to infect his wife.

The advice, therefore, that should be given to an adequately treated male is that he should not attempt to procreate children until five years after the date of his infection.

Since this paper was written Leeming (1947), quoting a personal communication from D. A. Erskine, gives reasons for the belief that men treated for gonorrhoea by adequate doses of penicillin may, in a prolonged sero-negative syphilis d'émblée, infect their wives.

My thanks are due to Dr. R. B. Nelson, whose notes on one of the cases have been quoted.

July, 1947.

A. O. F. ROSS.

REFERENCE

- Ross, A. O. F. (1939). *British Medical Journal*, **1**, 415.
Leeming, J. A. L. (1947). *Brit. J. Ven. Dis.*, **33**, 163.

The Minister of Health has received letters of thanks from the French Minister of Industrial Production and the Minister of Public Health for medical help given after the Lorraine mining disaster in January. The Ministry granted facilities for Sir Archibald McIndoe, the plastic surgeon, of Queen Victoria Hospital, East Grinstead, and Squadron-Leader F. T. Moore, R.A.F. Medical Service, who is working with him, to fly to the scene of the disaster. The Ministry also arranged for large supplies of plasma from the National Blood Transfusion Service to be flown across for use in treating casualties. The French Minister of Production in his letter to Mr. Bevan wrote: "This outstanding assistance, so spontaneously extended, has been appreciated in all its value, and both the Government and the nation feel it to be yet another sign of the friendship which binds our two countries."