4,600,000; Hb, 92%; haematocrit, 48. On April 27 arterial oxygen saturation was 88.5%.

Comment.—There seems little doubt that there has been considerable improvement in the cardiac condition, and the fall in blood count and haematocrit reading (both of which are now normal) and the rise in the oxygen saturation percentage are most encouraging. Unfortunately the persistence of the hemiplegia makes it impossible to assess the actual functional improvement. It is thought that the paralysis will soon improve enough to allow the patient to move about by herself.

Conclusion

These three cases are by no means precise, but the results are considered to be encouraging enough to be recorded in this preliminary communication. They serve, at any rate, to show that pulmonary valvulotomy is a feasible operation, and all three patients have survived the procedure. The result in the first may be classed as moderately successful so far; in the second the result is good; in the third it would appear to be good, but is as yet indeterminate functionally.

Two of the patients were of an age when the prognosis is ordinarily poor and when any operation carries a grave risk; both were very severely disabled. There can be no doubt that far better results are to be anticipated in younger patients, but it has not yet been possible to extend the operation to the younger age groups, although it is intended to do so.

The vascular complications in the first and third cases are disturbing and disappointing. The first case teaches us that a careful watch must be kept for signs of peripheral arterial embolism after cardiomyotomy so that immediate embolectomy can be considered. Arterial thrombosis, even to the extent of hemiplegia, occurs with the Blalock operation just as often. Presumably it is most likely to occur when the blood pressure falls to a low figure for some time, especially in the presence of polycythaemia.

Much still remains to be done in the development of this operation in the treatment of "blue children" suffering from pulmonary stenosis, but the results in these three cases are encouraging enough to justify proceeding further with it. It should be made clear that it is not suggested that this operation is meant to supplant the Blalock operation. As it stands at present it is of use in only a small proportion of cases; until a method can be found to relieve the subvalvular stenosis (if indeed this is possible), the Blalock operation must continue to take precedence. However, unless the Blalock operation is considered to have solved the problem of *morbus caeruleus* completely, it is essential to try to develop new and alternative measures.

I am indebted to Dr. Maurice Campbell, under whose care all these patients were, for most of the case histories, which are made from his own notes.

REFERENCES

- Allen, D. S., and Graham, E. A. (1922). *J. Amer. med. Ass.*, **79**, 1028.
 Blalock, A., and Taussig, H. B. (1945). *Ibid.*, **128**, 189.
 Crafoord, C., and Nylin, G. (1945). *J. thorac. Surg.*, **14**, 347.
 Cutler, E. C., and Beck, C. S. (1929). *Arch. Surg.*, **18**, 403.
 Doyen, E. (1913). *Pr. méd.*, **21**, 860.
 Gross, R. E., and Hubbard, J. P. (1939). *J. Amer. med. Ass.*, **112**, 729.
 Pribram, B. O. (1926). *Arch. klin. chir.*, **142**, 458.
 Rehn, L. (1897). *Ibid.*, **55**, 315.
 Souttar, H. S. (1925). *British Medical Journal*, **2**, 603.
 Tuffier, T. (1913). *Trans. Internat. Cong., Med. Sect.*, **7**, Pt. 2, p. 249. London.

The British Red Cross Society has published a neatly illustrated pamphlet (price 1s., by post 1s. 1½d.) by Air Marshal Sir Harold Whittingham, entitled *ABC of First Aid Treatment*. It is suitable as a short manual of reference or for refreshing the first-aiders' memory.

B.C.G. VACCINATION: IS IT OF ANY VALUE IN CONTROL OF TUBERCULOSIS?

BY

ARVID WALLGREN, M.D.

*Professor of Paediatrics, Royal Caroline Medical Institute,
Stockholm, Sweden*

Several eminent German physicians who are about to introduce B.C.G. vaccination in Germany, thanks to the Scandinavian Red Cross, have called my attention to the article on this subject contributed to the *British Medical Journal* by Dr. G. S. Wilson (1947). From various quarters, including England, I have been requested to answer his criticism of B.C.G. vaccination, which has given rise to a certain hesitation and concern, since Dr. Wilson, as a member of the Medical Research Council, is a critic whose opinion carries considerable weight. As I introduced the most widely practised and, by Wilson, criticized method of B.C.G. vaccination and am largely responsible for its present widespread use in Sweden, and as, furthermore, I am prepared to help in introducing it to combat tuberculosis in the devastated countries of Europe, I feel prompted to test the value of Wilson's criticism.

I have no reason to disbelieve Wilson when he says that he is not prejudiced against B.C.G. vaccination; my faith in his objectivity is, however, put to a severe test by the manner in which he argues against B.C.G. In several points the criticism is undoubtedly justified, although correct mostly on unessentials; in many respects it is erroneous and unjust, not giving the impression of being sufficiently unbiased; and in all respects it overshoots the mark.

Investigations on Children

Wilson's quotation of my investigation (Wallgren, 1934) is entirely correct; I pointed out in that paper that it was not feasible to establish how large a part B.C.G. immunity had played in the favourable results reported. Other factors were contributory. Although many children were isolated until 3 months of age before being subject to exposure, after this period they nevertheless were exposed to virulent infection without falling ill from primary tuberculosis, save in a single case. This is a result which surely the whole battery of prophylactic measures known to us cannot parallel when dealing with children of this age that have not been vaccinated before exposure to infection. Further experience has shown me that cases of clinical tuberculosis occurring after an adequately performed B.C.G. vaccination are extremely uncommon and remarkably benign. While primary pulmonary tuberculosis in a child unvaccinated with B.C.G. does not show any tendency towards radiographically demonstrable regression during the first six months, in a child who has been vaccinated with B.C.G. and who was still tuberculin-positive at the time of infection radiographs reveal that within this period the condition has virtually resolved. It is also remarkable that I have not seen a single adequately vaccinated child die from tuberculosis after having been infected, nor have I observed a single case of tuberculous meningitis among vaccinated children. Is, then, the scientifically calculated and statistically reported degree of immunity of such great import if the effect of the vaccination is satisfactory?

Rosenthal and his co-workers (1945) have used a different method of vaccination (multiple punctures) from that with which I have had experience (intradermal). Wilson is admittedly right in stating that the results of their experi-

ments are impressive—three times as many deaths among the non-vaccinated as in the vaccinated contact group. It is, however, difficult to follow Wilson in his refusal to accept the evidence of the Chicago investigation until he has received information with regard to certain minor particulars not mentioned in the report—which particulars, however, should not have any appreciable influence on the incidence of tuberculosis in both groups. Wilson contends that the similar investigation carried out by Levine and Sackett (1946) in New York does not support the results of the Chicago experiment. This is incorrect. Levine and Sackett, remarkably enough, obtained results nearly identical with those of Rosenthal and his co-workers, when the former, in conformity with the latter, took into consideration the fact that the immunity must have arisen after vaccination and before the children were exposed to infection. The results were: Levine and Sackett—non-vaccinated 96, with 3 deaths from tuberculosis; vaccinated 90, with no deaths from tuberculosis*; Rosenthal *et al.*, non-vaccinated 63, with 3 deaths from tuberculosis; vaccinated 98, with no deaths from tuberculosis.

Vaccination of Nurses

Wilson refuses to accept the favourable result from Ferguson's (1946) well-known experiment on the sanatorium personnel in Saskatchewan because the non-vaccinated control cases belonged to a period immediately prior to that in which the vaccinations were carried out. It is evident that the conditions relevant to infection and the ability of the personnel to protect themselves *may* be different in the vaccination and the pre-vaccination periods, and therefore it cannot be considered proved that the decrease of morbidity and mortality exclusively is due to the immunity conferred by B.C.G. But why should such a change in conditions arise simultaneously with the introduction of B.C.G. vaccination? One should surely ensure that all employees, before as well as after the period of vaccination, were given instructions on the danger of infection and also on how they could best protect themselves against it. Nor can the difference in the infectivity of the patients become so great between these periods as to explain the six times higher morbidity in the non-vaccinated personnel of the pre-vaccination period. Nordwall's (1944) experience in a training school for nurses in Stockholm was in agreement with that of Ferguson. Before the introduction of B.C.G. vaccination 40% of the tuberculin-negative student nurses developed tuberculosis during service and had to break off their studies for a more or less prolonged period; one of them died. Among those vaccinated there occurred only a single mild case of primary tuberculosis, and this pupil was able to resume her service after a couple of months.

The difference between the morbidity from tuberculosis in the period before and that after the introduction of B.C.G. vaccine in Ferguson's and in Nordwall's series is so great that it cannot be entirely explained even by a certain difference in the intensity of exposure in the different groups. In the Stockholm investigation, however, such a difference was not demonstrable, nor were the student nurses protected in any more effective way during the B.C.G. period. Evidence of the efficacy of the vaccine might be obtained if during a supplementary period one refrained from vaccinating the tuberculin-negative individuals of the newly appointed personnel; but I am convinced that neither Ferguson nor Nordwall would care to take the responsibility for such a test, since experiments to date definitely support the efficacy of the protection conferred by vaccination.

Investigation Among North American Indians

With reference to the well-known test of Aronson and Palmer (1946) among the North American Indians, Wilson seems to submit to evidence and accept the results as conclusive. He admits that the difference between the B.C.G. group and the control group is striking: the tuberculosis morbidity in the former group is seven times lower. Wilson detracts from the value of this test by denying the possibility of applying the results obtained in Indians to a civilized population. Perhaps the statistical results would not have been equally striking if the experimental subjects had belonged to another group of the population with inherited natural resistance, but this we do not know with certainty. To some extent he undoubtedly has a point.

In order to substantiate his opinion Wilson compares the incidence of tuberculosis among the Indians and among the total population of the United States. This comparison, however, is very lame. He points out that the ratio of non-pulmonary to pulmonary tuberculosis in Aronson and Palmer's non-vaccinated tuberculin-negative control group was 2.5 to 1, while among the 1-25 year group of the population of the United States it is 1 to 5. Here Wilson commits at least two errors. He compares the ratio of disease in one instance with the ratio of recorded deaths in the other. This would imply that pulmonary and non-pulmonary tuberculosis had the same prognosis with regard to mortality, and this is not at all the case. Furthermore, he compares two incommensurate groups: on one side the presence of tuberculous disease *during a period of six years of observation* in individuals aged 1-20 years, *all of whom were tuberculin-negative at the start* and who were *under annual control*; on the other side, the incidence of tuberculosis mortality during one *calendar year* among a group of individuals aged 1-25 years, *probably one-half of whom were already tuberculin-positive* at the beginning of the year of observation and would thus run a lower risk of developing, for example, tuberculous meningitis and miliary tuberculosis. The latter group, too, would in all probability not have had an *annual medical examination*.

Work of Hyge

Wilson claims to have selected for review those investigations which more closely approach the conditions of controlled laboratory experiments. He has, however, excluded the investigation which perhaps best corresponds to an animal experiment—that carried out by Hyge (1947). I gave an account of this study in my address at the London conference, and it has been published in *Acta Tuberculosea Scandinavica*, 1947. Hyge reported an epidemic of tuberculosis in a school for girls aged 12 to 18 years: 105 pupils were tuberculin-positive before exposure, 106 were vaccinated with B.C.G., and 94 were tuberculin-negative. The source of infection was a temporary teacher who visited the school for only a limited time. The duration of exposure was two months in all three groups. The pupils were followed up during the first three years after exposure. Of the tuberculin-positive group only two had fallen ill with pulmonary tuberculosis during the period of observation. Of the B.C.G.-vaccinated group two contracted pulmonary tuberculosis and were in need of pneumothorax treatment. Of the tuberculin-negative group 41 fell ill with tuberculous disease; one of them died and five others needed treatment with pneumothorax. The morbidity rate in the three groups was 1.9, 1.9, and 43.6%, respectively.

Pleurisy and Primary Tuberculosis

Wilson excludes all cases of pleurisy and primary tuberculosis from the reports on the results of B.C.G. vaccination. He considers these to be practically unimportant

*Wallgren, Arvid. *Amer. J. Dis. Child.*, 1948 (in the press).

diseases. They certainly are not so in our country. The prevention of clinical primary tuberculosis or pleurisy means a very great gain: children do not have to give up school for a period of three to six months or more; and adolescents and adults avoid unemployment because of illness during an equally protracted period, besides being subject to less risk of post-primary tuberculosis. This is perhaps the most demonstrable effect of the immunity conferred by B.C.G. vaccination. Therefore if one claims to give an unbiased and comprehensive evaluation of the effect of B.C.G. vaccination it is not fair to exclude these cases.

The fact that B.C.G. vaccination is particularly effective against the immediate dangers of virulent infection in the tuberculin-negative induced Heimbeck, in his group of pupil nurses, to transfer those who originally belonged to the tuberculin-negative group to the tuberculin-positive group a year after virulent infection. This has been criticized by Wilson, who claims that all the nurses should have remained in the group to which they belonged at the beginning of the study. Heimbeck (1947) publishes a table of his results with B.C.G. vaccination of pupil nurses during their three years' training; every nurse remains in the group in which she started at the beginning of her training period. He adds that these pupil nurses lived under exactly the same conditions, and that all of them were exposed to the same risk of contamination during training. The three groups were from the same period. It seems to me that this comparison thus fulfils Wilson's claim, and still the difference between the B.C.G.-vaccinated (morbidity 24.1, mortality 2.1 per 1,000 observation years) and the unvaccinated tuberculin-negative group (morbidity 141.2, mortality 14.6 per 1,000 observation years) is striking.

Alleged Difficulties and Disadvantages

The difficulties and disadvantages that the use of B.C.G. might entail are enumerated in seven points by Wilson. Let us discuss these alleged disadvantages.

1. *B.C.G. is a live vaccine that demands great care in preparation to avoid contaminants and must be used within one week.*—Is it not also necessary to exercise great care in the preparation of smallpox vaccine and therapeutic sera? B.C.G. has been used with success after the lapse of 10 to 14 days and thus does not necessarily have to be used exclusively during the first week after preparation, although this is preferable.

2. *The virulence of B.C.G. is not fixed.*—This is correct and is also true of smallpox vaccine; but the virulence of B.C.G. vaccine from one and the same laboratory varies only slightly and never suddenly, and it is tested continuously. There is no risk of a B.C.G. vaccine becoming so virulent that it induces frank abscesses in all individuals. That this alleged disadvantage is negligible is shown by the fact that the dosage of the B.C.G. vaccine prepared in Sweden (0.05 mg.) is the same now as it was 20 years ago, and that throughout this period it has remained constant. A factor that varies, however, is the individual sensitivity to the vaccine. In a subject with a low resistance abscess formation may occur, but with a high resistance the same vaccine may not induce any reaction whatsoever. It seems to me very likely that B.C.G. resistance in the individual runs parallel with his resistance against virulent tubercle bacilli, and therefore, notwithstanding the abscess formation, it must be considered beneficial and fortunate for the person with a low resistance that he was vaccinated with B.C.G. The person with a high resistance, who perhaps did not acquire any specific immunity after inoculation, has better prospects though lacking the immunity conferred by B.C.G.

3. *Injections have to be made intracutaneously with considerable care in order to avoid serious ulceration.*—As do most workers who for one reason or other are opposed to B.C.G. vaccination, Wilson exaggerates the disadvantages of too marked a local reaction. It has been generally stressed that it is

extremely seldom we see any serious ulceration, and still more rare is the occurrence of a swelling of the regional lymph glands. Törnell (1947) found a local reaction of 10 by 10 mm. in only 2.8%. The slight ulceration that very commonly occurs is of no consequence. If the inoculation gave rise to distressing local reactions it is hardly likely that in Sweden tens of thousands of mothers annually would of their own initiative bring their children to the doctors to have them vaccinated with B.C.G. At any rate, they have from personal experience come to consider these local discomforts trivial in comparison with the benefit conferred by inoculation.

4 and 5. *Infants of tuberculous mothers must be separated at birth until B.C.G. vaccination has produced tuberculin sensitivity, or, if taken away after exposure, they must be separated for at least six weeks before inoculation.*—This cannot be a disadvantage exclusively confined to B.C.G. Does any physician of our day permit a tuberculin-negative infant or a child to remain in an infectious home in the presence of a case of open tuberculosis if there is any possibility of segregating it? In our country this would be branded as almost criminal and would undoubtedly lead to an admonition from the Medical Board. The difference between the B.C.G.-vaccinated children and non-vaccinated children is that the former may return to their infectious homes when the vaccination has taken, while the non-vaccinated must stay away while the source of infection remains. This curtailment of the period of isolation we in Sweden consider rather as one of the advantages of vaccination with B.C.G.

6. *Infants separated and kept in residential nurseries may be exposed to serious risk of cross-infection and die.*—May I ask: Is it less dangerous to keep the infant in its home and let it become infected with virulent tubercle bacilli? Are not the infants even now, before the introduction of B.C.G. vaccination in England, separated from their infectious surroundings, and, if so, where and in what way are they cared for? Is the risk for an infant separated in order to be vaccinated with B.C.G. greater than for infants separated from their dangerous homes without being vaccinated? I should think that the danger is greater in the latter, because the period of separation must certainly be much longer than the few months in the case of a vaccinated child. Out of 117 infants who were vaccinated at birth in 1928-33 and who at the follow-up examination in 1934 were at least 1 year old, only four had died during their first year of life. These infants were segregated in a nursery from birth until they were a few months old. These figures do not indicate that separation for B.C.G. vaccination, when carried out under satisfactory conditions, constitutes any great danger for the child.

7. *It is desirable to revaccinate at intervals in order to ensure the continuance of immunity.*—This is of course a drawback. In Sweden revaccination is not commonly done. The tuberculin-sensitivity has a duration of three, five, or ten years or more, and probably the immunity continues as long, perhaps even longer. At any rate, the protection has lasted during the most susceptible childhood ages, even if revaccination is not carried out at the end of the tuberculin sensitivity.

In this discussion I hope that Wilson's seven points have been judged according to their deserts.

Conclusion

Wilson shoulders a great responsibility when he endeavours to decrease the interest in B.C.G. vaccination and to discourage its introduction into his own country. Although he cautiously enough concludes by recommending experiments with B.C.G. vaccination in specially threatened groups of the population—nurses and physicians—he does not do even this whole-heartedly, seeing that he at the same time tries to depreciate the danger menacing these same groups, thus intimating the superfluity of such an experiment. It seems as though he were in fear of an unbiased trial, while we who uphold B.C.G. vaccination know that to make an adequate trial with B.C.G. is automatically to become an adherent of the method. The erroneous conclusions drawn by Wilson from cited statistical reports with reference to the danger of

tuberculosis in England have already been taken up in an editorial comment in the issue of the *British Medical Journal* in which his article appears. I therefore need not enter upon this subject.

Wilson does not stand alone in criticizing B.C.G. vaccination. I myself, 20 years ago, strongly censured the manner in which B.C.G. vaccination was then carried out, as well as the conclusions drawn (Wallgren, 1927); but later, when it was performed in a satisfactory way, I became a fervent supporter of the method. Levine and Sackett have expressed very strong opinions on the B.C.G. studies of other workers, but Wilson does not accept the work even of these authors, as the vaccinations were carried out without consideration of the fact that protection does not arise until the vaccine has taken. These two B.C.G. opponents had finally to admit that if they comply with that condition the mortality among the non-vaccinated infants is three times greater than that of the vaccinated infants; how many times greater then is the morbidity, of which one does not receive any information? Aronson and Palmer were rather sceptical with regard to B.C.G. when they undertook their investigation, intended to demonstrate whether B.C.G. had any effect. They learned by experience that B.C.G. satisfies even rather large demands as a protective measure. I am confident that Wilson also, if he really is willing to perform adequate investigations with B.C.G. under scientifically unimpeachable conditions, will by degrees come to be a protagonist of this method of vaccination.

REFERENCES

- Aronson, J. D., and Palmer, C. E. (1946). *Publ. Hlth Rep., Wash.*, **61**, 802.
 Ferguson, R. G. (1946). *Canad. J. publ. Hlth*, **37**, 435.
 Heimbeck, J. (1947). *Irish J. med. Sci.*, **65**, 689.
 Hyge, T. (1947). *Acta tuberc. scand.*, **21**, 1.
 Levine, M. I., and Sackett, M. F. (1946). *Amer. Rev. Tuberc.*, **53**, 517.
 Nordwall, U. (1944). *Acta tuberc. scand.*, **18**, 45.
 Rosenthal, S. R., Bland, M., and Leslie, E. I. (1945). *J. Pediat.*, **26**, 470.
 Törnell, E. (1947). *Acta tuberc. scand.*, **21**, 241.
 Wallgren, Arvid (1927). *Acta paediat.*, **7**, 120.
 — (1934). *J. Amer. med. Ass.*, **103**, 1341.
 Wilson, G. S. (1947). *British Medical Journal*, **2**, 855.

THE EFFICACY OF B.C.G. VACCINATION

BY

H. MALMROS, M.D.

Physician-in-chief, Central Hospital, Örebro, Sweden

Last year hundreds of thousands of young children and adolescents in war-ravaged Europe were tuberculin-tested and the non-reactors vaccinated against tuberculosis with B.C.G. This extensive campaign, which was initiated and financially sponsored by the Danish Red Cross Society, does not appear to be very well motivated if B.C.G. vaccination does not confer an increased protection against tuberculosis. According to Prof. G. S. Wilson (1947) the value of B.C.G. vaccination is rather problematic and in any case not definitely proved.

Difficulty of Procuring Control Material

The great difficulty in getting fully comparable control material is the reason why there is still debate about the efficacy of a prophylactic measure which has been in use for many years. In his famous novel *Martin Arrowsmith* Sinclair Lewis describes the difficulties met with when a scientific experiment is consistently conducted with living human beings as objects and alternate patients are taken as controls. Where mass investigations are concerned,

which require a certain amount of propaganda to induce the public to take active part, it is soon found to be impossible to treat only every other person. There will be letters to the daily press from agitated members of the community who have been refused permission to share in the new medical discovery. If it be announced that the method is still uncertain and must first be tested, the rumour arises that it is something dangerous. The result will be that only a small proportion of the public will volunteer, and hence, statistically viewed, the material will be to some extent a selected one. Nor is it possible in our modern community to inject sterile saline instead of vaccine into every alternate subject without publicly giving notice of this from the outset. Should such a measure be adopted in secret, the whole investigation would immediately become discredited when the first cases of disease appeared among those who had received only a simulated vaccination.

The difficulties enumerated here have been so great that at all centres in Scandinavia where the Calmette method has been used it has been considered necessary to abandon the statistically most trustworthy method of vaccinating alternate cases.

Primary Infection at Adult Age

How, then, has B.C.G. vaccination become so popular in Scandinavia without definite proof of its reliability? First we are convinced that pulmonary tuberculosis is nowadays very often a direct consequence of primary infection in adults (Heimbeck, 1928, 1929, 1936; Arborelius, 1930; Kristenson, 1932, 1933, 1942; Malmros and Hedvall, 1938, 1940; Törnell, 1939; Ustvedt, 1932; Frostad, 1944; S. Holm, 1947; and others). Consecutive tuberculin-testing has perhaps been used to a greater extent in Scandinavia than elsewhere, and we have been able to observe continually that previously Mantoux-negative adolescents fall ill with tuberculosis. Sometimes the primary infection produces symptoms similar to those of influenza or a "heavy cold." As a rule the radiographs then disclose a primary focus in the lung and enlarged hilus glands. In other cases it may have its onset as pleural effusion. Often, however, it initially runs an entirely symptomless course. In some of these cases, and in many of those that have had an initial fever or pleural effusion, the first radiographic lung changes that can be classed as genuine pulmonary tuberculosis appear some months later. Malmros and Hedvall showed that genuine progressive pulmonary tuberculosis often begins in these cases as small cloudy spots in the supraclavicular part of the lung or in the first intercostal space (subprimary initial foci). According to Heimbeck, Frostad, Hedvall (1946), and S. Holm, pulmonary tuberculosis can develop in some cases direct from the primary focus by progression out into the surrounding lung tissues or by liquefaction of the primary focus. The last-named has shown that no less than 34% of the cases that exhibit x-ray changes at the primary infection will later develop into genuine pulmonary tuberculosis.

Systematic tuberculin tests carried out before the onset of the disease—e.g., on probationer nurses and medical students—proved that those who were formerly tuberculin-negative were more liable to succumb. Some had relatively mild changes, others became seriously ill, and some cases were fatal. Heimbeck showed that a positive tuberculin reaction can be produced by B.C.G. vaccination, and it was a quite natural step to make an attempt to protect these tuberculin-negative adolescents by means of vaccination. Experience of the past few years seems to confirm the assumption that B.C.G. vaccination confers a certain amount of protection. Very few cases of tuberculosis have been observed among the vaccinated.