Disk Lesions and Osteopathy

SIR,—Dr. G. Macdonald's partisan letter (April 3, p. 663) really cannot be allowed to stand unchallenged. A. T. Still (the founder of osteopathy) postulated that *all* diseases arose from pressure on arteries and nerves as the result of vertebral displacements. If medical research has led to the recognition of spinal disk protrusion, and if this is now claimed as the vindication of osteopathy, it must prove the opposite of Dr. Macdonald's contention. Surely it shows that the whole of osteopathic doctrine, except in that small particular, is false.—I am, etc.,

London, W.1.

JAMES H. CYRIAX.

Intervertebral Disk Lesions

SIR,—I was interested in the letter by Dr. J. Donaldson Craig and Mr. A. W. Lipmann-Kessel (March 20, p. 570) concerning sciatica produced by prolapsed intervertebral disks. They state that the pain produced by stimulating the fifth lumbar inter-spinous ligament is clinically and subjectively indistinguishable from that occurring spontaneously in an L5-S1 disk lesion, and also that sciatic pain might be regarded as a deep segmental pain due to joint disturbance.

I would suggest at the present time that we are in danger of considering that every case of sciatica must be caused by direct pressure of a displaced disk on a nerve root, thus causing direct pressure pain. It is true that in some cases where operative exploration is undertaken for sciatica the surgeon fails to see a disk prolapse, and yet to his pleasure the patient appears cured, at any rate for some time, of the distressing sciatic pain. Possibly the mere exposure of the dural sac and the traction of it in exploring the disks is sufficient to tear down adhesions in that area. Recently I dealt with a case that illustrates well the points brought out by Dr. Donaldson Craig and Mr. Lipmann-Kessel in their letter.

A well-built young man, aged 28, was referred to me at Fleet Hospital, Hampshire, by Dr. Sharkey on account of pain in the back and sciatica of five months' duration, following a fall on to his back at football. Diagnosing a prolapsed disk causing moderate pressure, for a further four months I carried out the so-called old-fashioned treatments for sciatica—namely, diathermy applied to the back and leg, stretching of the nerve, and manipulation of the spine. Immobilization both in bed and in a plaster spinal jacket was tried, all to no effect. Finally I decided to explore the spinal cord. After removing the ligamentum flavum between the 4th and 5th laminae, and the 5th lamina, I found no prolapse of the 4th disk. On proceeding, however, to remove the ligamentum flavum from the 5th lamina to the 1st sacral, with a view to removing the sacral lamina and exploring the 5th disk, I found that there was a local adhesion of the posterior aspect of the dura to one point on the flaval ligament, just at the level of the sacral lamina. On passing a hook around the adhesion to determine its size and desirability of cutting, it was noted that traction on it produced contractions of the posterior leg muscles. Further investigation showed that one of the roots of the corda equina was also adherent in the adhesion. The adhesion was freed from the ligament and resected off the underlying root, and the dural hole carefully sutured. No prolapse of the 5th disk was found. To my joy the next day the patient reported that he had completely lost his sciatic pain, and after a normal convalescence he returned to his job as a schoolteacher, standing, and riding a bicycle to school. He has had no return of backache or sciatica.

Such a case would bear out some of the early views held that backache and sciatica may be caused by adhesions of the soft tissues to the cord and roots rather than always being due to disk pressure, which appears to be the modern trend of thought.—I am, etc.,

London, W.1.

G. O. TIPPETT.

Bone Conduction in Otosclerosis

SIR,—I would like to refer to the letter by Mr. W. H. B. Magauran (March 20, p. 569) concerning air and bone conduction in otosclerosis. It has been known for some time now that improvement of hearing by both air and bone conduction occurs in the unoperated ear in a certain number of cases—but not by any means all—following the fenestration operation, but this improvement is never permanent. In fact in over 500

fenestrations I have not one case of sustained improvement by air or bone conduction over six months in the unoperated ear, though improvement be fully maintained in the operated ear.

I feel bound to confess my amazement that anyone practising the fenestration operation to-day should rely upon inaccurate tuning-fork tests—and the one fork of the least importance at that, namely 256—for selection of suitable cases. Mr. Magauran states that "bone conduction . . . was found to be lengthened by a few seconds and to be reduced by *about* 50% on the left mastoid bone." I cannot help feeling that if adequate audiometric analysis had been carried out before the operation both ears may have been found suitable for the operation and the patient would not have had to rely on the result of the first operation to determine whether the second ear would respond to the fenestration operation also.

I could not agree more with Mr. R. Scott Stevenson's letter (March 27, p. 618) in regard to the need for a careful, sober, and dispassionate estimation of the results of the fenestration operation for otosclerosis. I would like to reiterate the plea that I made in the *Post-Graduate Medical Journal* (November, 1947) for the institution of most careful records of the pre-operative examination, details of the actual operative technique, and post-operative records at intervals of one month, three months, six months, and then yearly for ten years. In addition to these records a questionnaire should be answered yearly by the patients regarding the state of the result from their own point of view.

It is advisable that this operation should not be performed sporadically, but only by those who are willing to take time and pains necessary to develop an irreproachable operative technique, and who will undertake the complete after-treatment either by themselves or by an assistant specially trained for this purpose. Those otologists regularly practising this operation under these conditions should in my opinion compare their results from time to time, especially in relation to the technique of the operation, for it is beyond doubt that those employing the best technique obtain the best results. It is inexpedient to come to conclusions of the value of the fenestration operation based on the results of those only occasionally performing it.—I am, etc.,

London, W.1

E. R. GARNETT PASSE.

SIR,—Mr. W. H. B. Magauran (March 20, p. 569) can only have seen an abstract of the paper which I read to the Royal Academy of Medicine in Ireland on "Bone Conduction in Otosclerosis," which no doubt accounts for a misconception about one of its main points.

In a study of cases before and after the fenestration operation I showed that, after omitting those cases in which pre-operative masked bone conduction was normal and those cases in which no improvement in air conduction took place, the acuity of hearing for masked bone conduction in the operated ear was increased in 90% of cases. There was no alteration of masked bone conduction in the unoperated ear, and in fact the unoperated ears were used as controls to show that the alteration of bone conduction in the operated ears was a genuine improvement. Mr. Magauran's finding of a "considerable lengthening of bone conduction in the unoperated ear" is directly contrary to my findings, and strongly suggests that the operated ear was inadequately masked during the test. Without adequate masking of the untested ear, bone-conduction tests in otosclerosis may be quite misleading. I have never seen the hearing of the unoperated ear altered in any way, either for air or bone conduction; and indeed it is inconceivable that an operation on one ear could affect the working of the opposite one. Such a result can only be explained as an error in testing, either on the part of the patient or the examiner.

There is one further point. In assessing the suitability of a case of otosclerosis for the fenestration operation, bone-conduction tests with a 256 tuning fork are quite valueless. These tests must be performed at 512, 1024, and 2048 cycles per second. Normal bone conduction is often found at 256, when there is severe loss in the higher frequencies. Furthermore, neither tuning forks nor conversation voice tests can be used for quantitative estimation of the degree of deafness without introducing errors so large as to make the results of little value. If the results are to be expressed quantitatively audiometric measurement must be used.—I am, etc.,

Dublin.

R. R. WOODS.