BRITISH MEDICAL JOURNAL

LONDON SATURDAY JUNE 5 1948

THE LIPOTROPIC FACTORS IN EXPERIMENTAL CIRRHOSIS

BY

E. A. SELLERS,* M.D., Ph.D. C. C. LUCAS, M.A.Sc., Ph.D.

AND

C. H. BEST, C.B.E., M.D., F.R.S.

(From the Banting and Best Department of Medical Research, University of Toronto)

[WITH PHOTOGRAPHIC PLATE]

The term "lipotropic" was introduced by Best and Ridout (1935) to describe substances which decrease the rate of deposition and accelerate the rate of removal of hepatic fat. This definition has now been broadened to embrace other established effects of dietary choline or its precursors. These include the prevention of renal lesions (Griffith and Wade, 1939) and of cirrhosis in experimental animals.

The production of cirrhosis in rats and dogs by dietary means afforded a new experimental approach to the study of liver disease (György and Goldblatt, 1939; Chaikoff and Connor, 1940). In rats it was shown that cirrhosis could be produced by feeding, for an extended period, a diet low in protein and devoid of choline. The appearance of the cirrhotic changes could be delayed or prevented by adding choline or its precursor methionine to the offending diet (György and Goldblatt, 1941; Webster, 1941; Daft, Sebrell, and Lillie, 1941; Blumberg and McCollum, 1941).

These observations aroused the interest of many clinical groups, and in the last few years methionine, choline, inositol, and combinations of these substances have been tested in the treatment of chronic liver disease. Owing partly to the inherent difficulty of controlling clinical studies adequately, and partly to the fact that the basal diets contained large and perhaps optimal amounts of the lipotropic factors, the results obtained are difficult to assess. In cirrhosis, especially that associated with alcoholism, it has been reported by a number of investigators (Russakoff and Blumberg, 1944; Barker, 1945; Beams, 1946; and others) that the addition of choline to the diet exerts a favourable clinical effect. The failure of this treatment to produce any beneficial result has also been reported (Yater, 1943; Man, Kartin, Durlacher, and Peters, 1945; and others).

It is well established that in animals in which cirrhosis has been produced by certain chemicals the hepatic lesions may disappear when the administration of the toxic agent is discontinued and a normal diet is supplied. It should be appreciated that most diets (whether for human subjects or experimental animals) which contain a large proportion of natural foodstuffs will provide adequate amounts of the lipotropic agents.

Cameron and Karunaratne (1936) observed that in cirrhosis produced by carbon tetrachloride extensive

regeneration of liver cells and return to a normal configuration occurred within a few weeks after the administration of carbon tetrachloride was discontinued. Edwards and Dalton (1942) made similar observations on the regression of the cirrhotic process in the livers of mice receiving carbon tetrachloride when its administration was stopped and a normal diet provided. Lowry *et al.* (1941, 1945) found that when rats with dietary (choline deficiency) cirrhosis were treated with choline, or given a high casein diet, removal of fat from the liver occurred and extensive regeneration of liver cells took place. Comparative studies on the influence of the various lipotropic agents in the repair of cirrhotic lesions produced by carbon tetrachloride have not hitherto been reported.

The results of the present studies demonstrate again the importance of the lipotropic factors in the repair process, and serve to indicate the relative merits of dietary protein, methionine, choline, and inositol, alone or in combination, in the treatment of carbon tetrachloride cirrhosis in albino rats under certain experimental conditions.

Production of Cirrhosis.—Cirrhosis produced by prolonged administration of carbon tetrachloride has some advantages for this type of study. It can be produced consistently and relatively rapidly, and resembles human portal cirrhosis in many respects (Lamson and Wing, 1926; Moon, 1934). In our first series of experiments the cirrhosis was produced by exposing rats to carbon tetrachloride vapour, and in the second by administering a solution of this substance orally.

Experiment 1

Male white rats (average weight 136 g.) of the Wistar strain were exposed to an atmosphere containing approximately 1,000 p.p.m. of carbon tetrachloride for four hours a day, five days a week, until moderate cirrhosis had been produced. This usually required from 25 to 35 exposures. The capacity of the exposure chamber was approximately 1,000 litres, and an air flow of 250 to 300 litres a minute was maintained.

The rats were kept in group cages and were supplied with a commercial stock diet ("purina fox chow") and fresh water. The 164 animals surviving the exposures were divided into 13 groups. One group of 20 animals was sacrificed immediately to provide initial histological and chemical data. One group (seven animals), to serve as a control, was placed on a hypolipotropic basal diet containing gelatin 12%, casein 8%, beef

*Now in the Department of Pharmacology, University of Toronto.

dripping 10%, corn oil 2%, salts 5%, cellu flour 2%, vitamin powder 1%, sucrose 59.75%, cod-liver oil concentrate 0.015%, cystine 0.24%. This amount of cystine was added in order to make the total organic sulphur content approximately equivalent to that of a diet containing 20% casein.

The vitamin powder was of such a composition that the intake per 10 g. of diet was as follows: thiamin hydrochloride 50 μ g., riboflavin 25 μ g., pyridoxin hydrochloride 20 μ g., calcium pantothenate 100 μ g., and niacin 100 μ g. The cod-liver oil concentrate supplied at least 300 i.u. of vitamin A and at least 75 i.u. of vitamin D per 10 g. of diet.

Ten different diets were prepared by adding to the basal ration various lipotropic supplements, as listed below. The quantities varied as the amounts in grammes per 100 g. of final mixture, the supplements in each case replacing an equal weight of sucrose in the basal ration: (1) casein 7, arachin 5, fibrin 3; (2) casein 7, arachin 5, fibrin 3, choline chloride 0.3; (3) casein 30; (4) casein 22, arachin 5, fibrin 3; (5) choline chloride 0.3; (6) methionine 0.9; (7) inositol 0.3; (8) choline chloride 0.3, inositol 0.3; (9) methionine 0.9, inositol 0.3; (10) choline chloride 0.3, inositol 0.3, methionine 0.9. There were 20 animals in Group 5, seven in Groups 3, 4, and 6, and 13 in all the others.

A final group (20 animals) continued to receive the laboratory stock diet (purina fox chow), which is referred to as Diet 11. All diets, with the exception of the last-mentioned, were essentially isocaloric. While receiving dietary treatment the animals were kept in individual cages. Each rat was offered 15 g. of the appropriate diet daily and the actual food consumption was recorded. When the average daily food intake of the group which ate the least (Diet 7) fell to 13 g. a corresponding reduction was made in the food offered to the other groups. After three weeks, and at intervals for the succeeding nine weeks, three or more rats from each group were sacrificed and their livers were removed for chemical and histological examination. Extraction and analysis of liver lipids were conducted as described by Best, Lucas, Patterson, and Ridout (1946).

Experiment 2

A 20% solution of carbon tetrachloride in corn oil was administered to female white rats (average weight 160 g.) of the Wistar strain two or three times a week over a period of four months. A dose of approximately 0.2 ml. of the solution was given to each animal by stomach-tube. During this phase of the experiment the rats were kept in group cages and were offered the stock diet and water ad libitum. They were then divided into five groups of 10 rats each, and one group was sacrificed immediately to secure initial values. The remaining groups were placed on rations the following components of which were common to all four diets: beef dripping 10, corn oil 2, cellu flour 2, salt mixture 5, vitamin powder 1, cod-liver oil concentrate 0.015. After the addition of the supplements listed below, sucrose was added to bring the mixture to 100 parts. Diet 21 contained casein 10%, cystine 0.2%; Diet 22, casein 10%, cystine 0.1%, methionine 0.1%; Diet 23, casein 20%, cystine 0.2%, methionine 0.2%; Diet 24, casein 10%, cystine 0.1%, methionine 0.1%, choline 0.3%. Four rats from each group were sacrificed after two weeks and the remainder after 13 weeks on the test diets.

Results of Experiment 1

In the group killed immediately after exposure to carbon tetrachloride considerable variation in the gross appearance of the liver was observed. Most of the livers were coarsely but evenly granular (Plate, Fig. 1), of a yellowish-brown colour, and of a rubbery consistency. Microscopically, wide bands of fibrous tissue separated the liver cells into pseudo-lobules with no visible central vein. No normal lobular structure was discernible. Some cells were laden with fat, while others showed hydropic or fatty degeneration (Fig. 2). Cellular necrosis, pyknotic nuclei, and haemorrhage into the parenchyma were also present (Fig. 3). Groups of liver cells with mitotic figures could be seen in many lobules. The average fat content, estimated on a pooled sample made up of large representative portions of each liver, was only 6.4% of the wet weight. In microscopical sections stained with Sudan IV the amount of fat

visible suggested a greater deposition of lipid than that demonstrated by chemical analysis.

The livers of rats maintained on the hypolipotropic basal diet became very fatty (20 to 30% of wet weight), and all but one of the animals died within three weeks. The surviving animal, which was sacrificed at three weeks, also showed extensive cirrhotic changes. The kidneys of all these animals showed the dilated tubules and atrophic degenerative "nephrosis" commonly seen in older animals receiving hypolipotropic diets (Hartroft, 1948). When choline (0.3%) was added to the basal diet, either alone (Diet 5) or in combination with the other supplements (Diets 2, 8, and 10), fat disappeared from the liver and a remarkable regeneration of liver cells took place. In rats killed after three weeks on these rations (Diets 2, 5, 8, and 10), parenchymal cells filled the lobules, but only a few mitotic figures could be seen at this time. This probably indicated that maximum regeneration of parenchymal cells had already taken place. A number of circumscribed lobules of deeply staining cells were seen. The flattened cells at the periphery of these lobules produced a whorled appearance. Pressure due to the bands of fibrous tissue may have been responsible for this condition (Fig. 4). The fibrous strands were thinner but appeared to be stained more deeply than in the livers of the controls. This appearance was probably due to "condensation" of the fibrous tissue. In specimens secured after 6, 9, and 12 weeks gradual improvement was noted, so that latterly (12 weeks) the livers were normal in colour and consistency (Fig. 5). Microscopically, many sections of the livers were essentially normal in appearance (Fig. 6). When special stains for fibrous and reticular tissues were applied thin strands were seen forming a network around and through the lobules (Fig. 7).

Supplements of methionine (0.9%, Diet 6), or of a quantity of casein (30%, Diet 3) which added approximately the same amount of methionine, produced favourable and essentially similar results. Equally beneficial effects attended the use of the laboratory stock ration (Diet 11). The gross and microscopical findings observed on all these diets (Nos. 3, 6, and 11) resembled closely those described above for the animals receiving choline.

Supplementation with inositol alone (Diet 7) possibly had a favourable effect on survival of the rats. Although the growth was poor the mortality rate in the group receiving inositol was of the same order as in those receiving supplements of other lipotropic substances. However, with inositol the livers became very fatty (21 to 29%), as shown in Fig. 5 (lower right), and the extent of the fibrosis apparently increased. In the kidneys definite changes similar to those seen in animals suffering from subacute choline deficiency (Griffith, 1941; Hartroft, 1948) were invariably present.

In rats receiving the very high casein supplement (Diet 3) or the mixed protein supplement (Diet 4) the livers were larger than those of the group supplemented with choline alone but did not differ in other important characteristics. The high-casein diet (No. 3) contained about 1,260 mg. of methionine per 100 g. of diet.

A high-protein diet (No. 1), which was lower in methionine than Diets 3 or 4, produced less beneficial results. While the 15% supplement of protein (casein 7%, arachin 5%, fibrin 3%) made a total of 35% protein in this diet, it provided only about 670 mg. of methionine per 100 g. This amount permitted rapid growth, but the fat content of the livers of these rats rose to high levels while this was taking place. It was only when a plateau in the growth curve was reached that the level of liver fat decreased. At three weeks fat was distributed evenly throughout the lobule, but subsequently, as the fat content fell, distribution became limited to the region of the central vein. The livers were large, light brown in colour (Fig. 5, lower left), and firm to cut. Fibrous strands could be seen on the cut surface. It was not surprising, therefore, to find microscopically that fibrous tissue was present in larger amounts than in groups supplemented more adequately with the lipotropic factors.

In the groups receiving adequate choline or methionine (free or combined in protein) the normal columnar appearance of the liver cells was quickly restored, so that no apparent compression of bile canaliculi or sinusoids existed. In the groups not supplemented adequately these structures appeared to be compressed by cells containing fat. There was no evidence that

E. A. SELLERS, C. C. LUCAS, AND C. H. BEST: LIPOTROPIC FACTORS IN EXPERIMENTAL CIRRHOSIS

FIG. 1.—Cirrhotic liver of the type produced after 28 exposures of 4 hours each to carbon tetrachloride, 1,000 p.p.m.

FIG. 2.—Cirrhotic liver (H. & E. $\times 180$). Fibrous trabeculae separate liver parenchyma into pseudo-lobules one-half size of normal lobule. Some hepatic cells laden with fat.

FIG. 3.—High power of same cirrhotic liver (H. & E. $\times 700$) shows nuclear degeneration and a mitotic figure (M). Vacuolation of cytoplasm indicates site of fatty deposits.

FIG. 4.—Liver from rat on Diet 5 (hypolipotropic basal diet plus choline) after 3 weeks' dietary treatment. Lobule of irregular shape made up of hyperchromatic cells is illustrated. Fat has disappeared from hepatic cells and extensive regeneration has taken place. Decrease in fibrous tissue is already apparent.

FIG. 5.—Illustrates appearance of livers after 12 weeks' dietary treatment. Upper: liver normal in appearance, as from animals receiving supplements of choline or its precursors. Lower left: large pale liver with surface irregularity, from Group 1 (high-protein diet of relatively low methionine content). Lower right: fatty cirrhotic liver from Group 7 (inositol supplement).

FIG. 6.—Photomicrograph of liver of rat on Diet 5 (choline supplement, 12 weeks) (H. & E. $\times 370$). Vacuolated appearance of cytoplasm revealed by special staining to be due to glycogen; fibrous tissue scarcely discernible.

FIG. 7.—Section shown in Fig. 6 stained to reveal fibrous tissue (Masson $\times 370$); a thin strand can be seen in the illustration.

FIG. 8.—Appearance of livers of rats of Experiment 2 after dietary treatment for 13 weeks. Upper left: cirrhotic liver of rat fed Diet 21. Upper right: dietary methionine, increased but still inadequate (Diet 22), effects some reduction in liver fat but no improvement of cirrhotic condition. Lower left: protein supplement enriched with organic sulphur (Diet 23, methionine adequate) results in liver of normal size and colour but with slightly irregular surface. Lower right: liver of rat fed Diet 24 (*i.e.*, Diet 22 supplemented with choline) appears normal.

combinations of the lipotropic substances produced either more rapid or more complete repair in this type of cirrhosis than did adequate intake of either choline or methionine alone.

Chemical estimation of total liver lipids on large portions of individual livers of rats killed after 3, 6, 9, and 12 weeks on the various diets revealed that no significant increase in liver fat occurred except in the rats fed the hypolipotropic basal diet and in the group of animals receiving a supplement of inositol alone (Diet 7). The transitory increase of liver fat in the rats receiving Diet 1 (high protein, of relatively low methionine content) has already been mentioned.

Results of Experiment 2

In this series, in which carbon tetrachloride had been administered orally, the livers of animals killed after receiving the low protein Diets 21 and 22 for only two weeks appeared somewhat altered—the fibrosis had apparently diminished to some extent but the fat content had increased. The apparent decrease in fibrosis may be largely attributed to condensation of the fibrous bands. Much more lipid had accumulated in the livers of the animals on Diet 21 than in those on Diet 22, in which a small (but obviously inadequate) amount of methionine was supplied (see Table). The liver lipids increased slowly in

12% gelatin fulfilled these requirements. This protein mixture is, of course, low in both organic sulphur and the lipotropic factors. As previously stated, cystine (0.24%) was added to correct, in large part, the sulphur deficiency. A deficiency of choline and its precursor methionine in the resulting basal diet is thus the factor limiting growth and survival. To this basal ration supplements of choline, methionine, inositol, proteins poor in methionine, and proteins well supplied with methionine were added, singly or in combination, as described above.

It has already been noted that on the low-methionine choline-free basal diet death ensued within a few weeks. Necropsy revealed that the fibrosis had persisted, and extensive deposition of fat in the liver, and kidney damage, were observed. These findings in the liver were, of course, not unexpected, as it has been shown that cirrhosis may be produced, without exposure to noxious agents, by feeding somewhat similar diets.

Supplements of choline alone (Diet 5) or of methionine alone (Diet 6) in appropriate amounts permitted repair to take place to an amazing extent. The results confirm findings of others already cited—that when an adequate diet is administered to rats with a moderate degree of cirrhosis a remarkable regeneration of liver cells occurs. The presence of choline or its precursor (methionine) in adequate amounts is essential for the removal of fat and the regeneration of parenchymal cells. Under these conditions diminution in amount of fibrous tissue is also observed. The effect of methionine is presumably due to its contribution of labile methyl groups for the biosynthesis of choline.

A supplement of inositol alone (Diet 7) had no favourable effect on the deposition of liver fat or on fibrosis. The apparently lower mortality in this group compared with that in the controls may indicate that inositol exerts some beneficial effect in a manner not yet understood.

A 30% casein supplement (Diet 3), making 50% total protein in the diet, or a mixed protein supplement (Diet 4) produced results as favourable as did diets supplemented with choline or methionine. These particular high-protein diets contained sufficient methionine for growth and maintenance and also for lipotropic activity. Another high-protein diet (No. 1), containing 35% total protein but lower in methionine content, permitted good growth. The livers from this group were, however, abnormally large and fatty during the period of more rapid growth. Although regeneration of parenchymal cells was evident, fat and fibrous tissue strands were seen on gross and microscopical examination even after 12 weeks of dietary treatment. In

Table showing Dietary Conditions and Results of Experiment 2

Group and Diet No.	Protein Content and Supplement	Duration (Weeks)	Degree of Fibrosis*	Total Liver Lipids % (wet wt.)	No. of Livers Examined
Control group (before treatment)			3-4	7.5	10
21	Casein 10% + Cystine 0.2%	2	2-3	24.6	4
		13	3-4	26.7	5
22	Casein 10% + Cystine 0.1% + Methionine 0.1%	2	2-3	10.8	4
		13	3-4	17.7	5
23	Casein 20% + Cystine 0.2% + Methionine 0.2%	2	2-3	6.5	4
		13	1	9.7	5
24	Casein 10% + Cystine 0.1% + Methionine 0.1% + Choline 0.3%	2	2-3	5.3	4
		13	1	7.8	5

* The degree of fibrosis in this series was graded arbitrarily from 1 to 4. Grade 1 = a slight increase above the normal amount of fibrous tissue, usually demonstrable only by special staining. Grade 4 = Extensive fibrosis involving 30% or more of every low-power ($\times 100$) field.

the rats on Diet 22 and eventually reached a moderately high level. The picture changed when the animals were maintained on these rations (Diets 21 and 22) for 13 weeks. The livers were obviously fatty and appeared coarsely granular or nodular (Fig. 8). A progression of the fibrotic process had occurred, so that the degree of fibrosis ultimately resembled that of the control group. Clearly, the cirrhotic lesions were not affected favourably by the low-protein diets.

The gross appearance of the livers of the rats receiving relatively high protein (Diet 23) or choline and methionine (Diet 24) was essentially normal. Microscopically and chemically it was shown that Diet 23 (20% casein supplemented with 0.2% cystine and 0.2% methionine) decreased the liver fat and caused diminution of the fibrosis as efficiently as did Diet 24 (10% casein supplemented with 0.1% cystine, 0.1% methionine, and 0.3% choline).

Discussion

All the experiments described in this study are concerned with the dietary treatment of young adult white rats in which a moderate degree of cirrhosis had been produced by administration of carbon tetrachloride.

In Experiment 1 the choice of protein for the basal mixture to which the test supplements were added was influenced by the desire to keep the amount of methionine in the diet at the lowest level that would maintain weight in the young adult rats. Beveridge, Lucas, and O'Grady (1944) had shown that a ration containing 8% casein and

Diet 1; high protein, relatively low methionine. The increases in body weight, liver weight, and total liver lipid content are plotted against time. A fall in total liver lipid content did not take place until a plateau in the weight curve had been reached.

the Chart the total liver lipids are seen to increase until a plateau is reached in the weight curve, after which a rapid decrease occurs. These data, and others to be published, suggest that when limited amounts of methionine are available it is used preferentially, but not exclusively, to meet growth requirements. The methionine content of this diet, which was insufficient for lipotropic action during the period of rapid growth, was adequate for removal of liver fat after rapid growth had ceased. While all essential amino-acids were present in Diet 1 in considerably greater amounts than are usually considered adequate, methionine was present in a proportionately smaller amount. Diet 1 contains about 670 mg. of methionine per 100 g., a quantity in excess of that usually accepted as optimal for growth. Since liver lesions persisted which would presumably have been eliminated by more methionine the amount supplied was obviously inadequate under these conditions. If sufficient methionine had been available to supply labile methyl groups for the biosynthesis of adequate amounts of choline the liver damage should have been repaired. Indeed, when choline was added to Diet 1 (Diet 2) the livers of the animals appeared normal on gross examination and nearly so microscopically. These observations illustrate the important nutritional interrelationship of methionine and choline, which will now be discussed briefly.

Methionine contributes to several metabolic requirements. The significance of the fact that these different metabolic roles exist is not always sufficiently appreciated. First, certain minimal quantities have been shown to be necessary, presumably as intact molecules, for maintenance and growth. Secondly, methionine serves as a source of organic sulphur for the formation of cystine and other sulphur-containing metabolites. Thirdly, it possesses a labile methyl group, and it is this characteristic which is of greatest interest in the study of lipotropic phenomena. Dietary choline cannot substitute for methionine in either of the first two roles. Thus in any diet choline can replace only that portion of methionine which is concerned with lipotropic phenomena; conversely, methionine supplements are not available for lipotropic purposes to a maximal extent until the first two (apparently more essential) roles of methionine have been filled completely. It would appear that in rats it is only under these conditions that dietary choline and methionine are maximally interchangeable. Moreover, not all dietary choline (or choline synthesized *in vivo* from dietary labile methyl groups) is used for lipotropic purposes, since it is obvious that some is utilized to satisfy structural and other needs. Only when the moieties of dietary choline and methionine that are used for lipotropic purposes have been assessed will it be possible to estimate a factor representing their lipotropic interchangeability in any diet. Similarly, cystine and methionine are interchangeable nutritionally to a maximal extent only when the minimal methionine requirement (i.e., as intact molecule) has been met and an adequate supply of choline (or dietary labile methyl groups) is present—i.e., when the deficiency involves only the supply of dietary organic sulphur. These interrelationships have recently become more clearly defined. Neglect to consider them has led to some confusion concerning the nutritional roles (particularly with regard to lipotropic phenomena) of choline, methionine, and cystine under different dietary conditions.

The apparent disappearance of fibrous tissue from the cirrhotic livers of animals which received the lipotropic agents deserves some further comment. After 12 weeks of treatment, in groups supplemented with choline or methionine, fibrous tissue could often be demonstrated only by using special stains. Decrease in volume of individual cells may have accounted for much of this apparent disappearance, yet the change from grotesque pseudo-lobula-

tion produced by the fibrous tissue to a more or less normal structure raises the question of actual removal of fibrous-tissue cells. This latter possibility is perhaps supported by the decrease in the number of nuclei of fibrous-tissue cells which can be seen. It should be emphasized that in our experiments the cirrhotic lesions which have been so strikingly affected by the lipotropic agents were of moderate degree and had not been present for long periods. A study of the action of the lipotropic agents on more advanced cirrhosis of long standing may be indicated, since clinicians are frequently confronted with this problem.

Certain potential clinical applications of the knowledge gained from the work on lipotropic factors, including that reported here, may be discussed briefly. Many clinical groups, stimulated in part by the findings of Patek and Post (1941), now use diets rich in protein and vitamins in the treatment of patients with cirrhosis. The favourable results which have been reported would appear to be attributable in large part to the occurrence of choline and methionine in these diets. We are not minimizing the obvious importance of inadequate amounts of minerals, other vitamins, and essential amino-acids. Estimations made in our laboratory and elsewhere show that normal diets may contain from 300 to 900 mg. of choline daily, and it may be calculated that the same rations would provide from 2 to 4 g. of methionine. If from one-fourth to one-third of this methionine were used for lipotropic purposes an additional 150 to 400 mg. of choline might be formed from it. Thus a normal diet may provide up to 1,300 mg. of available choline daily. The experimental evidence suggests that no beneficial effect in the treatment of cirrhosis should be expected from the addition of choline or methionine to a ration which is already adequately supplied with these factors—unless there should be interference with liberation of choline or methionine in the intestine or with the absorption or subsequent physiological actions of these compounds. No clinical conditions have yet been recognized in which the lipotropic actions of choline or methionine are inhibited, but this situation might conceivably exist in various diseases. However, there is convincing experimental evidence (Chaikoff, Entenman, and Montgomery, 1947) that interference with protein digestion by ligation of the pancreatic ducts or by removal of the pancreas decreases the liberation of methionine (and presumably the formation of choline). Excessive deposition of fat in the livers of duct-tied and insulin-treated depancreatized dogs has been observed repeatedly. In the depancreatized animals the progression of the lesion to cirrhosis has been followed (Chaikoff, Connor, Biskind, 1938). The clinician should therefore be on the look-out for signs of hepatic damage in cases of long-standing insufficiency of the external secretion of the pancreas.

Summary and Conclusions

Young adult rats in which a moderate degree of cirrhosis had been produced by the administration of carbon tetrachloride were placed on diets containing different amounts of the following lipotropic agents; choline, methionine, protein (containing methionine), and inositol.

None of the animals receiving only a basal hypolipotropic ration showed improvement in the cirrhosis. Deposition of fat in the liver increased, degenerative renal lesions appeared, and six of the seven rats in this group survived less than three weeks.

When adequate amounts of pure choline chloride or *DL*-methionine were added to the basal hypolipotropic diet a remarkable improvement in the gross appearance of the livers was noted within two or three weeks. After two or three months of this treatment the livers appeared normal, and on microscopical examination special stains were usually required to demonstrate any abnormal deposition of fibrous tissue. The

processes by which this fibrous tissue may disappear have been discussed briefly.

A high-protein diet, containing 35% protein but low in methionine, was not as beneficial as a ration containing 20% of similar protein to which adequate supplements of choline or methionine had been added.

A high-protein diet which contained an adequate amount of naturally occurring methionine produced results as favourable as those which were secured when suitable amounts of choline or methionine were added to diets of lower protein content.

While the experimental evidence indicates that the presence of the lipotropic agents is essential for the repair of the damaged liver under the conditions of our experiments, it does not suggest that the addition of choline or its precursor, methionine, to diets already containing adequate amounts of these factors will enhance the therapeutic effect unless some interference with the absorption or action of the lipotropic agents exists.

No evidence has been secured that inositol exerts a favourable effect on experimental cirrhosis.

On the basis of the experimental findings it is suggested that the favourable clinical results which have been obtained in patients with liver damage when an effective therapeutic diet is provided may be attributable in large part to the lipotropic factors which the diet supplies.

We wish to express our thanks to our colleagues Dr. Jessie H. Ridout and Dr. W. Stanley Hartroft for their generous assistance in certain aspects of this work. The expenses of these studies were defrayed in part by a grant from the Banting Research Foundation.

REFERENCES

- Barker, W. H. (1945). *Med. Clin. N. Amer.*, **29**, 273.
 Beams, A. J. (1946). *J. Amer. med. Ass.*, **130**, 190.
 Best, C. H., Lucas, C. C., Patterson, J. M., and Ridout, J. H. (1946). *Biochem. J.*, **40**, 368.
 — and Ridout, J. H. (1935). *J. Physiol.*, **84**, 7P.
 Beveridge, J. M. R., Lucas, C. C., and O'Grady, M. K. (1944). *J. Biol. Chem.*, **154**, 9.
 Blumberg, H., and McCollum, E. V. (1941). *Science*, **93**, 598.
 Cameron, G. R., and Karunaratne, W. A. E. (1936). *J. Path. Bact.*, **42**, 1.
 Chaikoff, I. L., and Connor, C. L. (1940). *Proc. Soc. exp. Biol. N.Y.*, **43**, 638.
 — Entenman, C., and Montgomery, M. L. (1947). *J. Biol. Chem.*, **168**, 177.
 Daft, F. S., Sebrell, W. H., and Lillie, R. D. (1941). *Proc. Soc. exp. Biol. N.Y.*, **48**, 228.
 Drill, V. A., and Loomis, T. A. (1947). *J. Pharmacol.*, **90**, 138.
 Edwards, J. E., and Dalton, A. J. (1942). *J. nat. Cancer Inst.*, **3**, 19.
 Griffith, W. H. (1941). *Biol. Symposia*, **5**, 193.
 — and Wade, N. J. (1939). *J. Biol. Chem.*, **131**, 567.
 György, P., and Goldblatt, H. (1939). *J. exp. Med.*, **20**, 185.
 — (1941). *Proc. Soc. exp. Biol. N.Y.*, **46**, 492.
 Hartroft, W. S. (1948). Personal communication.
 Lamson, P. D., and Wing, R. (1926). *J. Pharmacol.*, **29**, 191.
 Lowry, J. V., Ashburn, L. L., and Sebrell, W. H. (1945). *Quart. J. Stud. Alcohol*, **6**, 271.
 — Daft, F. S., Sebrell, W. H., Ashburn, L. L., and Lillie, R. D. (1941). *Publ. Hlth Rep. Wash.*, **56**, 2216.
 Man, E. B., Kartin, B. L., Durlacher, S. H., and Peters, J. P. (1945). *J. clin. Invest.*, **24**, 623.
 Moon, V. H. (1934). *Arch. Path.*, **18**, 381.
 Patek, A. J., jun. (1937). *Proc. Soc. exp. Biol. N.Y.*, **37**, 329.
 — and Post, J. (1941). *J. clin. Invest.*, **20**, 481.
 Russakoff, A. H., and Blumberg, H. (1944). *Ann. intern. Med.*, **21**, 848.
 Webster, G. (1941). *J. clin. Invest.*, **20**, 440.
 Yater, W. M. (1943). Discussion of paper by C. H. Greene, *J. Amer. med. Ass.*, **121**, 720.

When Mr. Bevan opposed the proposal to limit the opening of new pharmaceutical units it was dropped. "This half-hearted attitude does not augur well for the future," Mrs. Jean K. Irvine, President of the Pharmaceutical Society of Great Britain, told its annual general meeting. "The consequences remain to be seen, but one of the main causes of unsatisfactory conditions in pharmacy will continue to operate and, other things being equal, will have the same results." The proposal that the manager of a pharmacy as well as the owner should participate in the contract under the National Health Service had also not been further pursued, she said, when the Minister indicated his opposition. The manager was dependent on his employer for furthering his point of view, and it was not necessary to attribute a hostile attitude to employers to realize how unsatisfactory it was that approximately half the pharmacists taking part in the Service would be denied the status their qualification and responsibility deserved.

PNEUMOCONIOSIS OF COAL-MINERS*

BY

C. M. FLETCHER, M.D., F.R.C.P.

Director, Pneumoconiosis Research Unit, Medical Research Council

[WITH PHOTOGRAVURE PLATE]

The investigations of the Pneumoconiosis Research Unit have of necessity had a practical or *ad hoc* bias. There are many fundamental points of pathogenesis, of physiology, of cellular pathology, and so forth which remain obscure in coal-miners' pneumoconiosis, but, initially at any rate, we have had to set these points on one side and to seek information in the light of which appropriate administrative action might be undertaken in order to deal with the vast human and economic problems caused by the disease.

In South Wales, by the end of 1945, 12,000 men had been certified and new cases were still being certified at the rate of 100 a week. Nearly two-thirds of them were under 50 years old; one-quarter were under 40. The majority were coal-hewers, the most skilled class of mine workers. They found themselves, at an age when family responsibilities are at their heaviest, cast out of their accustomed work, with no training for alternative employment, and with weekly compensation amounting at most to £3 5s. for an unemployed man with a wife and three children. If the man obtained work his weekly compensation was reduced according to his earnings. No special system of treatment or after-care existed.

Post-mining Employment

Our first investigations, carried out under the supervision of Dr. Alice Stewart, were directed to studying these social problems. We asked the Social Survey to assist us, and during February, 1946, they carried out an inquiry, by house-to-house visits, into the post-mining employment of a one-in-ten sample of all the men who had been certified in South Wales before January, 1945. We also obtained a full employment history from the men whom we invited to attend for the clinical and radiological examinations which will be considered later. Lastly, we instituted inquiries at factories which were employing men with pneumoconiosis† in order to study the industrial performance of these men.

Generalization concerning the re-employment of men with pneumoconiosis is difficult because of the wide variety of the types of case to be considered. They range from young men with minimal disability to elderly men who are breathless at rest. However, 86% of the cases fall between the ages of 30 and 60, when any man should have a job, and there are two other important factors common to nearly all of them. They have all spent their adolescent and early adult years in coal-mining, at which they have become skilled, but they have no training for other forms of skilled employment, and they are, by reason of their breathlessness, incapable of undertaking strenuous physical exertion.

Bearing these points in mind, we may consider certain relevant statistics concerning the post-mining employment of certified cases of pneumoconiosis. Chart IV shows the percentage of the men in the Social Survey's sample of certified cases of pneumoconiosis and among all insured men and boys in South Wales who were employed each year between 1936 and 1946. It will be seen that under

*The second Goulstonian Lecture, given at the Royal College of Physicians of London, on Jan. 15. Lecture I appeared in last week's issue.

†In this paper the word pneumoconiosis includes silicosis.