REFERENCES

- Fishberg, A. M. (1934). *Hypertension and Nephritis*. Lea and Febiger, Philadelphia.
 Foa, P. P., Foa, N. L., and Peet, M. M. (1943). *J. clin. Invest.*, **22**, 727.
 Goldblatt, H. (1938). *J. exp. Med.*, **67**, 809.
 Jablons, B. (1944). *Rev. Gastro-enterol.*, **11**, 246.
 Klemperer, P., and Otani, S. (1931). *Arch. Path.*, **11**, 60.
 McFarlane, D. (1944). *Stain Technology*, **19**, 29.
 Moritz, A. R., and Oldt, M. R. (1937). *Amer. J. Path.*, **13**, 679.
 Morlock, C. G. (1939). *Arch. intern. Med.*, **63**, 100.
 Pickering, G. W., and Prinzmetal, M. (1938). *Clin. Sci.*, **3**, 357.
 Wilson, C., and Byrom, F. B. (1939). *Lancet*, **1**, 136.
 — and Pickering, G. W. (1938). *Clin. Sci.*, **3**, 343.

SILICEOUS GRANULOMA DUE TO TALC

A CAUSE OF POST-OPERATIVE PERITONEAL ADHESIONS

BY

W. ARTHUR MACKEY, M.B., Ch.B., F.R.F.P.S.G.
F.R.C.S.Ed.

AND

J. B. GIBSON, M.B., Ch.B.

(From the Departments of Surgery and Pathology of the
University of Glasgow at the Western Infirmary, Glasgow)

[WITH PHOTOGRAVURE PLATE]

Post-operative peritoneal adhesions have long excited the interest of surgeons and stimulated speculation about their pathogenesis. One is often surprised at operation by the scantiness, softness, and avascularity of intraperitoneal adhesions seen after the resolution in an unopened abdomen of a bacterial inflammation such as an appendicular abscess or a more generalized peritonitis. On the other hand, if it be necessary to reopen the abdomen after an aseptic laparotomy, adhesions may be found that are dense, vascular, and widespread. These findings suggest that laparotomy may on occasion introduce into the abdomen a substance more persistently irritant than bacteria. Observations over the past 15 years have identified one such irritant as talc—a complex magnesium silicate—the substance used almost universally in the preparation of gloves for operation. Two excellent reviews on the harmful effects of talc in the peritoneal cavity have appeared in the past year in America (Eiseman, Seelig, and Womack, 1947) and in this country (Roberts, 1947).

Within the abdomen the irritant action of talc elicits a proliferative response comparable to that found in traumatic siliceous granuloma (pseudotuberculoma silicoticum) (Shattock, 1916; Faulds, 1935). The effects are not limited to the peritoneum, and the implantation of talc at laparotomy may be followed by granulomatous formation in the scar, by sinuses or faecal fistulae, stenosis of entero-anastomotic stomata, and, as Roberts has shown, by chronic fibrosis of the Fallopian tube with consequent sterility. The patient's history and the density of the peritoneal adhesions found at operation may suggest the diagnosis of siliceous granuloma, but the only certain proof is to find, in microscopical sections of the collagenous tissue, doubly refractile material with the appearance of talc closely related to foreign-body giant cells. The use of "polaroid" screens on the ordinary microscope is simple and usually leaves little doubt about the diagnosis, but, unless sections are examined for anisotropic material, cases may easily escape recognition and be diagnosed as tuberculosis, regional enteritis, sarcoidosis, keloid scar, or non-specific foreign-body reaction, although the situation of the granuloma may well occasion comment.

Foreign bodies such as suture material are often surrounded by giant cells and can show in varying degrees

the property of rotating the plane of polarized light, but they are unlikely to be scattered diffusely in the peritoneum, and the structure of talc as seen by polarized light is fairly characteristic (see Plate, Fig. 3). Many years after its implantation in the tissues talc has been recognized in sections prepared by routine histological methods (Roberts, 1947).

Case History

Miss B., aged 74, was admitted to the medical wards of the Western Infirmary on Oct. 3, 1947, for investigation of a severe anaemia which had developed over the past four years. She was found to be passing considerable quantities of altered blood in the stools, but intensive clinical, haematological, and radiological investigation failed to establish a source. Transfusion of 9 pints (5.1 litres) of blood over a period of a month scarcely replaced the loss, and her condition remained precarious. The decision to perform laparotomy was taken in the hope of finding the bleeding-point. She had undergone no previous abdominal operation.

First Operation.—On Nov. 26 Prof. C. F. W. Illingworth made a right paramedian incision in the upper abdomen, and his note states that wide exploration of the abdomen was carried out. A Meckel's diverticulum was found attached by its fundus immediately deep to the umbilicus. The diverticulum was not apparently the source of bleeding and so was not removed, though its tip was freed from the abdominal wall. The spleen was enlarged to perhaps twice normal size. The previous haematological investigations having yielded no facts to contraindicate splenectomy, this was carried out and the arteries at the ends of both curvatures of the stomach were divided in the hope that the undisclosed lesion might be in the upper division of the portal system. The patient recovered uneventfully from this operation, but bleeding into the alimentary tract continued. On several occasions aspiration of the stomach was carried out over long periods, but no blood was recovered. By Dec. 9 the haemoglobin level had fallen to 38% and it appeared necessary to explore the abdomen once more. In the absence of Prof. Illingworth the responsibility for this step fell on one of us (W. A. M.). The patient was prepared by transfusion of 4 pints (2.27 litres) of blood.

Second Operation.—This was performed on Dec. 9, 13 days after the first operation. There were dense adhesions of recent origin involving the stomach, omentum, the Meckel's diverticulum, and the anterior abdominal wall. The diverticulum was removed and opened but no ulceration was found. The entire small bowel contained altered blood; nowhere throughout its length could a polyp or ulcer be palpated, but accurate examination of the stomach was prevented by adhesions. The abdomen was then closed; recovery from the operation was satisfactory, and four days later the haemoglobin value was 82%. Bleeding into the alimentary tract, however, continued as before, and to date is unexplained.

Pathological Report.—The specimen consists of a Meckel's diverticulum 3.5 cm. long. The peritoneum is dull and the wall thick and dense. The mucosa has a normal velvety appearance. Histological sections show zones of thickly woven collagenous connective tissue external to the muscularis and extending into it here and there for a short distance. This tissue replaces most of the subserous fat, is responsible for the increased thickness of the wall, and forms the outer surface of the diverticulum at several places (Plate, Fig. 1). Imbedded in it lie many foci of macrophages and foreign-body giant cells, arranged in some instances in the form of poorly defined follicles. These cells surround brownish crystalline doubly refractile particles of an average length of 15 μ and of a shape identical with surgical talc (Figs. 2 and 3). Similar material is seen underneath the serosa in varying concentrations in all parts of the diverticulum. The mucosa is of normal small-intestine type and no areas of gastric heterotopia are present.

Comment

There is no doubt that these lesions had their origin in the first operation, 13 days before the specimen was removed. It is comparatively rarely that the opportunity occurs to examine tissue removed from the abdomen such

a short time after laparotomy in an aseptic case, and we make this report to emphasize the speed with which dense fibrous tissues may form in response to a small quantity of talc. In the series of cases described by Eisman, Seelig, and Womack several patients complained only a few days after operation of symptoms later ascribed to talc granuloma, but histological examinations were not made in any of these until at least two months after the first operation. In our patient the reaction produced no symptoms despite its extent and intensity.

It is evident that the use of talc at operation constitutes a hazard potentially grave in its consequences for the patient. The authors of the two reviews quoted above recommend bluntly that it be abandoned. The suggested substitutes, potassium bitartrate and formalized starch, appear to be free of the irritant effect of talc, but unfortunately possess faults of their own, and the search for a completely satisfactory lubricant for gloves has not yet been successful. Until it is found we must obviously use talc with caution, make every effort to reduce the quantity to a minimum, and above all avoid depositing it in the peritoneal cavity.

Summary

The production of post-operative peritoneal adhesions by surgical talc is briefly discussed.

A case history is given of a patient whose abdomen was reopened 13 days after laparotomy. At the second operation extensive dense adhesions were found. Histological evidence is presented that these arose as a result of deposition of small amounts of talc at the time of the first operation.

We should like to thank Dr. W. R. Snodgrass and Prof. C. F. W. Illingworth for access to their records and permission to publish this case.

REFERENCES

- Eisman, B., Seelig, M. G., and Womack, N. A. (1947). *Ann. Surg.*, **126**, 820.
 Faulds, J. S. (1935). *J. Path. Bact.*, **41**, 129.
 Roberts, G. B. S. (1947). *Brit. J. Surg.*, **34**, 417.
 Shattock, S. G. (1916). *Proc. R. Soc. Med.*, **10** (Section of Path.), 6.

"TUBERCULOSIS" (COEXISTENT WITH CARCINOMA OF THE UTERUS) SHOWN TO BE SILICEOUS GRANULOMA

BY

G. HARVEY SMITH, M.D.

(From the Department of Pathology, University of St. Andrews and the Royal Infirmary, Dundee)

[WITH PHOTOGRAVURE PLATE]

The present case was previously reported as an example of the rare coexistence of carcinoma and tuberculosis in the same organ (Smith, 1945). The diagnosis of tuberculosis was made on histological grounds without much dubiety, despite failure to demonstrate acid-fast organisms in the sections. Roberts (1947), however, has proved that granulomata occurring in the Fallopian tube, with a histological picture very closely resembling tuberculosis, may in fact be siliceous and not tuberculous: these foreign-body granulomata apparently follow the introduction of talc (surgical gloves) into the peritoneum, and in all his five cases of siliceous granuloma of the tube a previous appendectomy had been performed and could well have been the occasion of implantation.

Restudy of my case in the light of Roberts's work suggests that the original diagnosis of tuberculosis is inaccurate and that of siliceous granuloma rather more likely, the more so as the patient also had undergone a previous appendectomy.

Case History

Mrs. A., aged 44, had a "chronically inflamed appendix" removed in November, 1943; at operation the uterus was noted to be enlarged, but no signs of tuberculosis were observed. In December, 1943, there was a sudden discharge of thick yellow pus from the vagina. The gynaecologist's report was: "Mass palpable through the anterior fornix." In April, 1944, she was readmitted with vaginal bleeding and abdominal pain; at laparotomy the uterus was seen to be enlarged and white nodules were visible in the walls of both tubes at the uterine ends. The uterus and right tube were removed and sent to the laboratory. No sign of pulmonary or renal tuberculosis was detected at the time of hysterectomy; she has been well since then (46 months), and at present shows no signs of tuberculosis or malignant disease. The patient's husband, however, has recently been found to have pulmonary tuberculosis.

Pathology.—The upper part of the uterine cavity is completely occupied by a bulky friable mass of tumour tissue covering most of the posterior wall. Two small, pale, button-like nodules are present on the posterior wall, nearer the cervix. The right Fallopian tube shows a firm white nodule 1 cm. in diameter at the uterine end. Microscopically, the appearances in the uterine mass are those of an anaplastic adenocarcinoma. At the margins of the main growth, and in the two smaller nodules of tumour below, there are follicular structures, histologically tuberculous, with giant cells, areas of necrosis, and surrounding lymphocytes. The sections of the Fallopian tube show that the nodule there is not neoplasm but a further tuberculous-looking granuloma with rather more connective-tissue proliferation.

Examination of sections by polarized light revealed the presence of doubly refractive particles both in the tissues and in a few of the giant cells (Plate, Fig. 2). It also revealed a superficial collection of talc crystals with accompanying round-cell reaction immediately below the serous coat of the tube (Fig. 1). This latter lesion, although not far from the main mass of the granuloma, is highly suggestive of direct implantation of talc on the serous surface of the tube as against transport into the lumen, suggested by Roberts to explain the striking localization of the talc in his cases. It is important to note that of 70 sections of the tube only 14 showed the presence of anisotropic material. This material, as Roberts also points out, when contained within the giant cells is not always acicular in shape. Fifty-six sections of the uterine lesions have also been examined through "polaroid" screens, but no acceptable anisotropic particles have been seen.

Discussion

It is suggested that talc from the surgeon's gloves was introduced into the peritoneal cavity during the appendectomy in November, 1943, and that this is the origin of the siliceous granuloma formed in the Fallopian tube. Although the granulomatous foci in the uterus failed to show anisotropic bodies it seems reasonable to regard them also as siliceous in nature.

The minimum period recorded by Roberts as elapsing between laparotomy and the discovery of a tubal lesion is two years; in the present case the interval was only five months. The discovery of the granuloma was, of course, incidental to the treatment of the cancer, and it is possible that the siliceous lesions would not otherwise have declared themselves for some time, if at all. The more actively necrotic character of the histological changes in my case compared with those described by Roberts may well be due to the earlier stage in development of the granuloma.

The period elapsing between the implantation of silica and the subsequent appearance of a superficial dermal granuloma has generally been lengthy—more so in the accident cases than in those with a lesion in a surgical scar. In most cases there appears to be a latent period, and in one traumatic case mentioned by Roberts the silica, driven mechanically under the skin as a result of a fall, lay there without provoking any noticeable reaction for 43 years. It

SILICEOUS GRANULOMA DUE TO TALC

W. A. MACKEY AND J. B. GIBSON

FIG. 1.—Section of wall of Meckel's diverticulum ($\times 35$). Dense fibrous tissue has replaced subserous fat. Complex of giant cells surrounds talc particles centrally and at serous surface.

G. HARVEY SMITH

FIG. 1.—Partly crossed "polaroid" screening shows mass of doubly refractive material with cellular reaction under serosal surface of Fallopian tube.

FIG. 2.—Syncytial giant-cell reaction enclosing talc particles ($\times 200$).

FIG. 3.—Same field as Fig. 2 seen through partially crossed "polaroid" screens.

FIG. 2.—Partly crossed "polaroid" screening of granulomatous lesion in wall of Fallopian tube. Note histological similarity to tuberculosis. Two giant cells show doubly refractive masses of talc.

WILLIAM WALKER

FIG. 1.—Adhesion between two loops of small intestine, one obviously hypertrophied. Thin collagenous fusion zone only slightly cellular, but giant cells are present ($\times 8$).

FIG. 2.—Partly crossed "polaroid" screening shows multinucleated giant cell containing doubly refractive material, slightly displaced ($\times 250$).

F. R. MAGAREY AND P. H. DENTON: TORULA
HISTOLYTICA INFECTION OF CENTRAL NERVOUS
SYSTEM

FIG. 1.—*Torula histolytica* in section of meninges ($\times 130$).

FIG. 2.—Torula body in subependymal giant cell ($\times 720$).