the blood in infective hepatitis appears to be secondary to the disease process.

The higher incidence of infective hepatitis among officers compared with men cannot be explained on the basis of avitaminosis. It is due to the greater incidence of "jaundice-precipitating factors" among them as already discussed in the previous paper.

Summary and Conclusions

The liver changes observed in fatal cases of infective hepatitis are described in detail. The sequence of these changes is traced from acute necrosis to portal cirrhosis. If death occurs within a fortnight the findings are those of acute necrosis. After a fortnight fibrous-tissue formation becomes apparent, especially at the portal tracts. The pattern is that of portal cirrhosis. In three months' time portal cirrhosis is obvious. In eighteen months the picture is that of portal cirrhosis, both macroscopically and microscopically. Jaundice waxes and wanes according to the intensity of inflammation of the liver cells.

This sequence occurs in less than 1% of cases. In the remaining cases the liver appears to recover ultimately.

Homologous serum jaundice is not a separate entity. The difference between apparent and real incubation periods is pointed out.

My thanks are due to Colonel Chandra, Colonel Lambert, Lieutenant-Colonel Daukes, and Lieutenant-Colonel Thomson for permitting this work to be carried out in the hospitals under their command; to Captain Ross and Captain Krishnaswami for valuable help in providing slides and records; and to Mr. Dodds for the photomicrographs.

BIBLIOGRAPHY

- Apley, J., and Wallis, H. R. E. (1948). *British Medical Journal*, 2, 197.
British Medical Journal. Annotation, Nov. 4, 1944, p. 602.
 Cameron, J. D. S. (1943). *Quart. J. Med.*, 12, 139.
 Harris, A. D., and Moore, T. (1947). *British Medical Journal*, 1, 553.

TORULA HISTOLYTICA INFECTION OF CENTRAL NERVOUS SYSTEM

BY

F. R. MAGAREY, M.D., M.R.C.P., M.R.A.C.P.

Department of Pathology, Welsh National School of Medicine

AND

P. H. DENTON, M.D., M.R.C.P.

Senior Medical Registrar, Cardiff Royal Infirmary

[WITH PHOTOGRAPHIC PLATE]

Tuberculous and torular meningitis give rise to very similar clinical pictures and to almost identical changes in the cerebrospinal fluid, so that the two diseases are not easily distinguishable. The clinical trials of streptomycin now being carried out make it important to differentiate between them. Levin (1937) summarized about 70 cases of torular meningitis, but the latest and most complete review of the disease is the monograph by Cox and Tolhurst (1946). Large numbers of cases have now been recorded, especially in America, Australia, and South Africa, but so far only three in Great Britain. Smith and Crawford (1930), in Harrogate, described the first case, which was diagnosed post mortem. Gelatinous swellings in the hypothalamic region and thickening of the basal and spinal meninges were found in which torula bodies were recognizable histologically. Greenfield, Martin, and Moore (1938), in London, recorded the second, which was also diagnosed post mortem. An adhesive meningitis around the basal cistern and pons was found, and torula bodies could be seen histologically in the meninges and in the perivascular spaces of the basal ganglia. Their presence was associated with granulomatous infiltration but no

caseation. Blair (1943), also in London, described the third case, which was apparently diagnosed during life by the finding of torula bodies in the cerebrospinal fluid. No details of the chemical or cytological changes were included. At necropsy the subarachnoid space over each cerebral hemisphere, together with the lateral ventricles, was filled with yellowish gelatinous material. A case has recently come to our notice in Wales.

Case Report

A clergyman aged 60 was admitted to hospital on July 26, 1947, having had severe headaches for ten weeks, with lassitude and loss of appetite. The headaches at first occurred at intervals of one or more weeks, but later became more frequent and were associated with vomiting and sometimes with blurring of vision and tinnitus in the left ear. The only abnormal sign on admission was a doubtful left extensor plantar reflex. Lumbar puncture yielded clear cerebrospinal fluid under normal pressure, with 66 lymphocytes per c.mm. and 95 mg. of protein and 684 mg. of chlorides (estimated as NaCl) per 100 ml., the Lange colloidal gold and Wassermann reactions being negative. Other investigations, including radiography of skull and chest, were negative. The headaches gradually disappeared and the patient was discharged at his own request, free from symptoms, three weeks after admission.

He was readmitted on Sept. 2, the headaches having recurred three days after he left hospital and become continuous, with daily vomiting. There was now slight mental retardation. His tendon reflexes in the right arm were brisker than those in the left, but both plantar reflexes were now flexor. There were no other significant physical signs. On Sept. 4 the cerebrospinal fluid was under increased pressure (220 mm.), but there was now only one lymphocyte per c.mm. The proteins had, however, increased to 150 mg., whilst the chlorides had fallen to 620 mg. per 100 ml. After admission his condition improved, but by Sept. 11 the headaches were again severe and stiffness of the neck and slight Kernig's sign were present. His right knee-jerk was brisker than the left, but the arm reflexes were equal. The cerebrospinal fluid pressure was now 300 mm. and the cell count less than one per c.mm. The proteins were 180 mg. and the chlorides 670 mg. per 100 ml. Ventriculography on Sept. 18 showed no abnormality. During the next month there was periodic mental confusion with drowsiness, and on Oct. 7 a left external rectus paresis and a right-sided papilloedema were noted. The cerebrospinal fluid findings on Oct. 22 were as follows: pressure, 180 mm.; cells, 25 per c.mm. (95% lymphocytes, 5% polymorphs); protein, 250 mg.; chlorides, 626 mg., and sugar 16 mg. per 100 ml. The colloidal gold curve was 3442221000. From this time onwards the patient's condition deteriorated, wasting appeared, and by Nov. 6 the right pupil was larger than the left and well-marked left hemiparesis was present. He passed gradually into coma and died on Nov. 13, six months after the onset of his symptoms. The whole illness was almost completely afebrile until the last four days.

Necropsy Findings.—Along the basilar artery and in some of the sulci over the cerebral hemispheres a few flakes of fibrinopurulent exudate were found. The pineal gland was imbedded in a yellowish gelatinous substance which filled the pial-arachnoid space in that area. In the right side of the pons there was an area of softening about 1.5 cm. in diameter. The lateral ventricles contained slightly cloudy fluid, but were not dilated. There was an area of apparent softening beneath the ependyma of the right lateral ventricle, involving the adjacent portion of the caudate nucleus. Another area of softening involved the anterior portion of the left external capsule.

Hypostatic pneumonia was present in the lower lobes of both lungs, but there were no significant lesions in any other organ.

Histology.—Sections of the meninges showed granulomatous lesions, chiefly around the large blood vessels. The granulomata consisted of central caseation surrounded by cellular zones in which there were small round cells, polymorphs, and pale endothelial cells, some of them large and multinucleated. The lesions closely resembled tubercles, but scattered throughout

them the highly characteristic *Torula histolytica* could be seen. These are round or oval yeast-like organisms surrounded by clear zones which represent their gelatinous capsules (Plate, Fig. 1). The bodies were in some parts clustered together, and were so numerous as to exclude other cells. Their staining properties were variable, some taking a pale blue with haemalum (as in Fig. 1), and with Gram stain a rather nondescript brownish colour. (We were unable to stain the capsules, and no budding was seen.) The lesions in the pons and left external capsule were areas of simple softening, but the subependymal lesion in the cerebrum was a torular granuloma. It differed from the meningeal lesions in that endothelial cells, especially the giant cells, were more prominent, while there was no caseation. Occasionally a torula body could be seen in a giant cell (Fig. 2). The vessels in the surrounding brain tissue showed perivascular lymphocytic cuffing.

Discussion

During life this case was thought to be either a cerebral tuberculoma or tuberculous meningitis. This diagnosis was supported by the decreased chloride and sugar content of the cerebrospinal fluid and by the normal ventriculogram, although repeated search for tubercle bacilli was unsuccessful. As the possibility of torulosis was not considered, the torula bodies were not specifically looked for, but they may well have been present and overlooked in the cytological examination, for they are easily mistaken for red blood cells or lymphocytes. If present in the cerebrospinal fluid they can be demonstrated by mixing the centrifuged deposit with a drop of indian ink. In a smear or wet preparation they then appear as small refractile bodies surrounded by broad colourless haloes.

There were certain features which, had they been appreciated at the time, might have helped in the diagnosis of this case. The illness lasted six months, during which the patient was for variable periods quite symptom-free. Also, except for the last four days when pneumonia was developing, he was almost entirely afebrile. These facts favoured torulosis rather than tuberculous meningitis.

Our thanks are due to Dr. Idris G. Jones for permission to publish the clinical details, and also to Mr. J. P. Napper for his technical assistance with the photomicrographs.

REFERENCES

- Blair, D. (1943). *J. ment. Sci.*, **89**, 42.
Cox, L. B., and Tolhurst, J. C. (1946). *Human Torulosis*, Melbourne University Press.
Greenfield, J. G., Martin, J. P., and Moore, M. T. (1938). *Lancet*, **2**, 1154.
Levin, E. A. (1937). *Arch. intern. Med.*, **59**, 667.
Smith, F. B., and Crawford, J. S. (1930). *J. Path. Bact.*, **33**, 291.

Medical Memoranda

Large Mesenteric Cyst

Mesenteric cysts are held to be the least frequent of intra-abdominal tumours. Warfield (1932) collated 129 cases published since 1920, to which he added two of his own. He estimated that by then about 500 cases had been published since the accidental discovery of a mesenteric cyst during a dissection by the Florentine anatomist Beneveni in 1507. Since the appearance of Warfield's paper another 104 cases have been reported. Most of these are described in the South American and Scandinavian literature, however, while in the British literature only eight cases are recorded—viz., those of Steel (1933), Hastings (1934), Cross (1935), Jackman and Mayston (1936), Jewesbury (1937), Luker (1938), Riggall (1940), and Jones (1943).

CASE REPORT

A boy aged 2½ was brought up to the out-patients clinic for investigation. He had been subject to attacks of abdominal pains, especially after a heavy meal, often culminating in vomiting. These attacks would pass off gradually. The mother, from whom

this history was obtained, had noticed that the child's abdomen frequently became distended during these attacks, and that a gurgling noise was audible towards their end, even though vomiting occurred. These attacks, obviously of intermittent intestinal obstruction, had been observed about every two to five weeks, starting when the child was eight months old, and were stated to have increased in severity.

X-ray examination, undertaken shortly after the beginning of these attacks, had been negative. They were repeated, and now showed considerable dilatation of the mid-coils of the small intestine, with definite delay in the passage of barium. An extrinsic pressure defect was noted on a coil of small intestine on the right side of the abdomen, suggesting compression by a soft tissue mass. Five hours after the barium meal had been given it was still seen to be held up in the small intestine, and none had reached the terminal ileum or caecum. "There is one very persistent coil," the radiologist reported, "closely looped upon itself, which suggests that this portion of the small intestine is actually partly rotated upon its axis and thus producing obstruction." Before this report had been delivered, however, the child had to be admitted for acute small-bowel obstruction.

Laparotomy was performed under general anaesthesia. A large cyst was found in the mesentery of a loop of the lower ileum. It extended for about 10 in. (25 cm.) along the mesenteric border of the ileum from a point approximately 4 in. (10 cm.) proximal to the ileo-caecal junction. The ileum was stretched over it almost to the point of obstruction. Apart from that the mesenteric base of the affected loop was very short and showed marked thickening and scarring, suggesting that volvulus had occurred previously, probably on several occasions. Indeed, twisting of this loop around its axis was noted at the time of operation. As the size of the cyst made it impossible to shell it out without interfering with the blood supply of the overlying gut, the whole loop was removed, a side-to-side anastomosis being made. Many enlarged glands were scattered throughout the mesentery. The child made an uneventful recovery, and was discharged home on the eighth day after operation.

Microscopical examination of the cyst showed the lining to consist of ciliated columnar epithelium with squamous patches. Muscular tissue was present in the wall, and the pathologist was of the opinion that the cyst was probably of teratomatous origin.

COMMENT

Warfield, and after him Luker, divided the cysts according to their origin into the following four groups: (1) Embryocystoma (from Wolffian remnants); (2) cystic dermoid or teratoma; (3) enterocystoma (foetal intestinal diverticulum); (4) chylus cyst.

The cyst described here was in a male child. This is of especial interest, since Warfield states that the cystic dermoids and teratomata always originate from displaced ovarian tissue. According to him no mesenteric cyst of that group has ever been recorded in a male patient, although teratomatous cysts have been met with in the spermatic cord. However, the testicle on its way downwards in the retroperitoneal tissues might occasionally leave some tissue behind, as it were.

I wish to thank Mr. H. H. Stewart, under whose care the child was admitted, and who performed the operation, for permission to publish this case. Also Dr. R. J. C. Campbell, who made the x-ray examination, and Dr. C. J. Young, who supplied the microscopical report.

G. M. HADDA, M.D.,
Late Clinical Assistant at the
Royal Infirmary, Bradford.

REFERENCES

- Cross, A. G. (1935). *Lancet*, **2**, 1409.
Hastings, J. P. (1934). *British Medical Journal*, **1**, 947.
Jackman, W. A., and Mayston, E. A. (1936). *Ibid.*, **2**, 1079.
Jewesbury, R. C. (1937). *Lancet*, **1**, 1170.
Jones, J. A. Seymour (1943). *St Thomas Hosp. Gaz.*, **41**, 7.
Luker, S. G. (1938). *Proc. roy. Soc. Med.*, **31**, 1388.
Riggall, F. and C. (1940). *British Medical Journal*, **1**, 345.
Steel, W. A. (1933). *Brit. J. Surg.*, **21**, 104.
Warfield, J. O. (1932). *Ann. Surg.*, **96**, 329.

Recognizing that chest surgery requires a specially trained staff—medical and ancillary—with special equipment, the Society of Thoracic Surgeons has issued a "Memorandum on the Provision of a National Thoracic Service." It emphasizes that the organization must be regional and that the work must be concentrated in centres. Primary centres should be attached, if possible, to teaching hospitals, and there may be secondary and tertiary centres located conveniently in each Region. The surgery of both pulmonary tuberculosis and other chest diseases should be carried out in the same centre, but the patients should be segregated into separate wards or blocks. In London the special chest hospitals would naturally play a prominent part in the service.

SILICEOUS GRANULOMA DUE TO TALC

W. A. MACKEY AND J. B. GIBSON

FIG. 1.—Section of wall of Meckel's diverticulum ($\times 35$). Dense fibrous tissue has replaced subserous fat. Complex of giant cells surrounds talc particles centrally and at serous surface.

G. HARVEY SMITH

FIG. 1.—Partly crossed "polaroid" screening shows mass of doubly refractive material with cellular reaction under serosal surface of Fallopian tube.

FIG. 2.—Syncytial giant-cell reaction enclosing talc particles ($\times 200$).

FIG. 3.—Same field as Fig. 2 seen through partially crossed "polaroid" screens.

FIG. 2.—Partly crossed "polaroid" screening of granulomatous lesion in wall of Fallopian tube. Note histological similarity to tuberculosis. Two giant cells show doubly refractive masses of talc.

WILLIAM WALKER

FIG. 1.—Adhesion between two loops of small intestine, one obviously hypertrophied. Thin collagenous fusion zone only slightly cellular, but giant cells are present ($\times 8$).

FIG. 2.—Partly crossed "polaroid" screening shows multinucleated giant cell containing doubly refractive material, slightly displaced ($\times 250$).

F. R. MAGAREY AND P. H. DENTON: TORULA
HISTOLYTICA INFECTION OF CENTRAL NERVOUS
SYSTEM

FIG. 1.—*Torula histolytica* in section of meninges ($\times 130$).

FIG. 2.—Torula body in subependymal giant cell ($\times 720$).