OCCUPATIONAL SKIN LESIONS DUE TO PITCH AND TAR

BY

PHILIP ROSS, M.D.

[WITH COLOUR PLATE]

The terms "pitch" and "tar" refer to a multitude of substances of varying composition and complexity, and a brief study of these compounds is desirable for the better appreciation of some of the problems they produce in industry. In addition to various asphalts and bitumens, Abraham (1945) in his recent treatise describes many types of pitch and tar, their origin and mode of preparation, but from our point of view we need consider only those derived from wood and coal.

Wood Tars and Pitches.—These are made by distilling various kinds of wood. Typical examples are the Stockholm, Norwegian, and Russian tars, all of which are used in the tarring of ropes, fishing-nets, boats, etc.

Coal-Tar and Pitch.—The main supply of tar in this country is from the destructive distillation of coal—a process carried out first and foremost in all gas-works, and, secondly, by the big iron and steel manufacturers. In the former instance we have the gas-works tar and in the latter the coke-oven variety. A small amount of blast-furnace tar is formed as a by-product in some of the Scottish iron-works where "splint" coal is used in the extraction of the pure metal from its ore; and an increasing quantity of tar is being made by the low-temperature carbonization of coal.

Chemistry

As the chemical composition of tar and pitch has a direct bearing on the production of skin lesions it is necessary to consider not only the mode of preparation but also some of the concomitant chemical changes. Tar is made by distilling coal, and pitch is the residue left after the distillation of tar.

Distillation of Coal.—At the gas-works coal is placed into retorts and heated to a temperature ranging from 1,200° to 1,400° C. (Martin, 1917), and, as the coal decomposes, ammonia, steam, coal-gas, and crude tar are evolved, leaving an amorphous mass of coke in the retorts. Similar decomposition occurs when coal is treated in the coke ovens, though the final products may vary somewhat from those produced in the retort houses. In both cases, however, the crude tar is collected for further distillation. In the low-temperature carbonization process the coal is distilled at a reading not exceeding 700° C.

Distillation of Crude Tar.—This process is carried out either at the gas-works by-products department or at the special tar distilleries. The results of a typical fractional distillation are indicated in Table I, but it must be stressed that the chemicals which are ultimately formed depend on several factors: (a) the type of coal in the original distillation; (b) the exact method of coal treatment—whether by vertical, inclined, or horizontal retort; (c) the temperature and conditions prevailing in the retort itself; and (d) the

actual technique adopted in the tar distillation. The remaining pitch, which represents about a half of the original quantity of tar, is composed of carbon and numerous polycyclic hydrocarbons.

Under normal working conditions tar is a black, heavy, viscous, and tenacious fluid, while pitch is a black, lustrous semi-solid. When subjected to an increasing temperature they become more and more labile, and tar begins to evolve fumes at about 80° C. The pitches fragment easily, and even the "softer" varieties may be ground to a fine dust when cooled to 10° C.

Industrial Hazards

Because of their unique properties and their cheapness tar and pitch have a widespread application in industry. Hazards from this source occur not only in the gas-works, coke ovens, blast furnaces, and tar distilleries, but also in the manufacture of producer gas, patent fuels, and "briquettes"; in asphalt-workers and road-repairers; in the making of certain electrical equipment such as cables, carbon brushes, carbon electrodes and insulations; in the lens industry, where pitch is used to imbed the lenses during polishing; in the building and allied trades, as in the preparation and the handling of roofing felt, special floor materials, paints, and weather-proofing compounds. Seamen and fishermen, too, handle a fair amount of tar and at times show typical lesions.

Pathology

When we consider the formidable array of coal-tar products, ranging from the light aliphatic and aromatic oils to the complex polycyclic hydrocarbons of pitch, it is not surprising to find that the pathological changes are both varied and bizarre. Our appreciation of these effects is assisted by the knowledge that the heavier oils, tar, and pitch have these characteristics in common: they are all skin irritants; are solvents of lipoids and cholesterol; are auxetic and keratoplastic; and, finally they are carcinogenic. Therefore a skin exposed to these materials may show evidence of epidermal and dermal inflammation; changes associated with an "eczematous" reaction; signs of excessive cell proliferation such as acanthosis, keratosis, and telangiectasis; and perhaps in the later stages malignant degeneration—usually a squamous-celled carcinoma.

An important physical reaction common to many polycyclic hydrocarbons is the property of fluorescence when their extracts are activated by ultra-violet light; and when these solutions are examined spectroscopically one can detect the carcinogens by their typical absorption bands (Mayneord, 1927).

Clinical Effects

Generally speaking, tar and pitch produce similar skin changes, but differences do occur by virtue of their respective physical and chemical properties. On the one hand, tar contains many light, middle, and heavy oils, while pitch is relatively richer in the polycyclic hydrocarbons. Furthermore, during the processing of pitch a fine dust can be generated with a high power of dissemination and penetration—a circumstance which explains the appearance of lesions (a) in remote parts of the body, and (b) in persons who work in the vicinity of, but not actually in, the particular workshop.

Regarding my own findings, I have been unable to differentiate between the effects of each material, as most of our employees are in contact with both pitch and tar either separately or mixed. Only rarely has it been possible to indict only one compound.

Tar Burns.—Owing to more thorough mechanization and enhanced methods of protection of the employees severe

TABLE I.—Distillation of Crude Coal-Tar

Temperature	Compounds Produced
170° C.	Light oils: benzol, toluol, xylols
170–230° C.	Middle oils: phenols, cresols, naphthalene
230–270° C.	Creosote oils, tar oils
270–400° C.	Anthracene oils
	Residue: pitch

Note.—Numerous other compounds are formed in addition to the above: aliphatic hydrocarbons, pyridine, anilines, quinolines, sulphur compounds, etc.

tar burns are uncommon in industry. Should, however, the skin receive splashes of hot tar it must be cleansed either by swabbing gently with trichlorethylene or by the use of a special spray containing this solvent. The burn, usually of the second degree, may then be treated in the routine way. The belief that a tar burn could be followed rapidly by an epithelioma (Bang, 1923) is not substantiated by recent observation.

Tar Erythema.—This reaction is referred to in industry under a variety of names: pitch burn, tar "flashes," and "the smarts." It occurs in those in contact with pitch or tar fumes, and may appear either at the end of the day's work, when the employee emerges into the open, or during the working spell if he is engaged on outdoor duties such as loading or trucking pitch, or tarring roofs. The condition is apparently precipitated by sunlight or a strong wind, and is characterized by severe smarting of the exposed parts of the body, particularly the face and neck. The eyes, too, are often affected, resulting in a burning sensation and photophobia.

Though the symptoms may be sufficiently intolerable to compel the victim to seek refuge indoors, only rarely is he incapacitated for more than a day, and in the majority of cases he is quite able to resume work on the next shift.

In appearance tar erythema resembles ordinary sunburn, and may vary from a slight redness of the skin to a severe dermatitis with oedema. This latter reaction is now uncommon, and is to be seen in those working out of doors in bright sunlight. Of interest, too, is the fact that repeated attacks of this intense form of pitch burn is followed by persistent pigmentation. Conjunctivitis of a varying degree usually accompanies this condition and may necessitate the wearing of dark glasses temporarily.

The causal agent or agents are still unknown, but there appear to be two predominating factors—a sensitization of the skin by some ingredient of the pitch or tar fumes, and the action of bright sunlight on a skin thus rendered susceptible. With regard to the former, the chemicals indicted are thiazine, thiazone, acridine, and other azines present in tar (Leymann, 1930).

To prevent this condition it is necessary not only to apply the general precautionary measures outlined below (under "Prophylaxis") but to use in addition a protective paste or cream which is impervious to tar fumes, pitch dust, and light rays. After experimenting with several applications I have found a kaolin paste based on a formula suggested by Jenkins (personal communication) most efficacious when applied liberally to the exposed parts before each shift (see Appendix).

Treatment of the dermatitis itself should be carried out according to general principles, though I have found that the patients prefer cold cream or similar emollients.

Allergic Eczematous Dermatitis.—In my experience this reaction has occurred most infrequently. During the past twelve years I have carried out periodic and regular inspections of about 170 employees who handle pitch and tar, and my records show that during that time only six cases occurred, four of which were sufficiently severe to warrant a transfer to alternative work. This may appear somewhat surprising when we consider the virulent effects common to many coal-tar derivatives, such as the phenols, cresols, anilines, nitrotoluene, nitrobenzene, etc.

Folliculitis, Acne, Comedones.—These phenomena, though quite common among these workers, rarely cause symptoms or disability. The lesions are, in the main, limited to the forearms, face, and neck, and they pursue a chronic course, waxing and waning at intervals. As with ordinary acne, any severe secondary infection may result in the

formation of small pitted scars, many of which can be seen on the faces and necks of some of our older employees. Of interest, too, is the presence in the scrotum and elsewhere of comedones from which one can express a small plug of sebaceous matter; and Henry (1946) has reported the presence of numerous sebaceous cysts in the scrotum of a man working in a low-temperature carbonization plant. These comedones are no doubt due to the mechanical obstruction of the glands by pitch dust.

In the selection of men for this type of work it is best to avoid engaging those who suffer from acne, as their condition is likely to be adversely affected by the materials they handle.

Pitch and Tar Melanosis.—This form of pigmentation has been recognized in industry for many years. Broadly speaking, there are two main types: (a) a fairly acute form where there is deep tanning of the exposed parts, especially the face, neck, and forearms, and (b) a more chronic variety accompanying other skin changes, such as keratosis and poikiloderma.

The first is a well-known event seen in those who work principally out of doors and who are in contact with pitch or tar fumes. The melanosis usually appears after a few months, is preceded by several bouts of tar "erythema," and persists throughout the summer months, with a tendency to fade during the winter or when the employee is transferred to indoor duties. An interesting concomitant phenomenon is the brown staining of the sclerotics. It is of interest to note also the presence of melanoderma in those in contact with creosote (Prosser White, 1934), anthracene (O'Donovan, 1929), and other coal-tar derivatives. The incidence of this form of pigmentation has been reduced considerably in recent years by the adoption of the various preventive measures directed against the development of pitch "burn."

The second variety of melanoderma evolves much more slowly and, in my experience, affects first the forearms and later the face, the neck, and occasionally the scrotum. After several years' contact with the materials in question one may perceive one of the following abnormalities: first, a brown mottling of the skin not unlike "freckles"; secondly, a diffuse dusky-red pigmentation; and thirdly, a patchy or reticulated deep reddish-brown discoloration. The two latter lesions either herald or accompany "shagreen" skin (Fig. 1).

The factors present in the causation of tar "erythema" appear to operate in the production of melanosis; for, according to Foerster and Schwartz (1939), who have carefully studied this latter condition, the tanning is due to the action of certain light rays on a skin made susceptible by the chemical photosensitizers in tar and pitch.

The mechanism of pigmentation has been investigated by Bloch (1927), who asserts that the epidermal dendritic cells of the basal layer of the skin and of the hair follicles (known as melanoblasts) produce the colourless precursor melanogen, which is transmuted into the pigment melanin by the action either of the ferment oxidase or of ultra-violet light. Some authorities believe that the chromatophores which normally carry pigment from the dendritic cells to the blood vessels may themselves generate melanin when affected by tar.

In contrast to this form of pigmentation is the observation of Dr. R. E. W. Fisher (personal communication) that tar workers frequently show on their forearms pale areas of skin which are the relics of burns caused by splashes of hot tar.

"Shagreen" Skin.—This was the name given by Prosser White (1934) to a pathological state of the skin characterized by keratosis, pigmentation, dilatation of the superficial capillaries, and alteration of its texture. It can be caused by the action of a variety of agents such as excessive sunlight, x rays, and prolonged exposure to oils, pitch, and tar;

and clinically the condition may be regarded as a poikiloderma, several types of which, according to Kinnear (1935), have been described by the following authorities: Jacobi, Petges, Civatte, Riehl, and Hoffmann and Habermann. Only the condition reported by the last two observers had an occupational aetiology. The appearance of "shagreen" skin is unmistakable (Fig. 1), and on palpation the integument feels hard, rough, craggy, and inelastic. So far as pitch and tar are concerned one may perceive the inception of this degeneration after about ten years' close contact, when there appear patches of keratosis and scattered areas of telangiectasis on a background of dusky-red pigmentation affecting the forearms. As the condition advances the lesions become more marked and widespread, and involve also the dorsa of the hands, the face, and the neck, until in due course we have the full-fledged "shagreen" skin with pigmentation, telangiectasis, keratoses, areas of epidermal hypertrophy and atrophy, superficial fissuring, and possibly papillomata. Except for occasional irritation or burning sensation, symptoms are absent; and as the changes are permanent treatment is essentially preventive.

Neoplastic Changes

As has already been indicated, these are manifest as keratoses, papillomata, and epitheliomata.

(a) Keratosis

The earliest macroscopic evidence of the auxetic effects of tar or pitch is perhaps to be seen in the stratum corneum, where after several months of exposure one may detect small, raised flattened areas of keratosis ranging from 1 to 5 mm. in diameter on the dorsal surfaces of the hands, on the forearms, or on the face. These lesions, which closely resemble flat warts, usually remain unaltered for many years. On rare occasions the hypertrophy can be sufficiently prolific to result in the formation of a cutaneous horn.

(b) Papillomata

Papillomatous growths can occur in these workers after one year, though I have seen instances where they have appeared after only six months' contact. Three types have been observed: first, the simple wart; secondly, the typical pitch wart; and thirdly, the soft papilloma. The regions affected are the dorsa of the hands, the forearms, the face, the neck, and the scrotum. Only rarely do we see lesions on other parts of the body.

The *simple wart* resembles the common verruca, and may be sessile or pedunculated. It is innocuous and, as a rule, retrogresses and falls off—the latter process not infrequently being assisted by some minor trauma such as shaving. Should any prove uncomfortable or begin to enlarge rapidly, treatment by low-voltage x rays, by carbon dioxide snow, or by minor surgery should be instituted. Caustics are best avoided in these cases, as their irritant action might stimulate further proliferation.

The *typical pitch wart* appears as a small conical or dome-shaped tumour having a base of apparently normal skin and a keratotic head. The most common site is on the face, particularly near the nostrils (Fig. 2), and on palpation one finds the base hard and infiltrated. The lesion should be regarded as precancerous or frankly malignant, and should be treated either by low-voltage x rays or by local excision.

The *soft papillomata* are to be found in those who have been in contact with these substances for several years, and they appear on the face, especially on the eyelids and adjacent regions. They form small, soft, multiple, pedunculated tumours not unlike skin fibromata, and they are quite innocuous.

(c) Epitheliomata

Every worker who handles pitch or tar and anyone exposed to pitch dust is liable to become a victim of this hazard. As with the production of experimental cancer in mice there is always a variable latent period between the initial contact with these materials and the appearance of the growth—an interval which in my group of cases has extended from 18 months to 34 years. The occasional and casual handling of the compounds is scarcely sufficient to precipitate neoplasms, but five years' exposure to small amounts of pitch dust has resulted in malignant disease of the skin of the face. The cessation of all handling of these carcinogenic substances does not necessarily render the individual immune from further trouble, as growths have appeared after a "lag" period of many years.

Though pre-existing papules, keratoses, and follicles may suddenly become malignant the majority of these neoplasms are cancerous from the start, beginning usually as a typical pitch wart and enlarging to a diameter of 1/4 in. (0.63 cm.) or more within two weeks or so. The most common site for their appearance is on the face (Fig. 2), in the neighbourhood of the nostrils, but they also occur with some frequency on the dorsa of the hands, the forearms, the nape of the neck, the ears (Fig. 5), and the scrotum (Figs. 3 and 4). One patient, a man with an industrial history of 28 years' contact with pitch and tar who had advanced "shagreen" skin, suddenly developed four epitheliomata simultaneously—one on the back of the neck, another on the right ear, and one on each forearm.

If left untreated the tumour enlarges and the keratotic head falls off, leaving the classical epitheliomatous ulcer, circular in shape, with a hard infiltrated base, indurated edge, and raw bleeding surface (Fig. 6). At first the surface is covered with a tenacious slough giving the lesion the appearance of an indolent boil—an error which, in my experience, has occurred on two or three occasions. At times the base advances rapidly, and by a process of eversion converts the ulcer into a mushroom-shaped mass (Fig. 4). The growth is of low malignancy, its spread is slow and localized for quite a considerable time, and glandular involvement is very late. Histological examination reveals a squamous-celled carcinoma, and with early tumours the problem is not so much to find a malignant growth as a cancerous cell.

The best treatment is by the contact low-voltage x-ray apparatus giving about 4,000 to 6,000 r over a period of 8 to 14 days. The epithelioma becomes necrotic, sloughs away, and heals with a thin supple scar which is almost invisible. Early cancers often disappear after a single dose of 1,500 r.

Pathogenesis of Pitch and Tar Cancer

The Exciting Cause.—The association between occupation and epithelioma was first recorded by Pott, who in 1775, in his classical description of chimneysweep's cancer (or soot wart) of the scrotum, suggested that the soot (which contains about 40% of tar) was to blame. It was not, however, till the end of the nineteenth century that the lethal effects of pitch and tar were recognized, and in 1907 the definition "scrotal epithelioma occurring in chimneysweeps and epitheliomatous cancer or ulceration of the skin occurring in the handling or use of pitch, tar, and tarry compounds" was added to the Workmen's Compensation Act (Henry, 1946).

Fresh light was thrown on the subject in 1915 when two Japanese research workers produced malignant skin growths in mice by painting their ears with coal-tar; and in 1924 Kennaway proved that a carcinogenic hydrocarbon was present in a fraction of coal-tar which distilled over a

temperature exceeding 500° C. Carrying this investigation further, Cook, Hewett, and Hieger (1933) were able to isolate from two tons of pitch a few grammes of a powerful cancer-producing agent which on spectrum analysis proved to be 3 : 4 benzpyrene—the now-recognized carcinogen of pitch and tar. In recent years numerous synthetic compounds having a similar action have been prepared—many of them related to the anthracenes (Cook, 1943)—and we now have about 300 such chemicals (Hieger, 1947).

Relative Potency of the Tars and Pitches.—It would appear, then, that the potency of a tar or pitch depends on the benzpyrene content, and any condition favourable to the appearance of this hydrocarbon would render its action more powerful. Such conditions are: (a) the particular type of coal that is mined in Britain and Belgium; (b) the distillation of this mineral at a high temperature; and (c) certain technical factors arising during the treatment both of the coal and of the tar. Generally speaking, gas-works tar and pitch are highly active; blast-works tar is believed to be less noxious, as are the pitch and tar derived from French coal; tar prepared during the low-temperature carbonization process is also relatively harmless, as is the tar obtained from the various woods; coke-oven tar and pitch prepared in America have low lethal effects (Cranch, personal communication); and the bitumens and asphalts, too, have a low carcinogenic action. Some observers, however, regard all types of tar and pitch with suspicion, and in any case from an industrial point of view it is almost impossible at present to obtain in sufficient quantities an inactive tar or pitch. Of the two, pitch is the more conducive to the development of malignant disease.

Ultra-violet Light.—The skin lesions poikiloderma and epithelioma caused by an excess of ultra-violet light resemble those produced by pitch and tar, but what the relationship is between the light rays and the polycyclic hydrocarbons is unknown. It is of interest to note, however, that the epitheliomata appear far more rapidly in rural districts than in urban—for example, in a country factory the latent period for a skin cancer is from 18 months to five years, whereas in London the analogous interval is from 9 to 34 years. It would appear, then, that an excess of ultra-violet light accelerates the action of the carcinogens in some mysterious way.

The "Lag" Period, or Delayed Action.—It has long been recognized that a malignant skin growth can appear several years after the individual has ceased all contact with pitch or tar. The longest interval in my series of cases is seven years, but Henry (1946) reports a "lag" period of 51 years in a patent-fuel worker. In what way the cutaneous cells have been "primed" is another puzzle to be solved.

Susceptibility.—There is no doubt that some workers are highly susceptible to the development of neoplasms of the skin, and this idiosyncrasy does not depend on the colouring alone, as blonds and brunettes are equally liable. Moreover, some employees who have well-marked shagreen skin have apparently never suffered from typical pitch warts, and there are others who have had numerous warts but never an epithelioma. On the other hand, an epithelioma was the first skin lesion (apart from attacks of tar erythema) to appear in seven of my cases. During the past 12 years, from a personnel averaging 170, including 10 women, I have noted skin growths in 102 individuals, and of these 16 showed malignant tumours. Among the women employees who come into contact with these compounds I have observed a corresponding ratio of papillomata but no incidence of malignancy. It is understandable why the exposed parts are affected, but why the scrotum too is involved is not easily explained. Contamination alone is

not the answer, as I have never seen an industrial epithelioma of the penis or of the adjacent parts of the thigh.

The Increasing Incidence of Occupational Cancer

Since 1920, when epitheliomatous ulceration due to pitch and tar became an industrial disease notifiable to the Chief Inspector of Factories, there has been a steady increase in the number of cases notified. This is clearly shown in Tables II and III, taken from the Annual Report of the Chief Inspector of Factories for the years 1945 and 1946.

TABLE II.—Notified Cases of Epitheliomatous Ulceration in Tar Distilling and Gas-works, 1921–45 (in Five-Yearly Periods)

Period	Tar Distilling	Gas-works	Total
1921–5	67 (9)	22 (6)	89 (15)
1926–30	132 (12)	85 (43)	217 (55)
1931–5	178 (11)	88 (51)	266 (62)
1936–40	299 (10)	59 (19)	358 (29)
1941–5	358 (5)	91 (4)	449 (9)

Note.—The principal figures relate to cases, the figures in parentheses to deaths.

TABLE III.—Extract from Table I of the Annual Report of the Chief Inspector of Factories for 1946

Industrial Diseases							
Epitheliomatous Ulceration	1920	1930	1940	1943	1944	1945	1946
1. Pitch	32	44 (1)	85 (1)	54	99 (1)	83	98 (3)
2. Tar	10 (1)	53 (9)	34 (3)	57 (1)	61 (2)	93 (1)	103 (4)

It is nevertheless encouraging to observe the decline during the past ten years in the number of deaths recorded, despite the somewhat alarming increase in the numbers notified.

What is the reason for the apparent increase in the incidence of industrial skin cancer? Several factors are involved: first, following the efforts of the Home Office Inspectorate, both the medical world and factory managements have been made aware of this particular hazard, and skin epitheliomata which were formerly thought to be a natural visitation are now being considered correctly in the light of occupational environment. Secondly, many workers have now passed the "latent" period and their skins are liable to develop this lesion. Thirdly, because of the increasing interest displayed by surgeons, pathologists, and industrial medical officers, many more biopsies are being performed, and tumours which were at one time thought to be innocent are now found to be malignant.

Prophylaxis

The widespread use of pitch and tar and the dangers attendant on their use render the question of prevention of the utmost importance in industry. This problem can be attacked in several ways.

1. Substitution

As gas-works tar and pitch are carcinogenic, attempts should be made to substitute other less noxious compounds—as, for example, wood tar, bitumen, tar from the low-temperature carbonization process, and fractions from petroleum. Unfortunately the application of these alternative products is not always practicable either because of their unsuitability or because of their prohibitive cost.

2. Protection of the Worker

One of the requirements of the Factories Act is that all injurious fumes and dust should be removed by mechanical means; and the strict observance of this regulation would go far in the prevention of skin troubles. Where practicable all the processes should be mechanized to avoid undue handling of the materials; machines should be enclosed, and efficient exhausts should be fitted to carry away any dust or fumes that are generated.

Employees should be made aware of the hazards involved, and be trained to work cleanly and to prevent the dissemination of effluvia. Also they should be encouraged to report early any abnormal skin condition or appearance.

The use of protective clothing, too, is of paramount importance. Overalls made of stout drill material or denim should fit snugly at the neck, wrists, and ankles, and they should be changed and cleaned daily. I would recommend also a daily change of underclothing; but this, unfortunately, is not feasible while clothes rationing remains so severe. The use of "any old suit" which rapidly becomes soiled and impregnated with pitch and tar is to be condemned.

Gauntlet gloves made of stout rubber or leather, when used judiciously in certain processes, have a definite protective value, as have suitable goggles in shielding the eyes.

Employees should be encouraged to maintain a high standard of body cleanliness, and time and facilities should be provided so as to ensure a wash and shower bath after every shift. The application of "barrier creams" and the plentiful supply of warm soft water should obviate the use of special cleansers—either chemical or abrasive—which of themselves may be a fruitful source of dermatitis. The restricted supply of soap is causing concern and the use of substitutes is not always desirable, though "teepol" or similar "wetting" agents may help to eke out the soap ration.

In some firms it has been the custom among those working with pitch, tar, and heavy oils to use soda ash for washing purposes, and some of the older employees are unequivocal in their assertion that the use of this alkali prevented the development of warts. The use of caustic soda has also been referred to by O'Donovan (1929) in the popular treatment of skin growths by men working with tar. Though weak alkali solutions may be beneficial, the use of them cannot be recommended, because of the serious risk of dermatitis.

I am of the opinion that barrier creams fulfil a useful function, and at our works several types are available, depending not only on the materials handled but on personal idiosyncrasy. In addition to the usual properties of a protective cream, the compound should be opaque to ultra-violet light. Several chemical and cosmetic firms now specialize in these preparations and are prepared to make suitable applications, especially if they are supplied with samples of the materials used and provided they are given information about the conditions under which these are handled—for example, the temperature of the process and the use of ancillary substances and solvents. The creams should be applied liberally to the hands, forearms, face, and neck before each working shift.

In a recent investigation by Kennaway (personal communication) it was shown that the use of a protective cream on the skin of a mouse delayed considerably the appearance of experimental tar growths.

3. Selection

While it is not yet possible to foretell those likely to develop skin lesions I nevertheless examine all prospective candidates for this type of work, and I exclude those with existing skin disease or with a bad dermatological history. The question of skin colouring is in my opinion not very helpful.

4. Inspection

I regard the frequent and regular inspection of employees an important step in prophylaxis. Not only am I able to detect early skin changes of which the worker himself may be unaware, but I am able also to educate and reassure him,

and discuss with him the various precautionary measures that are being adopted. I see almost every pitch and tar worker at three-monthly intervals, and those who have or have had skin lesions are examined as often as their condition warrants. Only in this way can we be sure of instituting early treatment and prevent serious complications.

5. Propaganda

Most industrial concerns are loath to broadcast indiscriminately the hazards obtaining in the use of these compounds, and do not favour the distribution of leaflets or the posting of warning notices (except the one recommended by the tar distillers), as these measures are likely to deter men from working in these processes. Instead, individual education and training are left to the medical officer and to the foremen and charge hands. In this way adequate welfare supervision is ensured without creating unnecessary alarm.

Statistical Surveys

As medical officer to a large engineering firm who manufacture various types of carbon products, I have had during the past 12 years the opportunity of observing some 170 employees who have handled pitch and tar more or less consistently during that period. My investigation has been facilitated by the small labour turnover, for at the end of 1947 there were 87 workers with at least 12 years' service to their credit and 53 with 20 years' or more.

An analysis of the incidence of their lesions is shown in the following tables. In Table IV I have recorded the number of personnel who have shown the appropriate condition for the first time—subsequent recurrences being ignored; and in Table V are noted the latent periods of the various manifestations. Tables VI and VII show the sites of election for papillomata and epitheliomata.

TABLE IV.—Incidence of Lesions Among a Personnel (including 10 Women) Averaging 170 from 1936 to 1947, Arranged in Four 3-Yearly Periods. Recurrences Not Included.

Lesions	1936-8	1939-41	1942-4	1945-7	Total
Allergic dermatitis	2	2	2	0	6
Folliculitis, acne, comedones ..	15	14	21	16	66
Melanosis	10	7	5	3	25
Shagreen skin	17	6	2	1	26
Keratosis	9	8	11	3	31
Papilloma	18	24	40	20	102
Epithelioma	2	2	8	4	16

The following points are worthy of note:

1. *1936 to 1938.*—I began my investigations during these years, and while collating the data I encountered conditions which had been in existence for several years. This was particularly the case with the 17 instances of shagreen skin, which had developed several years before 1936.

2. *1942 to 1944.*—The increased incidence of folliculitis, keratosis, papillomata, and epitheliomata during this period can be accounted for in two ways: first, the advent of war necessitated an augmented production of accessories derived from pitch and tar, and from 1940 onwards not only were greater amounts of these materials being handled but working hours were extended. Secondly, the establishment of a dispersal factory in a rural area brought a further 20 men into contact with these compounds from the years 1941 and 1942. By taking into account the latent periods one can appreciate the rise in the number of lesions recorded during these years.

3. *1945 to 1947.*—I attribute the reduction in skin troubles in this interval to the intensification of the campaign of prophylaxis, especially in the provision of clean overalls daily, in the more widespread use of barrier creams, in the supply of soft water for washing, and in the allowance of ample washing-time. We are indebted to the medical inspectors of factories for their assistance in the clean-overall service and to a firm of cosmetic manufacturers for their research work on suitable creams.

4. *Papillomata.*—Of the 102 cases 62 showed multiple warts.

5. *Epitheliomata*.—16 men developed malignant skin tumours (all confirmed by biopsy), and of these 9 had previously suffered from simple warts and 3 had multiple epitheliomata. There were no fatal cases. The 9 with simple warts have also been included under "papillomata."

TABLE V.—*Latent Periods*

Folliculitis, acne, comedones		1 month or more
Melanosis		6 months to 5 years
Shagreen skin		Over 10 years
Keratosis		6 months or more
Papilloma		6 months to 43 years
Epithelioma	{ Rural district	18 months to 5 years
	{ Urban district	9 to 34 years

Note.—The shorter latent period for the appearance of a skin carcinoma in the rural district is discussed under "Ultra-violet Light." Because of this latent period all employees who leave the firm are requested (where practicable) to report to us any rash, pimples, or warts that might appear at some future date.

TABLE VI.—*Site Incidence: Papillomata*

Employees affected	..	102 (including 6 women. Age from 29 to 66)
Single lesions	..	40
Multiple lesions	..	62
Site affected:		
Nose and lips	..	43
Cheeks, eyelids, chin	..	24
Ears	..	6
Neck	..	8
Scalp	..	1
Hands	..	29
Forearms	..	16
Scrotum	..	11
Leg	..	1
Anus	..	1
Total	..	140

It would appear that the site of election for these growths is the face (48%); the incidence on the hands and forearms is 32%, and on the scrotum 8%.

TABLE VII.—*Site Incidence: Epitheliomata*

Employees affected	..	16 (no women. Age from 34 to 62)
Single lesions	..	13
Multiple lesions	..	3
Site affected:		
Cheeks	..	5
Nostrils, lips	..	4
Ears	..	3
Neck	..	2
Scalp	..	1
Forearms	..	3
Arm	..	1
Scrotum	..	4
Total	..	23

The most frequent site for the appearance of epithelioma is on the head and neck regions (65%). The upper limbs and the scrotum each show a rate of 17%.

Summary

A brief account has been given of the preparation, the chemistry, and the industrial uses of coal-tar and pitch.

The following clinical effects have been described and discussed: tar burns; tar erythema; allergic eczematous dermatitis; folliculitis, acne, and comedones; melanosis; shagreen skin; neoplastic changes, and epitheliomata.

The pathogenesis of pitch and tar cancer has been discussed.

Various prophylactic measures have been described.

A statistical survey has been included.

APPENDIX

Formula for Starch and Kaolin Paste for the Prevention of Tar Erythema

Starch		28 g.
Kaolin		1,600 g.
Phenol		36 g.
Glycerin		50 ml.
Distilled water		2 litres

The starch is made into a thin paste with about 50 ml. of water and poured into 950 ml. of boiling water. The boiling is continued for a few minutes to ensure the breaking up of the starch cells. The phenol and glycerin are added to the mucilage, and finally the kaolin, in small quantities at a time, with constant stirring. The final mixing is best done with the hands to make certain that no lumps remain.

The phenol prevents the development of mould in the paste and the glycerin renders the paste more "supple" and prevents rapid drying and cracking. Colouring matter such as cochineal may be added for a better cosmetic effect.

I wish to thank the Controller of H.M. Stationery Office for permission to reproduce the tables from the Annual Reports of the Chief Inspector of Factories, and Professor Kennaway for permission to publish the results of his experiment with a barrier cream in the production of experimental skin cancer (1946). I also wish to express my indebtedness to Mr. A. E. Wiggs, B.Sc., Dr. Henry MacCormac, Dr. Davidson, late H.M. Inspector of Factories, and Dr. L. B. Bourne for their kind assistance and suggestions.

REFERENCES

- Abraham, H. (1945). *Asphalts and Allied Substances*, 5th ed., p. 67. Van Nostrand Co., New York.
- Annual Report of the Chief Inspector of Factories, 1945, p. 59; 1946, p. 49.
- Bang, F. (1923). Quoted by White (1934).
- Bloch, B. (1927). Quoted by Foerster and Schwartz (1939).
- Cook, J. W. (1943). "Chemistry and Cancer." Lecture to the Royal Institute of Chemistry.
- Hewett, C. L., and Hieger, I. (1933). *J. chem. Soc.*, p. 395.
- Foerster, H. R., and Schwartz, L. (1939). *Arch. Derm. Syph., Chicago*, 39, 955.
- Henry, S. A. (1946). *Cancer of the Scrotum in Relation to Occupation*. Oxford Univ. Press.
- Hieger, I. (1947). *Brit. J. Radiol.*, 20, 145.
- Kennaway, E. L. (1924). *J. industr. Hyg.*, 5, 462.
- Kinnear, J. (1935). *Brit. J. Derm. Syph.*, 47, 191.
- Leymann, H. (1930). *Occupation and Health*, 1, 467. Int. Labour Organ., Geneva.
- Martin, G. (1917). *Industrial and Manufacturing Chemistry*, 1, 392. Crosby Lockwood.
- Mayneord, W. V. (1927). Quoted by Heiger (1947).
- O'Donovan, W. J. (1929). *Arch. Derm. Syph., Chicago*, 19, 595.
- White, R. Prosser (1934). *The Dermatogoses or Occupational Affections of the Skin*, 4th ed. Lewis, London.

OBSERVATIONS ON TUBERCULOUS MENINGITIS IN CHILDHOOD*

WITH SPECIAL REFERENCE TO EARLY DIAGNOSIS

BY

W. S. CRAIG, M.D., F.R.C.P.Ed., M.R.C.P.

Professor of Paediatrics and Child Health, University of Leeds

Of all the clinical pictures met with in children there is probably none which has sustained such intense interest over so many years as that of tuberculous meningitis. The first account of the clinical course of the condition was given over 180 years ago by Robert Whytt during the time that he occupied the chair of medicine at Edinburgh University. His description, under the title "Observations on the Dropsy in the Brain," is still recognized as a classic.

Interest in the subject of tuberculosis in childhood, and of tuberculous meningitis in particular, has received added stimulus from the trials being given to streptomycin therapy. If this new form of therapy is to receive reasonable trial patients submitted to treatment must be diagnosed early. This constitutes something of a challenge to clinicians, because it must be admitted that we have not advanced greatly from the days of Whytt in the early diagnosis of tuberculous meningitis, no matter how improved are our methods of investigation in relation to the later stages of the condition.

Successive Clinical Phases

It is customary to describe the clinical course of tuberculous meningitis in terms of three stages. Sir Frederick Still used to say that while no fool could err when the middle and last stage had been reached, the wisest physician might fail to recognize that anything serious was amiss with a child in the first and early stage. It is this early

*Summary of an address given to the Harrogate Medical Society on Dec. 11, 1947.

P. ROSS: OCCUPATIONAL SKIN LESIONS DUE TO PITCH AND TAR

Fig. 1.—Shagreen skin of forearm. A. B., aged 52. Contact for 26 years. A "close-up" has been taken of the affected part to show the details of the skin degeneration.

Fig. 2.—Typical pitch wart on upper lip. C. D., aged 50. Contact for 4 years. Dome-shaped tumour with keratotic head proved innocent at biopsy. Two tiny warts near main lesion.

Fig. 3.—Pitch wart of the scrotum. E. F., aged 41. Contact with pitch and tar for 16 years. Tumour was innocent and disappeared with x-ray treatment.

Fig. 4.—Epithelioma of scrotum. G. H., aged 42. Contact with pitch and tar for 17 years. Mushroom-shaped lesion had been growing for one month and had been mistaken for indolent boil.

Fig. 5.—Epithelioma of pinna. I. J., aged 39. Contact with pitch and tar for 20 years. He had had the lesion for one month and it was beginning to enlarge rapidly. Note similarity to papilloma in Fig. 2.

Fig. 6.—Epithelioma of upper arm. K. L., aged 49. Contact with pitch and tar for 34 years. Lesion for 5 weeks; treated as indolent boil. Note typical epithelioma. Slight break in wall of ulcer marks earlier biopsy. No glands. Healing perfect after x-ray treatment.