his career in the interests of humanity." Unfortunately, your correspondent has completely missed the point of the story, which he appears to think was told against the "bad old days." The opposite is the case. The story is told by a physician who fears that the new Service will threaten the independence of medical opinion and action. He tells of this clash with the medical superintendent of his hospital, pointing out that his refusal to comply with the medical superintendent's advice did not involve any risk for him, since he was in no way dependent on the medical superintendent for his livelihood. He expresses a fear that similar independence might be costly in future.

It is to be regretted that your correspondent did not take the precaution of checking his facts before making this venture into radio criticism.—I am, etc.,

Broadcasting House,
London, W.1.

NESTA PAIN.

Excretion of Antimony

SIR.—With reference to your annotation (July 10, p. 83) on this subject, pharmacologists elsewhere have confirmed the records¹ made in Egypt in 1931 and subsequently showing that half the amount of antimony injected may be recovered from the excreta within a month. This would account for the re-appearance of ova in the excreta of patients undergoing the treatment with massive doses here in Natal with the sodium equivalent of tartar emetic as recommended in Southern Rhodesia. It is unreasonable to rely on the mere absence of ova from the excreta as evidence of the destruction of the adult worms until they have had time to recover from the effects of the drug or have been demonstrated by post-mortem examination.

I was asked to examine a tobacco planter whose skin test had given a positive reaction for schistosomiasis in Southern Rhodesia. As he showed no signs or symptoms of infection and his eosinophils were only 2.5% I did not consider that any treatment was indicated. He stated that his natives refused to submit to the treatment adopted for a rapid cure since several of their colleagues had died from its effects.—I am, etc.,

Durban, South Africa.

F. GORDON CAWSTON.

REFERENCE

- ¹ Khalil, M., *Arch. Schiff- u. Tropenhyg.*, 1931, **35**, 106.

Diabetic Coma

SIR.—I have read with great interest Professor R. H. Micks's article on diabetic coma (July 24, p. 200) and Drs. R. D. Lawrence and Wilfrid Oakley's letter on the same subject (Aug. 7, p. 310), and I should like to add one or two points to the discussion. Treatment of diabetic coma must be directed towards remedying not only the metabolic derangements but also the concomitant biochemical changes and associated disturbances which occasionally culminate in sudden death.

Dr. Lawrence and Dr. Oakley rightly stress the need for careful administration of insulin in the average case of pre-coma, whereas the case of profound coma requires heroic treatment with very large and repeated doses of the quickly acting soluble insulin. It must always be remembered that circulatory collapse is the most dangerous feature of diabetic coma, and the patient must be treated for shock—the maintenance of blood pressure being as important as in any other condition of shock. Fluid and food by mouth should always be withheld, since they are dangerous at this stage. The patient must be kept warm to prevent further heat loss, and aspiration of the gastric contents to prevent dilatation of the stomach with subsequent possible inhalation asphyxia—a definite cause of death in diabetic coma—should never be omitted. Professor R. H. Micks draws attention to the importance of gastric aspiration in the management of diabetic coma.

An intravenous saline drip, where necessary, to replace fluid loss, thereby correcting dehydration and the blood chemistry, is a very necessary part of the treatment of all cases of diabetic coma. Professor Micks states that Root's statistics strongly support his claim that normal saline is the only intravenous injection needed in diabetic coma and that other authors have asserted that alkalinizing solutions are necessary or that plasma should be used. Of the American authors, Root and Joslin do not advocate alkalinizing solutions, while Russell M. Wilder, of the Mayo Clinic, recommends the administration of alkali if the carbon dioxide combining power of the plasma is less than 20 volumes per 100 ml. I believe that the decision to give or withhold alkalis depends on simple basic principles.

The indication for the administration of alkali is marked hyperpnoea, and if this symptom is present the administration

of alkali can be a life-saving measure. In the average case of diabetic coma or pre-coma there is not the distressing hyperpnoea which is a feature of some cases of profound coma. Where there is marked hyperpnoea, however, administration of alkali—25 g. of sodium bicarbonate in 5% solution intravenously—should always be administered, as death from respiratory paralysis occurs in uncorrected severe diabetic acidosis through over-stimulation of the respiratory centre. It is true that insulin in the presence of adequate glycogen will correct the acidosis and therefore the hyperpnoea, but this is a comparatively slow process. Intravenous alkali, on the other hand, although not preventing acidosis will rapidly remove the dangerous symptom of hyperpnoea. The importance of aspiration asphyxia and respiratory failure due to marked hyperpnoea as causes of sudden death in diabetic coma have not received the prominence they merit and cannot be over-stressed.

Although there is a difference of opinion as to whether glucose should be given during the treatment of diabetic coma, it is certainly wise to give glucose in those cases receiving large amounts of insulin, particularly after the first 6–12 hours; 1–2 g. of glucose for each unit of insulin injected will guard against hypoglycaemia, which is a constant danger when repeated large doses of insulin are administered. Therefore, if it is not practicable to perform frequent estimations of the blood-sugar level, glucose should always be given in order to guard against hypoglycaemia, except in the earliest stages of treatment when the blood-sugar level is always high.—I am, etc.,

Maidstone, Kent.

GEORGE R. W. N. LUNTZ.

Peritoneoscopy

SIR.—I read with great interest the paper by Mr. John Hosford on peritoneoscopy (Aug. 14, p. 348), but before the term becomes part and parcel of our medical vocabulary I should like to follow Rendle Short's example and offer "coelioscopy" as an alternative. "Peritoneoscopy," in addition to being clumsy and cacophonous, would mean an instrument for viewing the peritoneal membrane and not necessarily the abdominal viscera. The other term suggested in the article, "laparoscopy," being derived from *λαπάρα*, the flank, is also a misnomer, as is "laparotomy"; the only true laparotomy is a kidney incision.

On the other hand, "coelioscopy" (or perhaps more correctly "coeliascopy")—a far neater word than the others—means the examination of the abdominal contents (*κοιλία*), which is the true purpose of the operation. In general, I feel that more care should be taken with our vocabulary. Had this occurred in the past, we should have been spared many glaring errors. I suppose it would be pedantic to quarrel with "appendicec-tomy," but surely "hypophrenic" or "subdiaphragmatic" are preferable to the hybrid "subphrenic," and "proctocoele" to "rectocoele"?—I am, etc.,

Guildford.

A. BARNSELY.

Choosing a Career

SIR.—It seems to me that in recent months we have devoted a great deal of attention to monetary considerations resulting from the passage of the Health Service Act. The Spens Report has still further accentuated the financial aspects of medical work and appears to have justified the demand for raising the standards of direct payment in all clinical posts. From the university point of view this has been followed by increases in the salaries of professors and lecturers in all faculties—medical and non-medical alike. The young graduate on the road to specialism has perhaps come best out of the deal, and if the younger generation of teachers in the preliminary subjects has been overlooked it broaches a question which demands an immediate answer.

In my view, however, elements other than those of remuneration should have the attention of the potential doctor after he has, from about the time of his third year onwards, surveyed the field of medical service. His object should be that of selecting from this field the type of work which would allow him on retiring to look back with satisfaction on his life's journey. They are, indeed, fortunate who can do this. In our profession at the present time there is a wide range of choice open to candidates. Given a certain standard of ability, coupled with