there is pain in the anterior aspect of the thorax, and strangely enough a nitroglycerin tablet relieves the pain in, say, 30 seconds.—I am, etc.,

London, W.5.

R. HALSTEAD DIXON.

Use of Ring Pessaries

SIR,—It is always a pleasure to read a broadside from Mr. Mortimer Reddington (July 24, p. 227), for although some of the shots may be wide, many undoubtedly score. It is certainly true that in the past far too many women have had ring pessaries fitted. When I was chief assistant of the department at St. Bartholomew's Hospital it was the custom for these patients to come up one afternoon a week to have their rings changed by students who were undergoing their gynaecological training. They came in great numbers, but I suppose it was because in those days beds were few and the operation was not as safe as it is nowadays.

The ring pessary has, however, in my opinion certain definite uses. It can quite reasonably be advocated in the case of the young woman with descent of the vaginal walls immediately following childbirth where physiotherapeutic measures fail, but it is not fitted to support the tissues till involution has occurred. It is fitted to support the tissues until the baby is weaned and the mother can be operated upon. I think it is also of value to patients who have symptoms but no signs of prolapse. It should be used in those rare cases unfit for operation, and in those rarer cases where the patient refuses operation.

Now a word about the Hodge pessary condemned to be "put away altogether, and for ever." No splint will correct a deformed limb. The deformity must be corrected and the splint applied. If the retroversion is first corrected, then the Hodge will keep it so, provided it is long enough to reach from the posterior fornix to the vulva. I personally regard the Hodge pessary as invaluable in elucidating the cause of backache. I would never contemplate shortening the round ligaments for backache unless either the uterus could not be replaced or the patient's backache not dispersed by her wearing a ring for a short time.—I am, etc.,

Blackheath.

KEITH VARTAN.

Foetus Papyraceus

SIR,—The case of twin pregnancy with acute hydramnios treated by abdominal paracentesis (July 24, p. 205) is most interesting, and Dr. R. W. Danziger is to be congratulated on the successful result. It is a pity, however, that more details were not furnished about the "blighted" twin, which is simply described as a foetus papyraceus. The interest lies in the aetiology of this condition, which usually follows mechanical compression of the foetus after loss of its liquor amnii; the compression may be effected by the fellow twin itself, or more usually by its bag of waters, but in order to produce a foetus papyraceus the compressing force must be adequate to cause collapse of all body cavities, so that maceration may be arrested. Thus the development of a rigid thoracic cage and firm cranial vault would prevent this pathological sequence. An analogy may be drawn between this phenomenon and the preservation of botanical specimens by pressure between sheets of blotting paper; the process of vegetable tissue autolysis cannot be prevented when the structures concerned are too bulky to be dehydrated by compression.

It is generally considered that the development of acute and subacute hydramnios in association with uniovular twinning is due to uneven foetal demands from the common placenta; the interplacental vascular anastomoses ("third circulation") allow the more powerful circulation to withdraw an excess of fluid into one foetus, resulting in some way in hydramnios for the larger and dehydration of the smaller twin. In most cases some sort of fluid balance is achieved, but rarely the smaller foetus may perish from fluid depletion, producing a dried, wrinkled specimen which does not undergo maceration and is not very much compressed. This is the mummified foetus, and many authorities fail to distinguish between it and the papyraceus.

In favour of a diagnosis of mummification in the case described is the fact that radiography some weeks after foetal death showed only failure of growth, with no unequivocal signs of compression. In addition, the foetal size was considerably smaller than the period of gestation should produce, suggesting

impairment of growth before foetal death. Additional evidence would be furnished by the presence of fluid in the affected amniotic sac, which is naturally absent with a foetus papyraceus, and complete infarction of that portion of the placenta supplying the morbid twin.—I am, etc.,

Birmingham.

W. G. MILLS.

Production of Rh Agglutinins

SIR,—In the article by Drs. George Discombe and H. O. Hughes (Aug. 14, p. 329) on the importance of transfusion as a cause of haemolytic disease, the paper by myself and others (*British Medical Journal*, 1945, 1, 584) is quoted as having expressed the view that patients afflicted with certain diseases might be especially apt to produce antibodies to antigens which they themselves lacked. No such point of view was expressed in our article. The article concerned itself simply with a report of facts—namely, the truly colossal Rh agglutinin titre produced in response to multiple transfusions of Rh-positive blood, the comparative absence of symptoms in the incompatible transfusions, the technical pitfalls in the typing and compatibility tests, the ultimate outcome of the case, and a brief factual report on the necropsy findings. No opinion was expressed as to the relationship of the ready production of the Rh agglutinins to the patient's disease, nor was it even considered. I have personally never held or expressed the view that patients suffering from certain ailments might be especially apt to produce antibodies to antigens, especially the blood group antigens, which they themselves lacked.—I am, etc.,

Cardiff.

R. J. DRUMMOND.

Independent Doctors

SIR,—Your annotation on "Independent Doctors" (Aug. 14, p. 347) concludes pathetically by saying, "There would appear to be no reason why such a committee [separate organization for independent doctors] should not be set up in the B.M.A." The B.M.A. has, apparently, as short a memory as Mr. Bevan. No doctor could avoid being shocked by the knavish way in which the B.M.A. deserted the sound stand it had taken (as a result of the strongly expressed views of the medical profession) when the appointed day drew near.

What is required by all doctors, and especially independent doctors, is an active body rather than a talking body such as the B.M.A. If Lord Horder can supply this dire need, good luck to him.—I am, etc.,

Bretby Hall, Burton-on-Trent.

R. LUNT.

Surgical Anatomy of the Parotid Gland

SIR,—I was very surprised to see Mr. Hamilton Bailey's article on this subject (July 31, p. 245) and to find that he persists in writing in the same vein as in a similar article in the *Journal* of March 29, 1947 (p. 404).

When I read the latter-mentioned article I carried out a series of dissections on the parotid region after injecting indian ink along the main parotid duct into the gland. To identify the small ducts within the gland I had to use the dissecting microscope. My findings with regard to the relation of gland to the facial nerve were entirely at variance with those of Mr. Hamilton Bailey. I could not display a bilobed arrangement of the parotid with the facial nerve running between two lobes. To illustrate the relationship briefly, let me compare the facial nerve plexus to a trellis-work fence, with the parotid gland, like a creeper, growing from the direction of the buccal cavity and winding itself indiscriminately around the nerve and its branches. I communicated my findings to Mr. Hamilton Bailey, previous to my reading a paper on the subject to a meeting of the Anatomical Society of Great Britain and Ireland in London in November, 1947.

I should like to comment on two other points in Mr. Hamilton Bailey's article. First, the size of the main parotid duct—surely it is larger than a fine hypodermic needle? During my investigation a year ago I could easily insert a hypodermic needle of a substantial calibre into the duct. Because I have not the specimens to hand I cannot say more than that the duct is definitely larger in diameter than that of a fine hypodermic needle; and that observation is in formalin-hardened subjects where the tissues normally tend to shrink under the influence of