which yielded another pint (570 ml.) of yellow pus. On this latter finding the cavity was drained after the instillation of 100,000 units of penicillin.

Convalescence following operation was uneventful. The pus from the abscess was sterile and yielded no amoebae. Repeated examinations of the stools failed to demonstrate the entamoeba, either in the cyst or vegetative form. X-ray examination of the bowel showed no pathological change. Sigmoidoscopy at this stage revealed the smooth, pale, glassy and rather rigid mucous membrane associated with chronic amoebiasis. A full anti-amoebic course was therefore given, the response being immediate and dramatic.

This patient was carefully questioned in order to elicit a possible source of infection. He lives in a Surrey suburb. Neither he nor any of his family has ever been out of the country, nor have any of his friends or business associates. He had no illnesses or symptoms until the day of admission to hospital.

C. F. CRITCHLEY, M.S., F.R.C.S.,
Honorary Surgeon,
Sutton and Cheam General Hospital.

Cure of Exomphalos in a 2-hours-old Infant

The following case is worth recording, as the patient, so far as we know, was the youngest to be anaesthetized successfully with cyclopropane; it also shows that children do stand a fair amount of handling of the viscera.

CASE REPORT

A full-term female infant was admitted to Dulwich Hospital on Dec. 7, 1947, with an exomphalos. On examination the abdominal

FIG. 1.—Showing the exomphalos before operation.

viscera were lying on the abdominal wall covered by transparent membrane (Fig. 1). The striking feature was that the major part of the liver was lying outside the abdomen.

Operation.—The child was operated on two hours after birth under cyclopropane-and-oxygen anaesthesia administered by Dr. J. Piney. An incision was made along the edge of the exomphalos and the membrane excised. The liver and then the intestines were returned to the abdominal cavity with some difficulty and the wound was closed with through-and-through silkworm-gut sutures. In

FIG. 2.—Showing the condition three weeks after operation.

spite of the considerable handling of the viscera the child made an uneventful recovery. Fig. 2 shows the condition three weeks later. At the time of writing the child is 6 weeks old and is progressing satisfactorily.

We wish to thank Dr. J. Piney for his excellent anaesthetic, Dr. O. W. Roberts for permission to publish this case, and the nursing staff for their post-operative care.

B. S. S. ACHARYA, M.B., F.R.C.S.
R. W. BURSLEM, M.D., M.R.C.O.G.

A Case of Rupture of Muscles of the Forearm

Rupture of muscles and tendons without breaking the skin is not uncommon. In the case of the long head of the biceps brachii, the plantaris, and the rectus femoris it presents well-recognized clinical features. I had not previously seen a case of rupture of muscles of the forearm, nor have I found it described in a somewhat limited search of the literature. Mercer (1943) does not mention any of the forearm muscles in a list of "those most frequently affected." The following case is reported because of its rarity, the clinical features seen, and the satisfactory result after a severe injury.

CASE HISTORY

A private soldier aged 21 was driving a 15-cwt. truck when it went over a milestone and overturned, late on the night of June 19, 1947. He is unable to tell how his arm was injured because "everything happened so quickly," but as the milestone was less than 1 ft. (30 cm.) high it is likely that the wheel was very powerfully jerked in his hand. When seen in hospital about 14 hours after the injury, he complained of "pins-and-needles" in the fingers of his left hand.

His general condition was good. The left arm was very swollen from a few inches above the elbow to the finger-tips. The skin was pink and flushed. There was only a flicker of movement in the fingers and thumb. The radial pulse was present at the wrist. Radiographs showed no bony injury. A provisional diagnosis of subfascial haematoma was made; the limb was elevated on pillows and efforts were made to do finger movements. Next day (June 21) the limb was rather more swollen, the paraesthesia was unaltered. The radial pulse was still easily felt. Exploration was considered advisable.

Operation.—At noon on June 21—i.e., about 36 hours after the injury—a longitudinal incision was made over the maximum swelling, on the antero-medial aspect of the forearm, from about 1½ in. (3.75 cm.) above the elbow to 4 in. (10 cm.) below it. The skin gaped immediately, revealing rather congested superficial veins and deep fascia which was tense over a haematoma. The fascia was incised in the same line, and it was then seen that the large blood clot lay among ruptured pieces of muscles. The clots were removed, and it was found that the palmaris longus lay intact over the tattered and separated ends of the flexor digitorum sublimis and the flexor carpi radialis. The rupture appeared to have occurred a little above the middle of the forearm. It was clearly impossible to make a good repair of the injured muscles, and in any case it seemed inadvisable to risk any avoidable post-operative swelling. Haemostasis was secured and the skin sutured; there was a little tension in places.

The limb was placed on pillows and active finger exercises were encouraged. Within a day the paraesthesiae had gone. In about four days the swelling of the hand had subsided. On the ninth day a plaster cast was applied to prevent flexion deformity at the wrist. Finger movements were improving but weak. After 24 days there were two hard masses of fibrous tissue in the forearm, and the fingers were showing a flexion deformity unless the wrist was palmar-flexed. Frequent stretchings were done by the patient and the physiotherapy staff. Splintage was at this time avoided as it was considered advisable to encourage finger movements continually. After seven weeks a night splint was used to keep the fingers straight with the wrist straight. After nine weeks the grip was 50 to 60% of normal. The index finger was flexed 30 degrees at each inter-phalangeal joint when the wrist was straight—the other fingers were straight. Ionization with potassium iodide was tried in the hope of reducing the scar tissue a little.

When seen three weeks later (Sept. 19)—i.e., three months after the injury—the index finger was straight when the wrist was straight, but only in the mornings after a night on a splint: after use it became flexed. With the wrist dorsiflexed all the fingers became slightly flexed. There was no palpable activity in the flexor carpi radialis. The patient was able to use the fingers well, both individually and together. He had an excellent grip, about 80% normal (he is right-handed). He was practising finer-finger movements on a typewriter and was making rugs.

Improvement was still taking place when he was last seen. While a degree of limitation of extension of the fingers and wrist seems inevitable, it is likely that he will have a very strong and useful hand with little disability after a period of two to three months.

This case is published by permission of D.M.S., B.A.O.R., and of O.C. 25 (Munster) Military Hospital.

G. M. SLEGGs, M.Ch.Orth., F.R.C.S.Ed.
Major, R.A.M.C.

REFERENCE

Mercer, W. (1943). *Orthopaedic Surgery*, 3rd ed., p. 876. Arnold, London.