SURGERY OF CONGENITAL HEART DISEASE*

BY

MAURICE CAMPBELL, M.D., F.R.C.P.

Physician, Guy's Hospital and National Hospital for Diseases of the Heart

The surgical treatment of some forms of congenital heart disease has raised many fresh questions of diagnosis and prognosis for the physician, as well as many problems for the surgeon. I shall deal with only some aspects of these.

Patent Ductus Arteriosus

The diagnosis of patent ductus is generally easy in the presence of the long continuous murmur through systole and diastole. The diagnosis may sometimes be made without these, but only after much experience; dilatation of the pulmonary artery, an increased pulse pressure at rest or after exercise, and a harsh systolic murmur will then be the main indications. In ordinary cases the two former are some guide to the amount of blood passing through the ductus. An adequate knowledge of the prognosis without operation is lacking, and we do not know with certainty whether the relative absence of cases over 40 is due to many deaths or to many cases closing spontaneously. A recent follow-up by Benn (1947) shows how few complications and symptoms arose in children over an eight-year period when the diagnosis was made at a routine school medical examination.

The first case of patent ductus arteriosus was ligated in 1939 by Gross, and experience has now been acquired in many centres. If there is no other congenital lesion the operation is entirely curative and the patient should be quite fit after it; the risks of infective endocarditis and of early heart failure should be entirely avoided. Surgical opinion is still divided between the question of ligation of the ductus and of resection with double ligation. The former is easier and should be the routine procedure, except in special circumstances, in spite of a small risk of recanalization.

The outlook is naturally different in the infected and non-infected cases. In the latter the mortality in skilled hands is low, perhaps 5%. In the infected cases the mortality is naturally higher, but even before the introduction of penicillin it was shown that surgical treatment was often an effective cure for such cases; the introduction of penicillin has made the risks in such cases much less. Many authorities recommend that in all cases in which the ductus has not closed spontaneously by the age of 7 an operation should be performed. Others, and I include myself among these, would limit it to those cases in which there is some enlargement of the heart, some striking fullness of the pulmonary artery, or some significant symptom.

The slightest degree of cyanosis or of right ventricular preponderance electrocardiographically should prevent a quick decision to operate, as the patent ductus may be an essential part of the circulation. After the age of 20 the

risks of operation are rather greater, but the chance of successful operation should not be refused if the symptoms appear to be troublesome or progressive. If infective endocarditis is present when the patient is first seen it should be arrested with penicillin treatment, and generally operation should be advised later because of the risk of recurrence.

Coarctation of the Aorta

The diagnosis of coarctation of the aorta will continue to be missed if a collateral circulation is always expected to be self-evident. The femoral pulse should be felt for and the blood pressure taken in the legs in every case in which high blood pressure, basal systolic or diastolic murmurs, or undue pulsation in the neck is present without an obvious cause. A high pressure in the arms and a low poor femoral pulse or the demonstration of a collateral circulation are pathognomonic of coarctation of the aorta. They are more important than notching of the ribs, not only because x-ray films are not always available but because the signs are present at an earlier age.

The collateral circulation can be more easily demonstrated or made more obvious if the patient stoops forward with his arms hanging down vertically while his back is examined in a good light. This is because the greatly dilated subclavian artery is compressed between the clavicle and the first rib and bending forward opens up this space and so releases the pressure on the subclavian artery.

We know more about the prognosis of coarctation. The average age at death is about 30. Two groups can be distinguished: those with symptoms before adult life, who will probably not live beyond 30; and those in whom the lesion is found by chance in adult life, the outlook being very much better. Most patients die from cardiovascular causes—heart failure, endocarditis, or a ruptured aorta—but one in ten may die from the rupture of a congenital cerebral aneurysm.

In October, 1944, Crafoord successfully resected a case of coarctation with an end-to-end suture of the aorta. In his first series (up to July, 1947) the results were entirely satisfactory in 20 and two patients died; in three patients after a thoracotomy no operation was possible and two of these died. In these 20 patients the blood pressure in the upper part of the body fell to or near to normal, the pressure in the legs rose, and the circulation was greatly improved as shown by oscillometric records. Some parts of the visible collateral circulation disappeared quickly and the patients felt better and were able to do much more.

It may be some time before the general operative risk can be assessed, but Crafoord's mortality of 16% and experience at other centres would justify a reasonable risk in those patients in whom the outlook seems unfavourable, considering the greatly improved expectation of life. The

*Read in opening a discussion in the Section of Diseases of the Chest at the Annual Meeting of the British Medical Association, Cambridge, 1948.

only sense in which the operation is not a complete cure is that bicuspid aortic valve or congenital cerebral aneurysms may be present. A fall of blood pressure to normal will greatly reduce the risks from the latter, but we cannot be sure that it will remove it entirely.

Atheroma develops in the aorta at a younger age in patients with coarctation; this adds to the risk of operation after the age of 20 or 25 years and is one of the reasons favouring early operation.

Cyanotic Congenital Heart Disease

Here the problems are much more difficult both for the physician and for the surgeon. Until recent years nothing could be done for these patients and they lingered on with increasing incapacity, some reaching a very limited adult life, but many died in infancy and more during childhood. Two centuries ago the association of cyanosis with pulmonary stenosis and with ventricular septal defect was recognized and there has been much discussion about their relative importance. It was Dr. Taussig's recognition of the inadequate flow of blood to the lungs as the fundamental cause of the disability that led to the idea of a successful operation, although the right-to-left shunt of venous blood actually produced the cyanosis.

Diagnosis

In early infancy many malformations of the heart may be responsible for gross cyanosis, but the more serious are not long compatible with life. As the children get older fewer forms of congenital disease are found with any frequency and this makes an attempt at diagnosis possible. Cyanosis from birth generally represents a right-to-left shunt either through a ventricular septal defect or because there is an overriding aorta that takes the blood from the right ventricle or, in an extreme case, where the aorta, being transposed, takes all its blood from the right ventricle.

The fundamental requisite for success of the Blalock-Taussig operation is that the cyanosis and disability should be mainly due to an inadequate blood-flow to the lungs, and when pulmonary stenosis is added to these other findings, as it is in Fallot's tetralogy, the operation is most certain to be successful.

Fallot's tetralogy consists of pulmonary stenosis (which may be valvular or infundibular), ventricular septal defect, an overriding aorta (that has often been called dextroposition), and right ventricular hypertrophy. The condition was very fully and accurately described by Peacock in 1858—20 years before Fallot's paper—and had been recognized earlier than this. These three abnormalities occurring together without pulmonary stenosis are known as Eisenmenger's complex. Pulmonary stenosis may also occur alone, but then there is generally no cyanosis from birth. Its later development probably depends on a septal defect which may be small and no more than a foramen ovale where the valve preventing a flow from left to right is overcome by the increased pressure on the right side so that there is a flow from right to left.

The majority of children with cyanosis from birth and clubbing of the fingers have Fallot's tetralogy, and often where the x-ray picture is characteristic this can be diagnosed with confidence. The main features are persistent cyanosis from infancy, with polycythaemia, clubbing of the fingers, and severe disability on walking. On radiography the heart, which may be sabot-shaped in half the cases, shows little enlargement generally but some enlargement of the right ventricle; there is no undue prominence of the pulmonary arc and no visible pulsation in the pulmonary branches. Most of them have a systolic murmur in the pulmonary area, often with a slight thrill. The

pulmonary second sound is never much increased and is often diminished. A diastolic murmur indicates some complication. Most of them have right ventricular preponderance in the electrocardiogram, with a large pointed P II. All are of normal mental development unless there are other causes for abnormality; most are underweight but of normal height. Squatting is noted in four-fifths of the patients, and panting after exertion is nearly as characteristic.

Although the diagnosis may be easy there are difficult borderline cases: the critical decision clinically is whether the pulmonary blood-flow as seen on radiography is diminished or not. Often it is easy to decide that this is diminished, as there is a hollow pulmonary bay and the lung fields look clear. In other cases (most familiar in the adult picture of auricular septal defect) it is obvious that the pulmonary arteries are dilated with increased pulsation. Difficulties are caused by the cases in which beyond the pulmonary stenosis there is some dilatation of the pulmonary arteries and in which a good collateral circulation produces increased density in the lung fields.

Fallot's tetralogy is the commonest finding among children who are obviously blue, but as one sees an increasing proportion of less cyanosed children there are more with other lesions and with a radiological picture similar to adult auricular septal defect. This defect alone cannot be the cause, because, the pressure being higher on the left than on the right, there is no cyanosis except perhaps terminally. Whether there are exceptions to this statement is one of the problems that remain to be solved, but more likely there is also transposition of the main vessels as well (Taussig, 1947). In Eisenmenger's complex also the pressure in the pulmonary artery is high, the blood-flow to the lungs is unimpeded, and operation will not help. In difficult cases great help may be obtained by the circulation times, cardiac catheterization, and by angiocardiology.

If the oxygen content of the blood in the right auricle is considerably higher than that in the superior vena cava there must be an auricular septal defect with a left-to-right shunt (a right-to-left shunt would not of course show by this method, though a catheter might pass through the septal defect). If the pressure is raised in the right ventricle and suddenly falls greatly as the catheter passes into the pulmonary artery, there is pulmonary stenosis, but it is not always easy to get the catheter into the pulmonary artery. A high pressure in the pulmonary artery often indicates Eisenmenger's complex, though it may occur in other conditions. A very high pressure on the right side, approximating to that in the aorta, suggests a septal defect with an overriding aorta and free communication between the two. Detailed calculations may often be made about the size of the blood-flow through the different circuits and shunts; while these are of great interest they are hardly reliable enough to be conclusive.

In angiocardiology quick filling of the left ventricle and of the aorta and its branches indicates a septal defect with a right-to-left shunt. Slow filling of the pulmonary arteries after the right ventricle has been visualized indicates pulmonary stenosis. In many cases other interesting details are observed, though considerable skill and experience is needed in the interpretation of the pictures. From such a simple investigation as the circulation times one may be certain of the presence of a large right-to-left shunt if the arm-tongue time is the same as or shorter than the arm-lung time.

Prognosis

There are few follow-ups of these children to give an accurate prognosis. If one may take the age incidence of a recent series as an indication, it is very unfavourable. In my first 137 patients (77 males and 60 females) with cyanotic

congenital heart disease there were almost equal numbers in each year up to 7, three-fifths of this number in each year from 8 to 11, and one-fifth of this number in each year from 12 to 22, after which there were only scattered cases. The figures suggest that those surviving infancy have no great risks until they reach the age of 7, though clinical experience hardly bears this out. After 7 the chances of surviving become less, and increasingly so after the age of 12.

If these figures can be taken as giving a true cross-section they mean that only one patient in ten reaches the age of 24 and only two in ten the age of 12. Many parents have been told that their children will not live to 7 or to 14 or to 21, and there seems some rough truth in these three round numbers as representing periods of increasing danger beyond which survival becomes less likely. During the last few months there have been some deaths after operation, but there have been many more among the other patients I have seen. These considerations seem ample justification for advising operation in spite of the relatively high mortality. The incapacity of many of these cases is hardly realized. A few may lead a reasonable but very quiet life, but if they are old enough to express their views they are generally the greatest enthusiasts for operation, even when the risks are put before them. Many, however, exist rather than live and are a handicap to their parents as well as to themselves, especially when they begin to realize what other children are able to do.

Operative Treatment

Following Taussig's suggestion about the importance of overcoming pulmonary stenosis, Blalock's experimental work on animals convinced him that a subclavian-pulmonary anastomosis was a feasible method, and in October, 1944, Blalock and Taussig introduced their operation for the relief of the inadequate blood-flow to the lungs in Fallot's tetralogy and related lesions.

The operation is essentially an anastomosis between a systemic artery and the pulmonary artery, most commonly an end-to-side subclavian-pulmonary anastomosis, though under special circumstances the innominate or carotid artery may be used or the anastomosis may be end-to-end. In one-quarter of these cases the aortic arch will be right-sided instead of curving back as usual to the left and this will of course change the position of the innominate artery. The operation should be performed on the side opposite to the aortic arch because this gives a greater choice of systemic vessels that can be used and the subclavian artery can generally be turned down to the pulmonary artery with a less acute angle.

In infants the small size of the vessels produces a surgical problem of the greatest difficulty, and the operation is much more dangerous and sometimes less lasting in its good effect because the anastomosis is too small as the child grows larger. For this reason the operation should never be performed before the age of 3, except as a life-saving emergency, and generally not till 5. From 5 to 10 is perhaps the ideal age, but there seem no special difficulties in older children, though the difficulties and risks are greater in those over 20—a group that should not trouble our successors, as suitable cases should have been operated on earlier. The direct division of the stenosed pulmonary valve by Brock (1948) has opened up fresh possibilities. The late results of this and the indications for it remain to be worked out.

Results of Operation

The operation is not of course curative in the same way as in the other two groups because the underlying abnormality remains, but the improvement may be so great that the child appears normal to the parents. The immediate

upset caused by the operation is less than might be expected, and intravenous fluid is often not required; when it is used only small amounts are needed. Many have a pleural effusion that often needs aspiration, but generally only once; other complications are not common. Of the first 38 cases operated on at Guy's Hospital 6 died, 7 could not have an anastomosis done or did not benefit much, and the remaining 25 were enormously improved almost at once, and this has been maintained.

The immediate results of the operation are excellent, and the improvement seems to be maintained over the period of two to three years for which it has been possible to follow up Blalock's patients. The colour becomes nearly normal except sometimes for slight cyanosis on a cold day or after vigorous exercise, and the patients' capacity is greatly improved so that they can walk some miles and get about normally all day. The polycythaemia disappears quickly, the clubbing of the fingers more slowly. The arm on the side of the divided subclavian artery gives no trouble.

The heart enlarges a little as might be expected from the increased work, but generally only slightly. Sometimes at the beginning the increase during the first few weeks made me feel that any such progressive increase would soon end in disaster, but I have seen no cases in which the increase has continued after the first month or two, and I understand that in the longer American experience this is a rare occurrence, though relapses in those operated on too young and a few instances of congestive heart failure have been seen.

REFERENCES

- Benn, J. (1947). *Brit. Heart J.*, **9**, 283.
Blalock, A., and Taussig, H. B. (1945). *J. Amer. med. Ass.*, **128**, 189.
Brock, R. C. (1948). *British Medical Journal*, **1**, 1121.
Crafoord, C., and Nylin, G. (1945). *J. Thorac. Surg.*, **14**, 347.
Gross, R. E. (1939). *Ann. Surg.*, **110**, 321.
Peacock, T. B. (1858). *Malformations of the Heart*, 1st ed., London.
Taussig, H. B. (1947). *Congenital Malformations of the Heart*, New York.

LARGE-INTESTINE COLIC DUE TO SYMPATHETIC DEPRIVATION

A NEW CLINICAL SYNDROME

BY

Sir HENEAGE OGILVIE, K.B.E., M.Ch., F.R.C.S.

Surgeon to Guy's Hospital

The subtitle of this note is intentionally provocative. Two cases are described, presenting essentially similar features. In each case the symptoms so strongly suggested obstruction of the distal colon by a carcinoma that laparotomy was eventually undertaken in spite of normal findings in a barium enema. In each case the operation disclosed a normal colon and extensive and entirely unsuspected malignant disease involving the region of the crura of the diaphragm and the coeliac axis and semilunar ganglia.

Whether such a syndrome has been described before I am unaware, but I have not encountered it apart from these two cases, nor have any of my friends whom I have consulted on the subject, and a search of the journals over the last ten years and of the literature of hypertension has failed to reveal similar observations.

Case 1

In 1944 a physician aged about 66, on the staff of an E.M.S. hospital, began to lose weight, and complained of increasing constipation. He was found to be diabetic, and was treated with diet and insulin. The diabetes was controlled, but the constipation persisted, and he began to have colicky pains in the hypogastrium. On several occasions waves of peristalsis,