July 1, 1946, to Dec. 31, 1947, are analysed, and show that, including two major epidemics during this period, out of 631 cases 98.4% were microsporon infections (97.1% *M. audouinii*) and 84.15% were in boys. These figures correspond closely to recent figures from other parts of Scotland and from the U.S.A. Infection most commonly appeared to occur in barbers' shops.

Wood's light was found essential in the diagnosis and management of microsporon and favus infections but of no use in trichophyton cases.

Epilation by x rays or thallium is still the most rapid and successful means of cure, but fungicides assisted by a wetting agent in carbo-wax 1500 offer a chance of cure in at least a fair proportion of cases after a considerably longer period of treatment. It must be realized that, especially with the animal microspora, a modified kerion reaction may occur leading to spontaneous cure, so that in assessing the value of any "cure" for ringworm the actual fungus causing the condition and the criteria of cure must be known.

A brief note is made of the organization required for a centre to deal with cases of ringworm and the standards necessary for ensuring that patients are non-infectious before their return to school.

REFERENCES

- Anderson, M. (1871). *Lancet*, 2, 672, 742.
 Cantanei, A. (1947). *Arch. Inst. Pasteur Algér.*, 25, 1. Quoted in *Excerpta Medica*, 1947, 1, Sec. XIII, 717.
 Chmel, L. (1946). *Bratislavské lekár. Listy*, 26, 601. Quoted in *Excerpta Medica*, 1947, 1, Sec. XIII, 716.
 Duncan, J. T. (1945). *British Medical Journal*, 2, 715.
 Keddie, J. A. G. (1947). *Hlth Bull. (Dept. of Health for Scotland)*, 5, 66.
 Lewis, G. M., Hopper, M. E., and Reiss, F. (1946). *J. Amer. med. Ass.*, 132, 62.
 Miller, J. L., Lowenfish, F. P., and Beattie, G. F. (1946). *Ibid.*, 132, 67.
 Schwartz, L., Peck, S. M., Botvinick, I., Leibovitz, A. L., and Frasier, E. S. (1946). *Ibid.*, 132, 58.
 Walker, N. (1899). *An Introduction to Dermatology*, 1st ed., p. 145. Wright, Bristol.
 — and Percival, G. H. (1939). *Ibid.*, 10th ed., p. 178. Green, Edinburgh.

CIRRHOSIS OF LIVER PRESENTING AS SEVERE ANAEMIA REPORT OF TWO CASES

BY

P. N. COLEMAN, M.B., Ch.B.

Pathologist, Townleys Hospital, Bolton, Lancs.

This paper concerns two cases of cirrhosis of the liver in which a severe anaemia, thought to be haemolytic, was the predominant feature. Both cases were sent to hospital because of anaemia, and cirrhosis of the liver was not diagnosed until after further investigation, in the second case not until necropsy. They are reported because haemolytic anaemia is said to be an uncommon complication of cirrhosis.

The anaemia accompanying cirrhosis of the liver is described as normocytic or macrocytic, and haemolysis is not a feature. The anaemia is seldom severe. Macrocytic cases resemble pernicious anaemia in some features (Whitby and Britton, 1946; Wintrobe, 1936; Wintrobe and Shumacker, 1933; Singer and Dameshek, 1941).

Davidson and Fullerton (1938) report a case of a macrocytic haemolytic anaemia in cirrhosis in which red-cell fragility was increased. Watson (1939), from studies of urobilinogen excretion, emphasizes that in some cases of cirrhosis a haemolytic element is present. He describes seven out of 38 cases of cirrhosis studied in which there was evidence of haemolysis. One of these had two severe haemolytic crises, during the second of which the patient died. Haemoglobin fell to the low level of 25%. Necropsy

revealed a diffusely cirrhotic liver, which on section showed portal cirrhosis. The liver was described as quite fatty. Femoral marrow was red.

Singer and Dameshek (1941) include disease of the liver among the relatively few conditions that can give rise to symptomatic haemolytic anaemia. They describe a case of subacute hepatitis in which a severe haemolytic anaemia was an important symptom: the haemoglobin fell to 27%, there was marked reticulocytosis, sternal puncture revealed a normoblastic hyperplasia, the anaemia was macrocytic, and fragility was normal.

Case 1

The patient, a man aged 66, was admitted to Townleys Hospital on Oct. 2, 1947, because of extreme pallor and shortness of breath. He had been working as a lorry driver up to three weeks before admission. He gave a history of vague abdominal pain and lack of appetite for several months. He had been a heavy beer-drinker for 30 years. He was extremely pale with an icteric tinge. Liver was enlarged to the umbilicus and the spleen was palpable two fingerbreadths below the costal margin. A blood count showed: Hb, 18%; red cells, 890,000; C.I., 1. The red cells appeared normal in size and shape. Biochemical findings supported a diagnosis of cirrhosis of the liver. The van den Bergh test gave a positive direct reaction; total serum bilirubin was 2.6 mg. per 100 ml. Thymol turbidity was 12 units. Serum globulin was raised to 4.1 g. per 100 ml., serum albumin was 3.3 g. per 100 ml., and the albumin-globulin ratio was reversed. Urine gave a strong reaction for urobilinogen and contained a trace of bilirubin. Tests for occult blood in the faeces were negative on three occasions.

The patient received a series of blood transfusions until he died 21 days after he was admitted to hospital. Although he showed no obvious reaction to these transfusions it soon became evident that the transfused cells were rapidly destroyed, because the rise in haemoglobin obtained was very small. Twelve days after admission to hospital he had received 11 pints (6.25 litres) of blood and his haemoglobin had varied between 28 and 35%. Shortly before he died he received the packed cells from 3 pints (1.7 litres) of blood and the haemoglobin was raised to 45%.

At necropsy the liver was of normal size and showed a diffuse cirrhosis; the spleen was enlarged to twice normal size. Section showed a portal cirrhosis with the architecture of the lobules destroyed by numerous bands of young fibrous tissue insinuated between the liver cells. There were considerable deposits of iron pigment in the liver and spleen.

Case 2

A woman aged 58 was admitted to Townleys Hospital on April 1, 1948, because of pallor and dyspnoea. She gave a history of vague ill-health for over a year and had recently had a small vaginal haemorrhage. She had been living on a very poor diet for a long time and said the only thing she really cared for was a glass of beer.

On examination the patient was pale, slightly icteric, and looked very ill. There were purpuric spots on forearm, thighs, and buttocks. She was obese, but it was thought that a tumour (? liver) could be felt in the right hypochondrium. Pelvic examination revealed nothing abnormal. The blood count on admission was as follows: Hb, 29%; red cells, 1,410,000; C.I., 1.03; reticulocytes, 22.5%; white cells, 4,600; normal differential; platelets, 160,000; bleeding and clotting time, normal. There was a little anisocytosis, but the majority of red cells appeared normal in size. Five orthochromatic normoblasts were seen while counting 100 white cells. Biochemical findings were as follows: van den Bergh test gave a positive direct reaction, serum bilirubin 3 mg. per 100 ml.; direct/indirect quotient (Gray, 1947) 68. Thymol turbidity was normal (4 units), alkaline phosphatase normal (10 units), and urine gave a very strong reaction for urobilinogen and contained a trace of bilirubin. No occult blood was found in the stools. The patient was given 3 pints of blood, which raised her haemoglobin to 55%, but her clinical condition showed

very little improvement. A week after the transfusion a fragility curve (Creed's method) showed a definite increase in fragility, the haemolysis in 0.5% saline being still 8%. Patient died 11 days after admission to hospital.

Post-mortem findings were as follows: the liver was slightly larger than normal and showed cirrhosis. The colour was a green-brown and the surface was covered with small nodules. The spleen was congested and enlarged to twice the normal size. Section showed a portal cirrhosis with young fibrous tissue breaking up the lobules. There was infiltration of the fibrous tissue with lymphocytes. Quite a large proportion of the liver cells were distended by globules of fat. There were deposits of iron in the liver and spleen. The femoral marrow was red throughout its length and showed a normoblastic hyperplasia.

Discussion

In both cases a severe anaemia has to be explained. Haemorrhage can be discounted. Tests for occult blood in the stools were negative. In the first case there was no history of bleeding; in the second there had been a small vaginal haemorrhage and purpuric spots, but nothing which could account for a haemoglobin of 29%. The appearance of the blood films was quite unlike that of pernicious anaemia. Case 1 received an urgent blood transfusion on admission and a reticulocyte count was not done, but there is evidence of a haemolytic process in the remarkable behaviour of this case on transfusion. A total of 14 pints (8 litres) of blood was given to this patient over a period of 14 days and relatively little increase was obtained in the haemoglobin. This can be explained only by a rapid destruction of transfused cells, which is one of the features of haemolytic anaemia (Sharpe and Davis, 1938). In the second case there is clear evidence of a haemolytic anaemia: a high reticulocyte count, numerous nucleated red cells, increased fragility, and red marrow in the shaft of the femur.

The biochemical findings do not assist in the diagnosis of the anaemia. The positive direct van den Bergh reaction and the presence of bilirubin in the urine are not, of course, the findings of haemolytic jaundice, but can readily be explained on the basis of the coexistent liver damage. The large amounts of urobilinogen in the urine are more likely to be due to the cirrhosis than to the haemolysis.

My thanks are due to Dr. E. S. Gawne, medical superintendent, Townleys Hospital, for permission to publish these case reports, and to Drs. D. Mooney and R. Kearney for the opportunity to co-operate in the diagnosis and treatment.

REFERENCES

- Davidson, L. S. P., and Fullerton, H. W. (1938). *Quart. J. Med.*, n.s., 7, 43.
 Gray, C. H. (1947). *Ibid.*, n.s., 16, 135.
 Sharpe, J. C., and Davis, H. H. (1938). *J. Amer. med. Ass.*, 110, 2053.
 Singer, K., and Dameshek, W. (1941). *Ann. Intern. Med.*, 15, 544.
 Watson, C. J. (1939). *Ibid.*, 12, 1782.
 Whitby, L. E. H., and Britton, C. J. C. (1946). *Disorders of the Blood*, 5th ed., p. 240. London.
 Wintrobe, M. M. (1936). *Arch. Intern. Med.*, 57, 289.
 — and Shumacker, H. S. (1933). *Bull. Johns Hopk. Hosp.*, 52, 387.

The village of Syrianokhorio in Cyprus has recently celebrated the eradication of malarial-carrying mosquitoes with religious services and a meeting at which the authorities were thanked, reports the Colonial Office. The offensive against mosquitoes was started by the Cyprus Government Medical Department in 1946, and is aimed at complete elimination of malaria from the island by 1952. The estimated cost is £300,000. The terrain is a difficult one to work in and covers about 3,500 square miles; the western half is mountainous, with steep precipices and rushing streams. *Anopheles superpictus* breeds in the streams and rock pools in almost inaccessible places. The country is divided into zones which are systematically sprayed with a solution of D.D.T. in oil.

TREATMENT OF ABORTUS FEVER WITH SULPHONAMIDES AND BLOOD TRANSFUSIONS

BY

J. MacD. HOLMES, M.D., M.R.C.P.

Physician, Staffordshire General Infirmary;
Consulting Physician, Crewe and District Memorial Hospital

AND

ROBERT HUGHES, M.D., M.R.C.P.

Visting Physician, Clatterbridge (County) General Hospital;
Honorary Assistant Physician, Royal Liverpool United Hospital
(Royal Southern Hospital Branch)

The treatment of undulant fever has remained unsatisfactory in spite of recent advances in chemotherapy, although occasional successes have been reported from the use of sulphonamides, suramin, vaccines, and injections of brucellin. Recent reports of treatment with penicillin and streptomycin are also very disappointing. In an annotation in the *British Medical Journal* (1947, 2, 700) attention was called to a report by Huddleson, made to the Fourth International Congress for Microbiology at Copenhagen in July, 1947, of four cases of undulant fever which responded quickly to the administration of small doses of sulphonamides together with blood transfusions. No special care was given to the selection of donors; it was assumed that all human serum contains bactericidal brucella antibody, but it is difficult to understand why a weak antiserum in conjunction with a sulphonamide should have a bactericidal effect. Huddleson explains this on the basis of the Neisser-Wechsberg phenomenon—a serum may be bactericidal in high dilution yet not in low dilution.

This method of treatment was tried in two cases of undulant fever with the encouraging results indicated in the following reports.

Case 1

The patient, a woman aged 32, first complained of feeling ill about Dec. 16, 1947. A week later she was found to have an evening temperature rising to 102° F. (38.9° C.). She was seen by her doctor on Jan. 17, 1948, when she complained of tiredness, headaches, and pain in the back. On examination at this time her temperature was 100° F. (37.8° C.) and her pulse rate 86 per minute; she had lost a little weight and her spleen was much enlarged.

On Jan. 8 she was admitted to Clatterbridge (County) General Hospital, where re-examination confirmed the above findings. Her evening temperature ranged from 100.4° to 102.6° F. (38°–39.2° C.), but apart from the fever and the splenic enlargement no abnormality was found. On Jan. 19 the blood urea was 26 mg. per 100 ml. A blood count showed red cells, 5,000,000 per c.mm.; haemoglobin, 92%; white cells, 3,600 per c.mm. (polymorphs 52.5%, lymphocytes 45%, monocytes 2.5%). Agglutinations: positive in all dilutions up to 1:1,400 against *Br. abortus*; negative against *Salm. typhosus* and *paratyphosus*. The urine was free from sugar and albumin. Blood culture: contaminated by *Staph. albus* (coagulase-negative), no other organisms found. X-ray film of the chest was normal. Blood Wassermann reaction was negative.

The following treatment was given: penicillin, 9 mega units (Jan. 25 to Feb. 3), sulphadiazine, 75 g. (Feb. 7 to 19), and blood transfusions of 10 oz. (280 ml.) on Feb. 11, 13, 15, and 17. There was no clinical improvement with penicillin, but once the sulphadiazine and transfusions were begun the temperature rapidly subsided and the patient's condition improved. She was discharged from hospital on Feb. 26 and reported to the outpatient department on April 14. She had been afebrile since discharge, felt well in every way, and there was no splenic enlargement.