BRITISH MEDICAL JOURNAL

LONDON SATURDAY NOVEMBER 27 1948

PHARMACOLOGY OF THE FAILING HUMAN HEART*

BY

JOHN McMICHAEL, M.D., F.R.C.P.

Professor of Medicine, Postgraduate Medical School of London

Digitalis and Strophanthus

It was a long time after Withering's introduction of digitalis into therapeutics in 1775 that its key position in the treatment of heart failure was recognized. Mackenzie (1905) analysed its action in the treatment of auricular fibrillation and ascribed the major benefits of the drug to its slowing effect on the heart in this condition. This view is still held in some quarters, and Lewis was long unconvinced that it had any other important action. Clinicians who used it in the treatment of failure with sinus rhythm paid a great deal of attention to its slowing action on the heart rate, and it was often held that any benefits achieved were the results of such slowing. A beneficial action on the heart independent of slowing effects, however, was postulated by observant clinicians like Christian (1919) and Gavey and Parkinson (1939). Rate reduction accompanying improvement of the circulation might in fact be the secondary result of such improvement.

The action of digitalis on the heart was a matter of considerable perplexity to pharmacologists. Cohn (1915) said that if digitalis stimulated the heart it did so by some means that pharmacological methods were unable to detect: the application of digitalis to the heart of the cat might or might not have a stimulating action. Cushny (1918) suggested that, while digitalis did not act directly on the normal heart, it had some specific action on the "malnourished" failing heart. Sir Thomas Fraser in 1890 observed: "The production (by strophanthin) of this increase in the movements of the heart . . . is significantly emphasized when the action of strophanthin is produced in an enfeebled and insufficiently acting heart. This experiment is therefore of peculiar value in demonstrating the therapeutical value of strophanthus in heart disorders." Evidence that the failing heart was stimulated by digitalis bodies was produced by the use of the heart-lung preparation by Bodo (1928), Visscher (1938), Gremels (1933), Anitschkow and Trendelenburg (1928), and others, although some workers such as Katz *et al.* (1938) produced some experimental evidence that failure of the isolated heart was uninfluenced by digitalis.

Into this perplexing confusion of ideas was thrown the observation by Harrison and Leonard (1926) that in the whole animal and in man the action of digitalis was to lower the venous pressure and with it the output of the heart. It was thus clear that there was no evidence of any stimulating action of digitalis on the normal heart. Although the pharmacologists had offered a useful working idea of

stimulation of the failing heart, the questions remained: (1) Does the "failing" heart which the pharmacologist produces in his laboratory by such methods as subjection of the animal heart to the action of poisons like camphor and large doses of chloral truly represent the condition of the failing heart as seen in clinical practice? (2) What is the explanation of the lack of response in some cases of heart failure to the administration of digitalis? These were among the questions which seemed to us to demand an answer when new methods of studying the cardiac output become available.

The technique of right heart catheterization, introduced by Forssmann (1929) and developed by Cournand and Ranges (1941) for the estimation of cardiac output, constitutes the most significant advance in cardiological method in the present generation. It might be thought that by such a method, which allows the measurement of the minute volume of blood pumped by the heart and the pressures at which it does this work, the problem of digitalis action on the failing heart would be readily solved. But, like so many new methods in medicine, extensive application of the technique simply revealed that the problem of congestive failure had unrealized complexities. The clinical picture of cardiac failure with congestion and oedema is now being resolved into a number of fundamentally different groups, the mechanism of each being differently from the others.

The first thing to realize is that general venous congestion may result from conditions which demand an increase in the cardiac output as well as from conditions which have long been known to impair the cardiac mechanism and which were assumed to reduce the output of the heart. We can thus subdivide cardiac failure into:

(1) *Failure with a Low Output.*—Due to hypertensive, ischaemic, and valvular heart disease.

(2) *Failure with High Output.*—Due to conditions which demand an increase in the output of the heart, such as: (a) diminished oxygen-carrying power of the blood—anaemia, emphysema; (b) metabolic disturbances—beriberi; (c) conditions producing a mechanical overload—arteriovenous aneurysms, Paget's disease of the bone.

In the high-output group the venous-pressure rise is not a back-pressure effect, and must rather be regarded as compensatory, the high venous pressure maintaining the output of the heart at a necessarily high level.

The reactions of these two groups to digitalis show fundamental differences. If we take emphysema heart failure as an example of the high-output group it is found that digitalis lowers the venous pressure and with it the output of the

*The Strickland Goodall Lecture delivered to the Society of Apothecaries on June 3, 1948.

heart (Fig. 1). The same type of response is seen to occur with venesection (Fig. 2). This is exactly similar to the response of the normal heart to digitalization (Fig. 3). It has long been known that digitalis reduces the venous pressure and cardiac output of normal animals and man (Harrison and Leonard, 1926; Dock and Tainter, 1930; Katz *et al.*, 1938; Rytand, 1933; Wood, 1940).

FIG. 1.—The action of digoxin in emphysema heart failure. Note that the cardiac output is raised above the normal average of 5.3 litres per minute. Digoxin lowers the filling pressure of the right auricle, and with it the cardiac output. (Reproduced by permission of *Clinical Science*.)

FIG. 2.—Emphysema heart failure. The cardiac output was 8.4 litres per minute. Venesection reduced the right auricular pressure, and cardiac output fell to 6.7 litres per minute.

FIG. 3.—The action of 1.5 mg. digoxin on the normal heart. Right auricular pressure and cardiac output fall in parallel. (Reproduced by permission of *The Quarterly Journal of Medicine*.)

The influence of the venous filling pressure of the heart in determining the diastolic volume, and thereby the capacity of the stretched fibres for work, is a fundamental part of Starling's law: the cardiac output of the normal heart rises with an increase, and falls with a decrease, of venous pressure. Digitalis reproduces on the normal heart exactly what we would expect to happen if digitalis had a primary action on the venous tone whereby the venous pressure was reduced: the heart would diminish in size and the cardiac output would fall. It seems reasonable to postulate, therefore, that digitalis has such a primary venous-pressure-reducing action in man, and that the normally responding heart diminishes its output in response to this venomotor effect.

Digitalis, then, reduces the venous filling pressure and with it the output of the normal heart. It acts in a similar manner in most instances of high-output heart failure where the rise of venous pressure should be regarded as "compensatory."

Action of "Digoxin" in Low-output Heart Failure

If we turn now to the group of patients with heart failure and *low output*, the usual late consequence of valvular, ischaemic, and hypertensive heart disease, we find a very different state of affairs so far as the response to digitalis is concerned. The most favourable responses in this group are obtained in patients in the early stages of hypertensive and ischaemic heart disease—a point which was emphasized many years ago by Marvin (1927). In the later attacks of failure the response to digoxin steadily diminishes. At the responsive stage the injection of digoxin reduces the venous pressure, and the cardiac output rises. It would be easy to say that this is evidence of a stimulating action of digitalis on the failing heart; but simple venesection, reducing venous pressure, in these subjects will also *increase* the cardiac output. Since we already have evidence of a primary venous-pressure-lowering action of digoxin in normal man, it is necessary to see whether such an action might not be responsible for some of the effect on heart failure. Sharpey-Shafer and I (1944a, 1944b) proceeded to do this by the application of constricting cuffs to the thighs, damming back blood in the legs, thereby producing a mechanical lowering of venous pressure in the right auricle similar to that achieved by venesection. When this was done we found that the cardiac output change produced by a given venous-pressure reduction was exactly the same as that produced by a subsequent injection of digoxin when comparison of effects was made at the same venous-pressure level. This was not only true in the individual case but also when a whole series of venesections were compared with a series of digoxin observations in similar cases (Fig. 4). It is of interest to note at this point that patients with auricular fibrillation, where rate control was added to the other actions of digitalis, showed no greater or better immediate cardiac output response than the group with sinus rhythm, in whom rate reduction was slight or absent.

If our comparison of the effects of digoxin with those of mechanical reduction of venous pressure by this method is a valid controlled experiment, then it would appear that a primary venous-pressure-reducing action of digoxin might be mainly responsible, in man, for its effects on cardiac output in low-output heart failure. With digoxin we have seen no evidence against this view. It seems that the failing heart may behave in a manner very similar to Starling's overloaded heart-lung preparation, where further rises of venous pressure induce a fall in cardiac output and conversely a fall in venous pressure may be accompanied by

FIG. 4.—The upper graph shows the effects of venesection in lowering right auricular pressure and raising cardiac output in low-output heart failure. The lower graph shows, on an identical scale, the effect of digoxin on right auricular pressure and cardiac output in a similar group of cases. There are no significant differences. (Reproduced by permission of *Clinical Science*.)

some recovery. As a working hypothesis this enables us to explain some of the apparent failures of digoxin in clinical practice. There are advanced cases of heart failure in which the heart does not respond to venesection, and in such cases we find that it does not respond to digoxin either. We have already demonstrated that in emphysema hearts the response to digitalization is often a reduction of output. The ineffectiveness of digitalis in cases of pulmonary heart failure was recognized by Vaquez and Lutembacher (1928): death has, in fact, resulted from the use of full doses of digoxin intravenously in emphysema heart failure.

Do Digitalis and other Cardiac Glycosides have any Stimulating Action on the Failing Human Heart?

Although the effect of digitalis on cardiac output and venous pressure may be exactly simulated by simple mechanical measures reducing the venous pressure, there is

in fact a difference when we consider the *work* of the heart after the two procedures. The work of the heart can be conveniently approximated for purposes of comparison by taking the product of output and mean blood pressure against which blood is ejected. Venesection is followed by a fall of blood pressure, while digoxin often produces a rise in pressure. Although the cardiac outputs are similar after the two procedures, it is clear that after digoxin the heart is doing more work than after venesection (Fig. 5). We have no easy means, however, of testing the venesected heart against an increased arterial pressure without introducing further complicating drugs. Detailed comparison of the responses, to see whether this increase in cardiac work truly indicates a stimulating action of digoxin, was not possible.

In recent months, however, we have approached this problem again, using ouabain (g-strophanthin). Two years ago it was suggested to me by Professor Rothlin, of Basle, that strophanthin might give answers which were different from those obtained by digoxin. Last year, in Boston, Drs. Bloomfield and Ellis showed me the results of a few experiments with ouabain which certainly suggested that the drug might have a direct stimulating action on the heart.

With my colleagues at the Postgraduate School we have pursued the matter further, and the present position may be stated as follows:

1. Ouabain has little measurable effect on the filling pressure or output of the normal heart.
2. It has *qualitatively* similar actions to digoxin on the failing heart.
3. In about half the cases of heart failure, however, there is a definite increase in cardiac output with only a very slight reduction of venous pressure. Detailed examination of these cases shows an undoubted cardiac stimulating action of ouabain. In the individual case the output of the heart may be raised very significantly beyond that achieved by mechanical venous-pressure reduction (Fig. 6).

FIG. 5.—The effects of venesection and digoxin on cardiac work (cardiac output multiplied by mean arterial pressure). The heart is doing more work after digoxin than after venesection.

4. The increases in cardiac work after strophanthin are often greater than anything seen after digoxin. This is partly the result of a greater pressor action, the rise in arterial pressure often being quite pronounced.

5. In emphysema heart failure ouabain tends to increase the output of the heart in contrast with digoxin, which very frequently lowers the output in such cases.

It seems, therefore, that ouabain has a direct stimulating action on the human heart in certain cases of failure. This stimulating action can be seen in some instances to be quite independent of any significant venous-pressure-reducing

the heart in high-output "failure" (where the venous-pressure rise is "compensatory"). This action raises the output of the heart in low-output heart failure in a manner exactly similar to that achieved by venesection.

(2) A direct stimulating action which is seen only in certain forms of failing heart and which is significantly more conspicuous with ouabain than with digoxin. The conditions of occurrence of this stimulating action are as yet indefinable.

(3) A rate-reducing action in auricular fibrillation which does not appear to enhance particularly actions (1) and (2), but which may be an excellent guide to proper maintenance dosage when the drug is given over a long period of time.

FIG. 6.—Comparison of venous-pressure reduction with ouabain in a case of hypertensive heart disease. Reduction of venous filling pressure by mechanical means (cuffs on thighs) had no significant influence on cardiac output. 1 mg. of ouabain intravenously raised the cardiac output considerably at a similar level of venous pressure (50–60 mins.). An immediate consequence of ouabain was a transient rise in arterial pressure from 136/90 to 190/114. During this phase the venous filling pressure remained high, but in spite of this cardiac output was definitely raised. This is clear evidence of a direct stimulating action of ouabain on the myocardium.

effect. Accepting analogous actions of digoxin, I think it is quite possible that the effect previously recorded of increased cardiac work after digoxin is the result of a similar but significantly weaker stimulating action on the myocardium.

Summing up the evidence, then, leaving out effects on vagus tone and cardiac rhythm, the actions of cardiac glycosides influencing cardiac contractility are as follows:

(1) Venous-pressure-reducing action, most pronounced with digoxin, which lowers the output of the normal heart and of

If asked whether the venous-pressure-reducing action or the stimulating action is more important, I should have to answer under two headings: (1) So far as digoxin is concerned, it is the venous-pressure-reducing effect which predominates, for the following reasons: (a) In certain forms of heart failure (emphysema heart) the venous-pressure action is predominant in controlling the response and the cardiac output usually falls. (b) Unless venous-pressure reduction is achieved the effect of digoxin on the cardiac output is negligible. In general, the response of the heart to venous-pressure change seems largely to determine the response of the cardiac output to digoxin. (2) With ouabain, on the other hand, it is the stimulating action which tends to predominate, the venous-pressure-reducing action being weaker.

The stimulating action of strophanthin on the heart was clearly demonstrated by Sir Thomas Fraser in his monumental study in 1890. Its use in therapeutics we owe largely to the sustained advocacy of Fraenkel, ably backed by Vaquez. Vaquez claimed that the intravenous injection of strophanthin would help patients in whom digitalis had failed. Subsequently Vaquez popularized ouabain, which was of similar composition but more highly soluble in water. Vaquez summed up this situation in a manner which is completely in keeping with our recent researches. He recognized the uselessness of digitalis in pulmonary heart disease, and said, "Here again, with less success, however, ouabain can be advantageously substituted for digitalis" (Vaquez and Lutembacher, 1928).

Vaquez also claimed that a few days' use of ouabain might restore the responsiveness of a failing heart to digitalis when the latter drug had apparently completely failed. I can certainly say that I have observed this sequence.

Before drawing the conclusion, however, that restoration of digitalis sensitivity is specifically achieved by ouabain, I think we should keep in mind the recent work of Ray and LaDue (1945), who found that many patients who went into failure while on maintenance dosage of digitalis were in fact not receiving adequate dosage of the drug. Even in the presence of digitalis effects in the electrocardiogram, further intravenous injections of a preparation of *Digitalis lanata* might bring the failure under control. A great deal more work would therefore have to be done to prove

that ouabain is more effective than further digitalis dosage in these circumstances.

One of the reasons why strophanthin preparations have so long been neglected in this country is that they are very irregularly absorbed when given by mouth: consequently intravenous injection is necessary, and this makes their continuous use a matter of difficulty. Our own observations to date are concerned with immediate responses and do not deal to any extent with the more prolonged therapeutic use of the drug. Although it is clear that it is a potent cardiac stimulant, it has also an active blood-pressure-raising action which may actually precipitate further breathlessness after injection. It seems probable that small repeated doses may avoid this sudden extra arterial load on the heart and thus may be more beneficial than a single full dose.

Theophylline

The xanthine derivatives have had an honourable place in the therapeutics of heart failure for a long time. In the immediate past two main actions have been ascribed to them

only a slight increase in cardiac output the subsequent injection of theophylline produces a much greater rise in the minute-volume of the circulation (Fig. 7). Theophylline may be given with or after digoxin with mutual enhancement of their beneficial actions.

This stimulating action of xanthine derivatives on the heart was noted by Curschmann in 1873, when he observed that the apex beat became forcible and heaving in a woman who drank a decoction of half a pound of coffee in an effort

FIG. 7.—Comparison of venous-pressure reduction and theophylline-ethylene-diamine. Venous-pressure reduction produced a slight rise in cardiac output; theophylline-ethylene-diamine ("cardophylin") produced a greater rise.

—diuretic and coronary vasodilator. In combination with ethylene-diamine it has a remarkable action in the abolition of Cheyne-Stokes respiration (Marais and McMichael, 1937). Using it initially for this respiratory action, Sharpey-Schafer and I observed that it had a profound venous-pressure-lowering influence accompanied by an equally remarkable increase in cardiac output in a patient with hypertensive heart failure. We followed up that observation with a study of this action in a variety of different conditions (Howarth, McMichael, and Sharpey-Schafer, 1947). Theophylline has a stimulating effect on the normal heart as well as in heart failure; the stimulating action is better seen in hypertensive heart disease than in mitral stenosis. It also has a venous-pressure-reducing effect which may sometimes be seen to occur in mitral stenosis unaccompanied by any significant change in cardiac output. When we make a comparison between the effects on the heart of venous-pressure reduction and of theophylline there is no doubt of the direct stimulating action on the heart. When venous-pressure reduction brings about

FIG. 8.—The effect of sodium mersalylate on cardiac output and right auricular pressure; 0.2 g. injected intravenously at the arrow. The diuresis reached a peak of 500 ml./hour at the fourth hour, and thereafter continued at a lower rate up to the tenth hour. By the seventh hour the right auricular pressure was considerably reduced and cardiac output had risen from 2.5 to 3.7 lit./min. (Reproduced by permission of Drs. L. G. C. Pugh and C. Wyndham.)

to produce a miscarriage. The cardiac stimulating action was also frequently noted in animal experiments, and this effect was emphasized by Boyer (1943), who thought any coronary vasodilatation seen in experimental work was a secondary consequence of the increased activity of the heart. The action of the drug is extremely rapid, coming on within five minutes of the injection. In view of the widespread use of the substance as a coronary dilator it is perhaps as well to sound a warning note that stimulation of the heart in a patient with a recent coronary thrombosis may have untoward consequences. Cases of sudden death have been reported following its use with this purpose in mind.

Theophylline, while of immense value in emergency treatment of severe heart failure, is of less value in the maintenance therapy of the cardiac sufferer. Its action tends to be transient, and when given by mouth it is often disappointing and has the further disadvantage of being a gastro-intestinal irritant. It has frequently to be discontinued on account of abdominal discomfort and diarrhoea.

Mercurials

The introduction to human investigations of the techniques of determination of kidney function in terms of renal blood flow and glomerular filtration has recently led to a bold reassessment of the part played by the kidney in the whole mechanism of heart failure. Merrill (1946) and his colleagues have shown that the oedematous cardiac patient retains far more sodium than the normal individual. Mercurial diuretics reverse this sodium retention, and for the time being the kidney gets rid of large amounts of salt as well as water.

At the Postgraduate School recently Drs. L. G. C. Pugh and C. Wyndham have made observations on cardiodynamics during the action of mercurials. As most of these mercurial diuretics contain theophylline it was necessary first to eliminate any special effects of this drug. The theophylline part of the injection does in fact have a slight venous-pressure-reducing and cardiac stimulating action, which, however, is transient. When a mercurial is injected without any theophylline diuresis follows in a period of 4 to 10 hours. At the peak of the diuresis the venous pressure is reduced and the cardiac output is raised (Fig. 8). As the diuresis passes off, the venous pressure climbs again towards the value prior to the injection. The action is very much like that of a venesection. This is the first clear demonstration of a circulatory action of mersalyl, and the value of the substance in the treatment of heart failure is further appreciated.

Discussion and Conclusion

There still exists a great deal of confusion about the meaning and significance of venous congestion, and the view is prevalent that venous congestion is a damming back of blood behind a failing pump, and that any improvement in cardiac output will automatically reduce this pressure. Against this view there are many arguments, some of which are: (1) Direct cardiac stimulation by minute doses of adrenaline below the threshold of dosage necessary to influence arterial pressure or heart rate increases the cardiac output by about 50% without affecting the venous pressure (McMichael and Sharpey-Schafer, 1944b). (2) Venous congestion is now abundantly recognized as an accompaniment of numerous circulatory disorders with a high cardiac output. (3) In normally acting hearts cardiac output and venous pressure tend to rise and fall in parallel and not inversely as would be expected on the older view. (4) Drugs like digitalis and theophylline may produce a rise, no change, or a fall in cardiac output; but, whatever the cardiac reaction, the venous pressure nearly always falls. The venous-pressure change is therefore not determined by the cardiac response but rather the reverse.

Thus it is clear that the application of Starling's law, which established venous filling as a primary regulatory mechanism in controlling cardiac output, forms a much more satisfactory working hypothesis for the consideration of the problems of circulatory dynamics in health and disease.

The lessons we have learnt from our investigations should induce a deep sense of humility. We used to believe that we were "scientific" when we subdivided the therapy of heart failure into the use of substances which acted on the heart (digitalis), while diuretics (theophylline and mercurials)

were supposed to act on the kidneys. It now seems that these latter substances also act beneficially, directly or indirectly, on the circulation itself, while digitalis seems to have an important action on venomotor tone, possibly of greater significance than any direct cardiac effect. We must also plead guilty to neglect of the study of the actions of the strophanthins in this country.

The sick man must be studied as a functioning whole and not as a series of isolated organs. The application of direct investigative methods to man further makes it clear that human pharmacology must stand on its own feet as a science yielding results at least as valid as those of the animal laboratory. In the latter, investigations are considerably hampered by the use of material from varying biological species and in no sense guaranteed to reproduce the conditions which obtain in clinical medicine. The story of Chassar Moir's discovery 16 years ago, by direct observations on the human puerperal uterus, of an active principle in ergot previously overlooked in decades of pharmacological work was a lesson we should not readily forget (Moir and Dale, 1932).

Clinical medicine tends to make use of many valuable measures long before their scientific rationale is fully understood. The origin in an old wives' remedy of Withering's introduction of digitalis shows that medicine has a responsibility to give fair trial to new therapeutic ideas even though their origin may be somewhat dubious. The polypharmacists who applied all remedies—digitalis, mercurials, and venesection—simultaneously to the cardiac patient probably did better for their patients than their more scientific fellows who tried to observe the actions of these substances one by one. But the blind haphazard use of drugs is equally to be deplored, and we have a predominant duty to analyse the manner in which each remedy may act.

Finally, I should like to paraphrase very slightly the words of my fellow countryman James Mackenzie (1905):

"We do not for one moment suggest that we have found out the whole matter of the action of digitalis, because there are many phenomena produced by digitalis with which we do not deal. The interpretation here given may not be found sufficient because our knowledge is very limited, and new facts are cropping up continually. . . . I have endeavoured so far as possible to give a careful record of the facts with such interpretations as appear most reasonable in our present state of knowledge."

I gratefully acknowledge the help of my colleagues, Drs. E. P. Sharpey-Schafer, Sheila Howarth, P. Wood, and, more recently, S. Ahmed, R. I. H. Bayliss, W. A. Briscoe, A. J. Palmer, L. G. C. Pugh, and C. Wyndham. The work could not have been carried to its present stage without the technical help of Messrs. A. H. Latham, R. W. Halls, and W. Manders.

REFERENCES

- Anitschkow, S. V., and Trendelenburg, P. (1928). *Dtsch., med. Wschr.*, **54**, 1672.
 Bodo, R. (1928). *J. Physiol.*, **64**, 365.
 Boyer, N. H. (1943). *J. Amer. med. Ass.*, **122**, 306.
 Christian, H. A. (1919). *Amer. J. med. Sci.*, **157**, 593.
 Cohn, A. E. (1915). *J. Amer. med. Ass.*, **65**, 1527.
 Cournaud, A., and Ranges, H. A. (1941). *Proc. Soc. exp. Biol., N.Y.*, **46**, 462.
 Curschmann, H. (1873). *Dtsch. Klin.*, p. 377.
 Cushny, A. R. (1918). *J. Pharmacol.*, **11**, 103.
 Dock, W., and Tainter, M. L. (1930). *J. clin. Invest.*, **8**, 467.
 Forssmann, W. (1929). *Klin. Wschr.*, **8**, 2085.
 Fraenkel, A., and Thauer, R. (1934). *Ergebn. inn. Med.*, **46**, 208.
 Fraser, T. R. (1890). *Trans. R. Soc. Edin.*, **36**, 343.
 Gavey, C. J., and Parkinson, J. (1939). *Brit. Heart J.*, **1**, 27.
 Gremels, H. (1933). *Arch. exp. Path. Pharmacol.*, **169**, 689.
 Harrison, T. R., and Leonard, B. W. (1926). *J. clin. Invest.*, **3**, 1.
 Howarth, S., McMichael, J., and Sharpey-Schafer, E. P. (1946). *Clin. Sci.*, **6**, 41.
 ——— (1947). *Ibid.*, **6**, 125.
 Katz, L. N., Mendlowitz, M., and Kaplan, H. A. (1938). *Amer. Heart J.*, **16**, 149.
 ——— Rodbard, S., Friend, M., and Rottersman, W. (1938). *J. Pharmacol.*, **62**, 1.
 Mackenzie, J. (1905). *British Medical Journal*, **1**, 587.

- McMichael, J., and Sharpey-Schafer, E. P. (1944a). *Quart. J. Med.*, **13**, 123.
 — (1944b). *Brit. Heart J.*, **6**, 33.
 Marais, O. A. S., and McMichael, J. (1937). *Lancet*, **2**, 437.
 Marvin, H. M. (1927). *J. clin. Invest.*, **3**, 521.
 Merrill, A. J. (1946). *Ibid.*, **25**, 389.
 Moir, Chassar, and Dale, H. H. (1932). *British Medical Journal*, **1**, 1119.
 Ray, T., and LaDue, J. S. (1945). *Amer. Heart J.*, **30**, 335.
 Ryland, D. A. (1933). *J. clin. Invest.*, **12**, 847.
 Vaquez, H., and Lutembacher, R. (1928). *Pr. méd.*, **36**, 129.
 Visscher, M. B. (1938). *Minn. Med.*, **21**, 85.
 Withering, W. (1785). *An Account of the Fox-glove and some of its Medical Uses*. Robinson, Birmingham and London.
 Wood, P. (1940). *Brit. Heart J.*, **2**, 132.

PROCAINE PENICILLIN: EFFECT OF SINGLE DAILY INJECTIONS

BY

P. F. JONES, M.B., B.Ch.

Chief Assistant, Surgical Professorial Unit,
St. Bartholomew's Hospital

AND

R. A. SHOOTER, M.D.

Assistant Bacteriologist, St. Bartholomew's Hospital;
Lecturer in Bacteriology, St. Bartholomew's
Hospital Medical College

Among the devices for prolonging the presence of penicillin in the body, suspensions of penicillin in oil and beeswax have until recently been those most commonly used. These suspensions are apt to give rise to local reactions, for which the beeswax is blamed. A more recent preparation, also designed to prolong the presence of penicillin but thought to be free from undesirable complications, is procaine penicillin.

This substance, described by Salivar, Hedger, and Brown (1948) and Sullivan, Symmes, Miller, and Rhodehamel (1948), is a true salt formed by the equimolecular combination of procaine and sodium or potassium penicillin G. It has a solubility in water of 0.43%, and is administered as a suspension in oil—sesame, arachis, and cotton-seed oils all having been used. The use of oil as a vehicle enables the compound to be dispensed ready for use in a stable form, and further prolongs absorption. From the reports of Hobby, Brown, and Patelski (1948) it appears that *in vitro* and in laboratory animals it is little if at all inferior to sodium penicillin G, and the persistence of penicillin in the blood of animals following a single injection, which they describe, has been shown to occur in humans by Herrell, Nicols, and Heilman (1947), by Boger, Oritt, Israel, and Flippin (1948), and by Hewitt, Whittlesey, and Keefer (1948). Following a single injection of 300,000 units of procaine penicillin Herrell *et al.* (1947) found detectable amounts of penicillin in the blood of nine out of ten patients after 24 hours. Boger *et al.* (1948) showed that after an injection of 300,000 units of procaine penicillin in ambulatory patients detectable amounts of penicillin remained in the blood from six to thirty hours, while in patients with pneumonia the range varied from seven to seventy hours. Of 11 cases of pneumonia treated with a single injection of procaine penicillin eight required no further treatment. Hewitt *et al.* (1948) showed that after an injection of 300,000 units of procaine penicillin, out of 63 patients 58 had detectable amounts of penicillin in their blood at twenty-four hours and 21 at forty-eight hours.

Tests in a Series of Cases

This paper presents information derived from the treatment of 45 ambulant patients and four ward patients with procaine penicillin G in arachis oil.

All the ambulant patients received five daily injections of 1 ml. of procaine penicillin in arachis oil which contained 300,000 units of penicillin. No difficulty was experienced in administering the compound so long as, after shaking, it was withdrawn from the bottle with a wide-bore needle, not less than a No. 1 serum needle, and it was recognized that it would take 10 to 15 seconds to withdraw 1 ml. For administration a No. 1 hypodermic needle was satisfactory.

The ambulant patients comprised 45 persons of both sexes attending the casualty department of the hospital, their ages ranging from 17 to 83. Although ambulant in that they attended the hospital daily, the majority of them were unfit for work, and were instructed to rest as much as possible at home. In addition 12 children have been treated with procaine penicillin, the youngest aged 14 days, from whom it was not desirable to obtain blood and who are therefore not included in this report. The diseases for which treatment was given were all inflammatory in nature and included carbuncles, furunculosis, breast abscesses, cellulitis, pulp infections, and infected abrasions with lymphangitis and adenitis. The injections were said to be painless, and no local or general reactions were seen.

Penicillin blood levels were determined by the Oxford slide method, using the Oxford H staphylococcus, immediately before the second injection of penicillin in each patient, and in as many cases as possible further levels were determined before the third, fourth, and fifth injections and at approximately 24 hours after the fifth injection. The results are shown in Charts 1 and 2.

It will be seen that it has been possible to find detectable amounts of penicillin in the blood of 37 out of 45 patients at times varying from 18 to 26 hours after their first injection of 300,000 units of procaine penicillin. Determinations made between 18 and 26 hours after the second or subsequent daily injections have shown detectable amounts of penicillin in 93 out of 109 samples. Neither