STEROID METABOLISM AND FRONTAL LOBES

BY

F. REITMAN, M.D., D.P.M.

Director of Clinical Research, Netherne Hospital, Coulsdon,
Surrey

The 17-ketosteroids are end-products of steroid metabolism, their source being the adrenal cortex. The daily urinary 17-ketosteroid output is fairly constant—3.5–14.6 mg. in 24 hours for adult normal women and 9.4–20 mg. in 24 hours for adult normal men (Patterson, McPhee, and Greenwood, 1942). Hemphill, MacLeod, and Reiss (1942) investigated the 17-ketosteroid values before and after brain operations, in particular bilateral prefrontal leucotomy (lobotomy), in six cases, and found that after the third post-operative month the urinary 17-ketosteroid excretion rises, indicating an increased steroid metabolism. Commenting on their results, they advanced the theory that an interference with the nervous pathways between the frontal lobes and the hypothalamus may influence the anterior pituitary, which in its turn alters the pituitary adreno-gonadal functions.

The cerebral control of steroid metabolism is of clinical as well as physiological importance, and the above-quoted conclusions have been reinvestigated in 12 cases. All the patients were females at the period of sexual maturity, and all had undergone bilateral prefrontal leucotomy, performed with a rotating-blade leucotome of 25 mm. cutting diameter. The urinary 17-ketosteroid output was followed up fortnightly for three to four months, except in the case of one patient who, after leaving the hospital, was inaccessible for the repeated investigations. The pre-operative and 3–4-monthly post-operative values are tabulated below.

Case No.	17-Ketosteroids in mg./24 hours	
	Pre-operative	3–4 Months Post-operative
1	15.1	11.4
2	6.1	4.4
3	14.6	10.9
4	10.0	6.5
5	13.2	—
6	11.0	4.8
7	14.2	4.2
8	10.9	10.5
9	23.6	14.0
10	20.3	19.5
11	12.4	9.6
12	9.3	8.4

The absence of the post-operative value in Case 5 has been explained; Cases 9 and 10 show high pre-operative values, but as the sample was taken at random they are retained in the series. It can be seen that the 17-ketosteroid output taken three to four months post-operatively is lower than that taken pre-operatively, in contradistinction to Hemphill and MacLeod's findings.

Further examination of our figures shows that the standard error of the mean difference is 1.05 mg./24 hours and Student's *t* ratio 3.71. For 10 degrees of freedom this is significant at the 1% level; thus the odds are 99 to 1 against its arising by chance. In other words, statistical evidence suggests very strongly that after bilateral injury to the prefrontal lobes the 17-ketosterone excretion is likely to diminish.

The hypothesis of other workers—namely, that the prefrontal lobes influence steroid metabolism—thus seems to be confirmed, but the direction of quantitative alterations is found to be contradictory. While no concrete explanation can be offered for these differences, attention should be drawn to the fact that the technique of leucotomy

(lobotomy) varies from hospital to hospital, and the extent and localization of the operative incision at various levels may be responsible for the difference in findings. Furthermore, Hemphill and his collaborators' reported cases were males, whereas the case material just presented consisted entirely of females.

Reports on frontal-lobe injuries and on lobotomies emphasize a frequently increased sensuality. In the above-reported case material three patients showed increased sensuality but no corresponding increase in the 17-ketosteroid excretion could be observed—a possible indication that the increased sensuality is not of neuro-hormonal genesis. In other words, the 17-ketosteroid values seem to be independent of the patients' psychosexual state.

To ascertain whether patients with frontal-lobe injury respond to stimulation in the same manner as "normals," 17-ketosteroid values have been studied on patients who had undergone leucotomy and on patients who suffer from a mild neurosis. These patients were stimulated by 8 gr. (0.53 g.) of pituitary whole gland in 24 hours, by 6 mg. of intramuscular "prostigmin" in 24 hours, and finally by the administration of an erythema dose of ultra-violet light on the day the urinary 17-ketosteroids were collected. These experiments are still incomplete and numerically do not lend themselves to statistical examination, but the main tendencies of the results so far are as follows:

Administration of pituitary whole gland raises the 17-ketosteroid excretion, but administration of the prostigmin results in lowered 17-ketosteroid excretion. These findings might confirm a cortico-pituitary control of the 17-ketosteroid excretion rather than an influence through autonomic mechanisms. On administration of an erythema dose of ultra-violet light the 17-ketosteroid excretion increases, as was reported by Myerson and Neustadt (1939). No differences in responses were found between operated and non-operated cases.

I wish to thank Dr. W. T. Griffith and Mr. J. P. S. Robertson for their collaboration and help.

REFERENCES

- Hemphill, R. E., MacLeod, L. D., and Reiss, M. (1942). *J. ment. Sci.*, **88**, 554.
Myerson, A., and Neustadt, R. (1939). *Endocrinology*, **25**, 7.
Patterson, J., McPhee, I. M., and Greenwood, A. W. (1942). *British Medical Journal*, **1**, 35.

Medical Memoranda

Haematoma of the Rectus Abdominis Muscle

Haematoma of the rectus abdominis muscle, especially the variety in which trauma is an inconspicuous factor and which is referred to as spontaneous to distinguish it from the traumatic variety, is very uncommon but of considerable clinical interest because it simulates very closely an acute intraperitoneal lesion.

CASE REPORT

A widow aged 71 was admitted to hospital on Aug. 3, 1947, complaining of severe abdominal pain of ten hours' duration. There was sudden onset of "tearing" pain in the right lower abdomen. The pain was continuous, aggravated by movement, and became steadily worse up to the time of admission. She vomited twice soon after the onset of pain. The bowels were opened before the attack and flatus had passed after the attack came on. Micturition was normal. There was a previous history of chronic bronchitis.

On examination the temperature was 99.2° F. (37.3° C.), pulse rate 120 and of poor quality, and respiration normal but somewhat laboured. The facies was pale and anxious, and the tongue moist and clean. The general appearance was one of moderate shock. Inspection revealed an ovoid swelling of the right lower abdomen. Abdominal respiratory movements were limited. The shoulder-raising

test was too painful to carry out properly. Palpation confirmed that the swelling extended from the symphysis to the umbilicus, and from the midline to the right as far as the linea semilunaris. The shape was oval, with the large diameter in the vertical axis. The swelling was smooth, firm, and tense, without impulse, extremely tender, and did not move with respiration. Percussion of the swelling gave a dull note. Rectal and vaginal examinations and urinalysis were negative.

The case resembled an acute surgical abdomen, and torsion of an ovarian cyst or torsion of the omentum was a possible diagnosis. As mentioned before, the shoulder-raising test was too painful to carry out and no deduction was made whether the swelling was in the abdominal wall or was intraperitoneal, but it seemed more superficial than would have been expected of an intraperitoneal lesion. However, operation was decided upon mainly because a serious intraperitoneal lesion could not be discounted.

Operation.—A right sub-umbilical paramedian incision over the swelling revealed a bulging and dusky-coloured rectus sheath. Dark, tarry blood clot exuded immediately the sheath was incised. A clot the size of a large grape-fruit was removed from in front of and behind a longitudinal split in the rectus muscle which extended from umbilicus to symphysis. There was considerable bleeding welling up from behind the rectus muscle in the region of the deep epigastric vessels, and this was only partially controlled by under-running sutures, so that packing was necessary. The wound was closed on either side of the pack.

The pack controlled the bleeding well and was removed after 36 hours. A full course of penicillin was administered. The wound healed satisfactorily and the patient was discharged 21 days later. The follow-up revealed no obvious abnormality of the rectus muscle, which functioned very well.

DISCUSSION

That trauma in some degree or other is the exciting factor in all cases appears certain. But where this traumatic factor is slight—e.g., coughing (the factor in this case)—then there will be a predisposing factor, and the main ones seem to be (1) senility, with its degenerative changes in the walls of blood vessels; (2) debilitating illness, with its degenerative changes in muscle; and (3) pregnancy, with its effect of stretching the muscle fibres of the rectus abdominis and the deep epigastric blood vessels.

Correct diagnosis is seldom made, the most popular one being torsion of an ovarian cyst. Diagnostic points are a history of paroxysm of coughing followed by pain, with the early appearance of a mass which may steadily increase in size and which becomes more prominent and fixed on raising the shoulders. Operative treatment is usually undertaken because of the doubt in diagnosis, but where a definite diagnosis of haematoma of rectus muscle is made then there is the choice of conservative treatment. If the haemorrhage is already considerable or is progressing operation should be undertaken; Teske (1946) gives an overall mortality of 4%, and Thomas (1945) states that there is a 13% mortality rate in the obstetric group.

G. H. MOORE, F.R.C.S. Ed.,
Surgeon, North Lonsdale Hospital,
Barrow-in-Furness.

REFERENCES

- Teske, J. M. (1946). *Amer. J. Surg.*, **71**, 689.
Thomas, R. C. (1945). *J. Obstet. Gynaec. Brit. Emp.*, **52**, 580.

A Case of Melaena Complicating Congenital Pyloric Stenosis

The following case of congenital hypertrophic pyloric stenosis is worthy of record on account of an unusual complication—melaena—occurring after operation.

CASE REPORT

A male infant aged 6 weeks, an only child, was admitted to the Royal Salop Infirmary on July 16, 1947. He had had projectile vomiting for over a week. He had been breast-fed, but his mother had given him complementary feeds of half-cream milk food after each feed for a few days. The weight had risen from 7 lb. 15 oz. (3.6 kg.) to 9 lb. 6 oz. (4.25 kg.) before the onset of vomiting, but had fallen to 7 lb. 4 oz. (3.29 kg.) on admission.

Congenital pyloric stenosis was diagnosed and operation was decided upon (July 21). The stomach was first washed out, and then under local analgesia a Rammstedt's operation was performed. There was a well-marked pyloric tumour. In separating the circular muscle

fibre the duodenal mucosa was inadvertently opened and was sutured with fine catgut. In view of this perforation an intravenous saline drip was set up and feeding by mouth withheld until July 23, when half-strength half-cream milk food was given, the drip being discontinued on July 24. Apart from two small vomits he made good progress until Aug. 1, when his weight was 8 lb. 3½ oz. (3.73 kg.). He then started vomiting bile-stained material, and this lasted two days, being relieved after gastric lavage and rectal saline. Feeding by mouth was resumed.

A temporary improvement occurred, but on Aug. 5 vomiting recommenced and the stools were relaxed. There had been no pyrexia throughout, and no signs of peritonitis. On Aug. 6 the stomach was washed out and rectal saline given. A course of phthalylsulphathiazole, one tablet four-hourly, was started. Next day he was given glucose feeds. A "dark brown" stool was reported. His weight was 7 lb. 8 oz. (3.4 kg.).

On Aug. 8 (the 18th day) the baby was collapsed, pale, and generally oedematous. A plasma transfusion was started. Soon afterwards a large melaena stool was passed. The haemoglobin was 30%. Blood was substituted for the plasma, and about 450 ml. run in slowly. Next day a remarkable improvement had occurred and the oedema had disappeared. The haemoglobin was 92%. Phthalylsulphathiazole was discontinued. After several dark stools the motions had now become a normal colour. Feeding was resumed with half-cream milk food. No further vomiting occurred. Apart from one melaena stool on Aug. 10 the motions remained normal. Unfortunately it was not possible to estimate the prothrombin index.

Further convalescence was uninterrupted, and the baby was discharged on Aug. 26 weighing 8 lb. 5 oz. (3.77 kg.). When seen again on Oct. 7, as an out-patient, his weight was 13 lb. 14 oz. (6.29 kg.) and his mother stated that he had remained well.

DISCUSSION

I have found only a few references to bleeding into the bowel in cases of pyloric stenosis. Meader (1936), quoting a case in which "occult blood" had been found in the stools on the seventh day—gradually decreasing after a few days—commented: "Now it is not unusual for these patients to vomit fresh blood within 24 hours following operation, or for some evidence of blood to appear in the first or second stool, but in our experience this phenomenon has not occurred as late as in this instance."

In my case the bleeding was severe, occurring on the 18th day, and was preceded by gastro-enteritis. It was presumably due to ulcer. Infection was probably an aetiological factor. Possibly trauma, particularly as the duodenal mucosa was punctured at operation and subsequently sutured, determined the location of the ulcer.

If infection and trauma are regarded as aetiological factors in peptic ulcer (Illingworth and Dick, 1941) it is surprising that this complication is not more common in pyloric stenosis, since infections of various kinds are not infrequent after operation and the mucous membrane is widely exposed after incising the seromuscular layers.

The importance of breast-feeding in pyloric stenosis in relation to prognosis is well established (Herzfeld and Wallace, 1935; Thompson and Gaisford, 1935; Tallerman, 1938; Levi, 1941; Williams, 1942). Levi (1941) reported a consecutive series of 100 breast-fed babies who underwent a Rammstedt's operation without a death, and contrasted this with a series of 46 babies who were bottle-fed, of whom five died—all from gastro-enteritis. Bottle-fed babies present greater risk of cross-infection owing to their longer stay in hospital; they are less well nourished and are more liable to suffer from dietetic disorders (Williams, 1942).

It is probably significant that my case was artificially fed immediately before and after operation.

I wish to thank Dr. A. D. Rope and Mr. A. D. Haydon, under whose care this case was admitted, for the opportunity of treating it.

H. C. FLETCHER-JONES, M.B., B.S., F.R.C.S.,
Late Resident Surgical Officer,
Royal Salop Infirmary.

REFERENCES

- Herzfeld, G., and Wallace, H. L. (1935). *Lancet*, **2**, 385.
Illingworth, C. F. W. and Dick, B. M. (1941). *A Textbook of Surgical Pathology*, 4th ed. Churchill, London.
Levi, D. (1941). *British Medical Journal*, **1**, 963.
Meader, H. B. (1936). *Ohio St. med. J.*, **32**, 531.
Thompson, W. A., and Gaisford, W. F. (1935). *British Medical Journal*, **2**, 1037.
Tallerman, K. H. (1938). *J. Pediatr.*, **13**, 787.
Williams, H. (1942). *Med. J. Aust.*, **1**, 303.