ENDOCRINES IN THEORY AND PRACTICE

This article is one of a series on Endocrinology contributed by invitation

HYPERPARATHYROIDISM

BY

DONALD HUNTER, M.D., F.R.C.P.

Until twelve years ago we had no proof of the relationship between parathyroid activity and the metabolism of phosphorus and calcium. The discovery of the parathyroid hormone led at once to studies of its effects in normal individuals. When injections of an active parathyroid extract are given over a long period of time they cause a decrease in the plasma phosphorus, an increase in the serum calcium, and at the same time a mobilization of calcium salts from the skeleton, which are then passed in the urine. This knowledge has led to the discovery of cases of a bone disease associated with hyperfunction of a parathyroid tumour—the generalized osteitis fibrosa of von Recklinghausen.

In 1891, in a Festschrift in honour of Virchow's seventy-first birthday, von Recklinghausen gave the first accurate account of the bone disease which now goes by his name. Although his contribution is involved and he described more than one bone disease, there can be no doubt that his Cases 5 and 7 constitute the discovery of generalized osteitis fibrosa. Thirteen years later Askanazy reported a case associated with a parathyroid tumour, but no special significance seems to have been attached to the association, despite the fact that Erdheim had already noted enlargement of the parathyroid glands in osteomalacia and had suggested that they were functionally related to calcification. When opinion differed on whether the increase in parathyroid secretion was primary or secondary to the bone disease, Erdheim himself defended the view that the skeletal change was primary and that the parathyroids enlarged merely to meet the greater demand for calcium. Acting on this assumption, Mandl in 1925 transplanted four parathyroid bodies into the abdominal wall of a man with generalized osteitis fibrosa. No improvement followed, however, and he accordingly explored the neck, where he found a parathyroid tumour measuring 2.5 by 1.5 by 1.2 cm. This was removed, and the subsequent disappearance of pain in the bones, the reduction of excretion of calcium in the urine, and the general improvement in the patient's health constituted the first clinical evidence that in this disease the parathyroid lesion is primary and the skeletal changes secondary. By this time it was well established that the parathyroid glands are specifically concerned in regulating the metabolism of calcium.

The isolation of an active parathyroid extract which raises the calcium of blood and urine, and the recognition that injection of this extract causes loss of calcium from the bones, were the next important steps in deciding that in generalized osteitis fibrosa the high calcium content of the serum, the increased calcium excretion in the urine, and the diminished density of x-ray shadows of the bones, point with certainty to a parathyroid lesion.

Morbid Anatomy

Hyperparathyroidism leads to changes which are generalized throughout the skeleton: the bones are soft and cut with a knife like rotten wood. Both cortex and spongiosa are too thin and are largely replaced by tough, grey, fibrous tissue. Multiple grey, spongy, osseous swellings

occur, and sometimes expand the cortex. Firm, round, red tumours and single or multilocular cysts may be present. The great resorption of bone may lead to deformity in any part of the skeleton, and also to spontaneous fractures. Histologically the main changes are lacunar resorption, apposition, fibrosis of marrow, and the formation of osteoclastomata (giant-celled tumours) and cysts. Lacunar resorption is the predominant change. There is a general osteoporosis which varies in degree in different cases. Apposition, however, does not cease, and where present it is usually abnormally active.

The condition found in the parathyroid glands may be a hyperfunctioning adenoma of a single gland or rarely of part of two glands. The greatest diameter of such tumours may be from 0.7 cm. to 7.5 cm. In some cases there is a generalized hyperplasia of all the parathyroid glands, and it is possible that this is a result of an increased secretion of the parathyrotropic hormone of the pituitary gland. Bilateral renal calculi, composed principally of calcium phosphate, are commonly present. In advanced cases areas of metastatic calcification have been found in the lungs, stomach, kidneys, and myocardium.

It is possible by injecting parathyroid extract into puppies in increasing amounts over a long period to produce hyperparathyroidism experimentally. Such injections lead to progressive resorption of bone, fibrous replacement of the marrow, and to the production of other features characteristic of osteitis fibrosa. Prolonged overdosage may give rise to metastatic calcification in the tissues.

Clinical Features

Besides pain and tenderness of the bones, and even spontaneous fractures, hyperparathyroidism gives rise to symptoms and signs outside the skeleton. These include hypotonia and muscular weakness; anorexia, sometimes with nausea, vomiting, and abdominal cramps; polydipsia and polyuria; renal calculi, sometimes with colic and haematuria; and wasting in advanced cases. There is no one symptom which is diagnostic. Because of the insidious onset and the extremely varied symptomatology the disease is often not suspected. Anorexia, nausea, vomiting, lassitude, and hypotonia may all arise early, and are sometimes presenting symptoms. Polyuria and polydipsia are the rule, and may be so marked as to suggest diabetes insipidus. Increased frequency of micturition during the night may be a distressing feature, and occasionally there is enuresis. The large quantity of water necessary for the increased excretion of calcium salts is of course responsible for this group of symptoms.

Hyperparathyroidism is a progressive disease, and if untreated it is usually fatal. It is more than twice as common in women as in men and it occurs at all ages, but most commonly in middle life. The youngest case known is in a boy of 10 and the oldest in a woman of 69. It is possible to describe several different types of hyperparathyroidism, for the clinical features vary according to whether the skeletal, renal, or gastrointestinal symptoms predominate.

CLASSICAL TYPE WITH FIBROCYSTIC SKELETON

In this type the skeletal symptoms and signs predominate. They consist of pain, bone deformities, generalized decalcification, and sometimes fractures. Though this type may begin with weakness, thirst, and polyuria, most

patients complain early of pains in the bones, particularly in the back, pelvis, and legs. Deformities of the skeleton appear and slowly progress. They are of the type commonly seen in conditions associated with softening of the skeleton. The thoracic cage and pelvis are often grossly deformed, and the spine is shortened owing to narrowing of the vertebral bodies. Kyphosis occurs, with loss of several inches in height. The chest has a deep antero-posterior diameter and distorted ribs, and the abdomen is shortened. The limbs may be irregularly curved and deformed, so that walking becomes at first difficult and then impossible. Occasionally bone tumours or cysts are visible or palpable, especially when they appear in the limbs or jaws. The fingers may show a characteristic deformity, in which the terminal phalanges become shorter and broader than the normal. The appetite fails, there are nausea and vomiting, and the patient becomes extremely weak and may go on to a cachectic state with extreme deformity and helplessness. Treatment at this stage of the disease will not restore the normal function of the skeleton, and it should be our aim to make the diagnosis of hyperparathyroidism before such complications have arisen.

TYPE WITH OSTEOPOROTIC SKELETON

The skeletal changes in this type may be indistinguishable from those of osteomalacia, hyperthyroidism, or senile osteoporosis. In all such cases studies of the blood chemistry and the calcium balance are essential.

TYPE WITH RENAL CALCULI

The presenting symptom in this type may be renal colic or haematuria. Skeletal changes may be absent, minimal, or severe. In all patients with renal calculi, especially where they are bilateral, the serum calcium and plasma phosphorus should be estimated in order to rule out hyperparathyroidism.

TYPE WITH METASTATIC CALCIFICATION OF THE KIDNEYS

In this type calcium phosphate is deposited in the substance of the kidney, and can be demonstrated in radiographs. Chronic nephritis occurs, and as it progresses the plasma phosphorus may rise and the increased excretion of calcium salts may be diverted from the urine to the faeces. Patients with this type of renal lesion may be wrongly diagnosed as cases of chronic nephritis.

Radiological Features

Radiologically the most striking feature is the great lack of calcium salts throughout the skeleton. The cortex of the bone loses its ivory character and may be reduced to a thin, uneven, linear shadow. The medulla becomes less dense, with resulting accentuation of the bone pattern at first, but later cancellous bone tends to fade entirely. In the skull there is a uniform miliary granular mottling with small translucent cyst-like areas. The vertebrae may become biconcave like fish vertebrae, and they are then reduced in height. There may be evidence of cysts, but these only rarely expand the corticalis. At times the cysts are difficult to see, so great is the lack of calcium in the surrounding bone. The cystic changes come on late in the disease, and their presence is not essential to diagnosis. All the bones of the body may be affected, even the phalanges of the hands and feet. Areas of subperiosteal resorption in the long bones and phalanges are common. The pelvis undergoes pressure deformities due to the thrust of the femoral heads: cyst-like areas are common in the pelvic bones. The femora and ribs may bend and collapse.

In estimating the density of the radiographic shadows of bones it is found necessary to use a controlled method. A control subject is chosen of the same sex, age, weight, height, and build as the patient. The corresponding limbs of patient and control are placed side by side and exposed simultaneously on the same negative. Where a series of radiographs is taken in one patient the same control subject is used each time. The urinary tract should always be examined for calculi. Where metastatic calcification has occurred in the kidneys it will show in radiographs. On examination of the chest and neck it must be borne in mind that a parathyroid tumour may vary greatly in size and position, and that its shadow may be found in the mediastinum.

Calcium and Phosphorus Metabolism

Increased production of the parathyroid hormone leads to characteristic changes in the calcium and phosphorus metabolism. The serum calcium is raised, and the figures vary between 12.6 and 23.6 mg. per 100 c.cm. Values as high as 14 mg. per 100 c.cm. are not uncommon. Any serum calcium figure of 11 mg. per 100 c.cm. should be regarded as suspicious, and therefore be repeated. The plasma phosphorus figure is low, and may vary between 1 and 2.7 mg. per 100 c.cm. The combination of this high calcium and low phosphorus figure almost always indicates hyperparathyroidism, but in cases in which there is renal impairment the plasma phosphorus figure may become elevated. The plasma phosphatase figure is raised in all cases of hyperparathyroidism in which bone disease can be detected. There are cases without detectable bone disease in which the plasma phosphatase is normal.

An increased calcium excretion in the urine is always present, and varies from slightly above the normal to eight times the normal figure. In an uncomplicated case the increased calcium excretion is entirely urinary, the faecal excretion being unaffected. In individuals with renal impairment, however, part of the increased excretion is faecal. There is also an increased urinary phosphorus excretion.

Differential Diagnosis

If the history and physical signs are suggestive of hyperparathyroidism complete confirmation of the diagnosis may usually be obtained in the chemical findings—namely, a high serum calcium, low plasma phosphorus, and increased calcium excretion in the urine. In more difficult cases removal of a portion of bone for histological section may be necessary. Without such chemical or histological confirmatory evidence it is not justifiable to explore the parathyroid glands.

The following skeletal diseases are sometimes confused with hyperparathyroidism: osteomalacia, senile osteoporosis, thyrotoxic osteoporosis, osteitis deformans, osteoporosis with renal glycosuria, secondary carcinomatosis of the skeleton, multiple myelomatosis, osteogenesis imperfecta, renal rickets, focal osteitis fibrosa, dyschondroplasia with multiple enchondromata (Ollier's disease), xanthomatosis of bones (Schüller-Christian's disease), and Gaucher's disease.

Treatment

By means of a high calcium and high phosphorus diet it is possible to demonstrate improvement in skeletal calcification, but such treatment is contraindicated because it increases the danger of renal complications. No beneficial clinical or metabolic results have been observed in the treatment of hyperparathyroidism by x-irradiation. The treatment is surgical. The parathyroid tumour itself has been palpable in less than 20 per cent. of cases.

Even the largest tumour removed, which measured 7.5 by 5 by 1.8 cm. and weighed 26.2 grammes, could not be felt in the neck as it was behind the trachea. The patient should be prepared for operation by giving for some weeks a high calcium diet and vitamin D. The neck should be explored through a collar incision and the four sites of the parathyroids examined. The chief operative difficulty is that of finding the tumour. Search must be made from the bifurcation of the carotids to the arch of the aorta. If a single parathyroid tumour is discovered it should be excised, but if two tumours are found there is some risk in removing both. In at least six cases the tumour has been removed from the mediastinum. In eight cases two parathyroid tumours were removed; five of the patients survived this drastic treatment and the other three died in tetany. In three cases subtotal parathyroidectomy was performed, for the good reason that in each of them two normal parathyroid bodies had already been excised.

When no tumour at all has been discovered removal of one or two normal parathyroid glands has had no permanent good effect. Such an operation does not in fact seem to be permissible, for in every case of hyperparathyroidism yet recorded there has been a tumour measuring at least 0.7 cm. in its greatest diameter. Moreover, 30 per cent. of the deaths recorded after operation have been due directly or indirectly to interference with one or more normal parathyroid bodies. In less than 4 per cent. of recorded cases has death immediately followed operation, but in 10 per cent. tetany or a renal complication killed the patient at periods varying from nineteen days to fourteen months later. Even if we allow that unsuccessful ventures do not always find their way into print the operative mortality is not high, and this in spite of the fact that the technical problems confronting the surgeon may be complicated. Thus in 1931 Walton reported a case in which search in the usual sites of parathyroid glands did not reveal a tumour, and a small mass in the lower part of the thyroid gland was removed in the belief that it might be a parathyroid tumour. However, on section this proved to be thyroid tissue, and since its removal was not attended by any material chemical changes in the blood Walton opened the wound five days later. He then incised a deep layer of the cervical fascia which forms the bed of the thyroid gland, and two parathyroid tumours were found and removed. One tumour was lying behind the oesophagus, and the second—and larger—retrosternal tumour was lying in front of the second and third thoracic vertebrae. Since learning of this experience others have realized that previous unsuccessful explorations have been incomplete. In one such case seven exploratory operations were performed in the course of six years, the tumour being found ultimately only by dividing the sternum (Churchill and Cope, 1934). The ordinary post-operative treatment should include a high calcium diet, vitamin D, and ultra-violet irradiation. Rarely, in cases in which giant-celled tumours have destroyed a bone, amputation is necessary. Operation upon the kidneys may be required for renal calculi.

EFFECTS OF TREATMENT

Removal of a parathyroid tumour brings about dramatic changes. Many crippled patients have been able to dispense with sticks and crutches. Usually, for some reason not yet clear, pain in the bones goes away immediately. Spontaneous fractures often heal as readily as cases of fractures in healthy bones. Decrease in size of osteoclastic tumours of bone within a few weeks of operation on the neck has now been many times recorded. The

effect of treatment in increasing the density of bone is really remarkable, and controlled radiographs have demonstrated this point now many times. There is marked gain in weight. It is interesting to note that the symptomatic changes bear no relation to the size of the tumour. Thus a woman of 37 after four years of total crippledness, including great bone deformities and destruction of hip-joints and shoulder-joints, had a tumour weighing only 1.3 grammes removed, and is now able to walk with sticks and to sew and knit.

As a rule the level of the serum calcium and plasma phosphorus and the excretion of calcium and phosphorus in the urine are restored to normal. Sometimes, indeed, there is temporary hypocalcaemia and even oliguria. Latent tetany after operation has been common, and manifest tetany has often occurred—followed by death in four cases. In one published case tremor and acute mania came on three weeks after operation, at a time when the serum calcium had dropped to 6.6 mg. per 100 c.cm., but the condition was completely relieved in three days by giving injections of the parathyroid hormone in conjunction with a preparation of calcium administered intravenously. This must undoubtedly be regarded as the essential treatment of severe post-operative tetany after removal of a parathyroid tumour.

BIBLIOGRAPHY

- Bauer, W. (1933). *J. Bone Jt. Surg.*, **15**, 135.
Churchill, E. D., and Cope, O. (1934). *Surg. Gynec. Obstet.*, **58**, 255.
Hunter, D., and Turnbull, H. M. (1931-2). *Brit. J. Surg.*, **19**, 203.
Walton, A. J. (1931-2). *Ibid.*, **19**, 285.

VOLUNTARY HOSPITALS: A FEDERATION SCHEME

RECOMMENDATIONS OF SANKEY COMMISSION *

Nearly two years ago at its annual conference the British Hospitals Association resolved to appoint a Commission to consider the present position of voluntary hospitals and how far, in view of legislative and social developments, steps should be taken to promote their interests and safeguard their future. A commission of ten members was set up under the chairmanship of Viscount Sankey. The members of the medical profession on the Commission were Sir Henry Brackenbury, Professor L. S. Dudgeon, Colonel D. J. Mackintosh, and Provost A. W. Sheen, and the others were Miss Bartleet, an alderman of the Birmingham City Council, Mr. Alan Davies, a former mayor of Wolverhampton, Mr. H. L. H. Hill, a past-president of the Institute of Chartered Accountants, Miss E. M. Musson, chairman of the General Nursing Council, and Sir Reginald Poole, a past-president of the Law Society. Mr. R. H. P. Orde, secretary of the British Hospitals Association, acted as honorary secretary.

The Commission began its work in January, 1936, and it was intimated that a report might be expected by the following June. The fact that the Commission has taken fifteen months over its report indicates the complexity of its task. The hearing of evidence occupied twenty-one sittings. The Royal Colleges, the British Medical Association (through Dr. Peter Macdonald, chairman of the Hospitals Committee, and Dr. Charles Hill, Deputy Medical Secretary), the Medical Superintendents' Society, the

* *Report of the Voluntary Hospitals Commission.* British Hospitals Association, 12, Grosvenor Crescent, London, S.W.1. (1s. net, 1s. 2d. post free.)